



Correspondence

Notes on Early Land Plants Today. 1. A new name for *Cryptothallus hirsutus*

LARS SÖDERSTRÖM¹, ANDERS HAGBORG² & MATT VON KONRAT²

¹Department of Biology, Norwegian University of Science and Technology, N-7491 Trondheim, Norway; lars.soderstrom@bio.ntnu.no

²Department of Botany, The Field Museum, 1400 South Lake Shore Drive, Chicago, IL 60605–2496, USA; hagborg@pobox.com, mvonkonrat@fieldmuseum.org

Cryptothallus mirabilis von Malmberg (1933), which is the type species for *Cryptothallus* Malmberg (1933), has been shown to belong to *Aneura* Dumortier (1822; Wickett & Goffinet 2008). One more species (*Cryptothallus hirsutus*) needs to be transferred. It needs a new name, however, since *Aneura hirsuta* already exists.

***Aneura crumii* L.Söderstr., A.Hagborg et von Konrat nom. nov.** Replaced name:—*Cryptothallus hirsutus* H.A.Crum, *Bryologist* 99: 438, 1996 (Crum & Bruce 1996). Holotype:—COSTA RICA. Cerro de la Muerte, growing underground among shrubby bamboos at 3100 m elev., Bruce s. n. (MICH). Blocking name:—*Aneura hirsuta* Furuki, *J. Hattori Bot. Lab.* 70: 317, 1991 (Furuki 1991).

References

- Crum, H.A. & Bruce, J. (1996) A new species of *Cryptothallus* from Costa Rica. *The Bryologist* 99: 433–438.
Dumortier, B.C. (1822) *Commentationes botanicae*. Tournay, 118 pp.
Furuki, T. (1991) A taxonomical revision of the Aneuraceae (Hepaticae) of Japan. *Journal of the Hattori Botanical Laboratory* 70: 293–397.
Malmberg, S. (1933) *Cryptothallus* nov. gen. Ein saprophytisches Lebermoos. *Annales Bryologici* 6: 122–123.
Wickett, N.J. & Goffinet, B. (2008) Origin and relationships of the myco-heterotrophic liverwort *Cryptothallus mirabilis* Malmb. (Metzgeriales, Marchantiophyta). *Journal of the Linnean Society, Botany* 156: 1–12.