



Smilax bolavenensis, a new species of Smilacaceae from southern Laos

SHUICHIRO TAGANE^{1,4,*}, PHETLASYS SOULADETH^{2,5} & MINORU N. TAMURA^{3,6}

¹Kagoshima University Museum, Kagoshima University, 1-21-30, Korimoto, Kagoshima 890-0065, Japan

²Faculty of Forest Science, National University of Laos, Dongdok Campus, Xaythany District, Vientiane Capital, Laos

³Department of Botany, Graduate School of Science, Kyoto University, Kitashirakawa-oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan

⁴✉ stagane29@gmail.com; <https://orcid.org/0000-0002-1974-7329>

⁵✉ p.souladeth@nuol.edu.la; <https://orcid.org/0000-0001-8563-1773>

⁶✉ mntamura@sys.bot.kyoto-u.ac.jp; <https://orcid.org/0000-0002-5774-3353>

*Corresponding author

Abstract

A new species of *Smilax*, *S. bolavenensis* (Smilacaceae), is described from the Bolaven Plateau, southern Laos. It is similar to *S. elegantissima* in its elongated inflorescence with slightly zigzagged rachis longer than 20 cm, but the new species is distinguished by its terete stem and petiole (vs. 4-angled with each corner narrowly winged in *S. elegantissima*) and 42–54-flowered umbels (vs. 7–12). A description, photographs and preliminary conservation status are also provided for *S. bolavenensis*.

Keywords: Bolaven Plateau, Liliales, Southeast Asian flora

Introduction

Smilax Linnaeus (1753: 1028) became the sole genus of Smilacaceae after *Heterosmilax* Kunth (1850: 270) was reduced to *Smilax* (Cameron & Fu 2006, Qi *et al.* 2013a, b). It consists of about 300 species, collectively distributed in tropical, subtropical and temperate regions (Chen & Koyama 2000, POWO 2022). The species are dioecious, often woody climbers and shrubs, sometimes herbs, with simple lamina with 3–7 main veins arising from the base, netted leaf venation, axillary inflorescences are composed of 1(–3) umbel(s) or paniculately, racemosely or spicately disposed several umbels, 6 tepals, superior ovary and drupes or berries (Chen & Koyama 2000, Qi *et al.* 2013a). In Laos, 19 species have been recorded (Gagnepain 1934, Koyama 1983, Newman *et al.* 2017 onwards, Kladwong *et al.* 2018, 2022).

The Bolaven Plateau, southern Laos, is one of the biodiversity hotspots in Indochina (Profile 2012). Indeed, our recent extensive fieldwork carried out in the area 2018–2020 has resulted in discovery of more than 17 new taxa endemic to the Bolaven Plateau and 70 new records for Laos (e.g., Souladeth *et al.* 2020, Tagane *et al.* 2020, Phonepaseuth *et al.* 2021, Souvannakhommane *et al.* 2021, Vongthavone *et al.* 2021, Yamazaki *et al.* 2021, Noyori *et al.* 2022). During our botanical inventories, we collected seven species of *Smilax*, of which six were identified as *S. bockii* Warburg ex Diels (1900: 252), *S. corbularia* Kunth (1850: 262), *S. extensa* Wallich in Hooker (1892: 309), *S. glabra* Roxburgh (1832: 792), *S. lanceifolia* Roxburgh (1832: 792) and *S. megacarpa* Candolle (1878: 186). However, one species did not match descriptions of species recorded in Laos and surrounding countries.

The two unknown specimens were collected during our field surveys in Bolaven Plateau in 2018 and 2019. We examined the putative new species and referred to herbarium specimens at FOF and KAG and digital specimen images available online (e.g. JSTOR Global Plants <https://plants.jstor.org>). Floral measurements were made on rehydrated specimens, and we consulted the relevant literature (Gagnepain 1934a, Koyama 1975, 1983, Hô 2000, Li *et al.* 2011, Heckroth *et al.* 2014, Sun *et al.* 2015, Jin *et al.* 2016, Qi *et al.* 2016, Newman *et al.* 2017 onwards, Baruah *et al.* 2018, Kladwong *et al.* 2018, 2020, Feng *et al.* 2022). Based on these studies, it was clear this is an undescribed species, which we formally described below.

Taxonomy

Smilax bolavenensis Tagane & Soulad., *sp. nov.* (Figs 1, 2)

TYPE:—LAOS. Champasak Province: Paksong District, near Nong Luang Village, Dong Hua Sao National Park (Bolaven Plateau), 15°04'44.53"N, 106°12'27.36"E, 1249 m a.s.l., 17 Dec 2019, *Souladeth et al.* L3342 (holotype: FOF 0006673; isotypes BKF, KAG 155707).

Smilax bolavenensis is similar to *S. elegantissima* Gagnepain (1934b: 619) from Vietnam in having an elongate inflorescence with slightly zigzag rachis longer than 20 cm, but clearly distinguished from it by its terete stem and petiole (vs. 4-angled with each corner narrowly winged) and 42–54-flowered umbels (vs. 7–12).



FIGURE 1. *Smilax bolavenensis* (holotype, *Souladeth et al.* L3342, FOF0006673).




FIGURE 2. *Smilax bolavenensis*. A. Leafy branch. B. Branch with staminate inflorescence. C. Abaxial leaf surface. D., E. Staminate inflorescence. F. Longitudinal section of a staminate umbel. G. Bracteole. H. Staminate flower bud with bracteoles (left), staminate flower bud (middle), and staminate flower bud outer three tepals and one inner tepal removed showing stamens (right). I. Outer tepals (top) and inner tepals (bottom), adaxial side (left), abaxial side (middle and right). A–C from Tagane *et al.* L1964 (KAG); D–G from Souladeth *et al.* L3342 (KAG). Scale bars: F = 5 mm, G–I = 2 mm. Photographed by S. Tagane.

Climbers to 4 m tall, all parts glabrous. Stems and branches subterete, without prickles and wings. Petioles 3.5–4.5 cm long, subterete; sheaths 1.2–1.7 × 0.7–1 cm. Tendrils up to 10.0 cm long, 1.0–1.5 cm above from the base of sheath. Lamina ovate-oblong to narrowly ovate, (11.0–)15.2–22.4 × (3.5–)5.8–13.6 cm, shiny brownish red to brownish yellowish on both surfaces when dry or paler and glaucous abaxially, apex acuminate, acumen up to 0.6 cm long, base cordate, 5-veined, major veins diverging at base of lamina, prominent abaxially, (0.9–)1.7–2.5 cm interval (at widest point of lamina), connected with reticulate veinlets. Staminate inflorescence composed of spicately disposed umbels, to 33 cm long, 2–4-branched, rachis slightly zigzagged, not winged; umbels sessile, 22 per inflorescence, 42–54-flowered, prophylls ovate-lanceolate, 3.0–4.5 × 1.0–1.2 mm, apex acuminate; receptacle of an umbel globose, 2.8 mm

in diam.; bracteoles 4–5, ovate-triangular to oblong-obovate, 1.6–2.5 × 0.6–1.2 mm, apex acute. Staminate flower (before anthesis), tepals 6, free from base, outer ones ovate-oblong, 2.5–2.9 × 1.3–1.5 mm, inner ones narrowly ovate, 2.0–2.2 × 0.9–1.0 mm long, apex acute, stamens 6, filaments sessile, anthers oblong, 1.3 mm long; pedicels 0.3–0.7 mm long. Pistillate inflorescence, fruits and seeds not seen.

Distribution:—Thus far known only from the Bolaven Plateau.

Habitat:—Montane forests, 1240–1250 m a.s.l., dominated by Fagaceae, Lauraceae and Myrtaceae, along with some subcanopy trees such as *Taberaemontana bovina* (Apocynaceae), *Aporosa yunnanensis* (Phyllanthaceae), and *Camellia bolavenensis* (Theaceae).

Phenology:—Staminate inflorescence with flower buds collected in December.

Etymology:—Referring to the type locality, Bolaven Plateau.

Vernacular name:—*Kheau kheaug Bolaven* (ເຄືອເຂືອງບໍລະເວນ; Lao), suggested here. *Kheau kheaug* in Lao refers to a common name for *Smilax*.

Preliminary conservation assessment:—Critically endangered (CR). During our five intensive field surveys in Bolaven Plateau 2018–2020, we found only one population with a few individuals in lower montane forest on the top of a plateau where we collected the above two specimens. Therefore, the most appropriate initial assessment of the IUCN conservation status for this species is critically endangered (CR) under criteria D (IUCN 2012). In the Bolaven Plateau, the forests at higher elevation have been decreasing due to dam construction and coffee plantations, which may produce negative effects for *S. bolavenensis*. Further information on its distribution and number of individuals/populations is needed to accurately assess its status and conserve this species.

Note:—Among the *Smilax* species in Laos, *S. bolavenensis* resembles *S. glabra* in leaf texture and venation patterns, and both species occur in Bolaven Plateau (specimens of *S. glabra*: *Souladeth et al. L3754*, FOF, KAG). However, *S. bolavenensis* is easily distinguishable from *S. glabra* by its long-spicately disposed sessile umbels (vs. only 1 umbel axillary in *S. glabra*) and its cordate leaf base (vs. cuneate to rounded).

Additional specimen examined:—LAOS. Champasak Province: Paksong District, near Nong Luang Village, Dong Hua Sao National Park (Bolaven Plateau), 15°04'19.26"N, 106°12'38.67"E, 1248 m a.s.l., 10 Dec 2018, *Tagane et al. L1964* (FOF 005293, FU, KAG 128065).

Acknowledgments

Authors thank the manager and staff of Dong Hua Sao National Park for permitting our botanical inventories in the protected area. This study was supported by Nagao Natural Environment Foundation, Japan and partly by JSPS KAKENHI (21K06307 and 21KK0132).

References

- Baruah, S., Sarma, J. & Borthakur, S.K. (2018) *Smilax sailenii* (Smilacaceae) – a new species from Assam, North East India. *Taiwania* 63: 32–36.
<https://doi.org/10.6165/tai.2018.63.32>
- Cameron, K.M. & Fu, C.X. (2006) A nuclear rDNA phylogeny of *Smilax* (Smilacaceae). *Aliso* 22: 598–605.
<https://dx.doi.org/10.5642/aliso.20062201.47>
- Chen, X. & Koyama, T. (2000) Smilacaceae. In: Wu, Z.Y., Raven, P.H. & Hong D.Y. (Eds.) *Flora of China*, vol. 24. Missouri Botanical Garden Press, St. Louis and Science Press, Beijing, pp. 96–115. Available from: http://www.efloras.org/flora_page.aspx?flora_id=2 (accessed 1 March 2022)
- De Candolle, A. & De Candolle, C. (1878) *Monographiae phanerogamarum 6: Smilacaceae, Restiaceae, Meliaceae*. Masson, Paris, 783 pp.
- Diels, L. (1900) Die Flora von Central-China. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 29 (2): 169–659.
- Feng, J.Y., Jin, X.J., Zhang, S.L., Yang, J.W., Fei, S.P., Huang, Y.S., Liu, Y., Qi, Z.C. & Li, P. (2022) *Smilax weniae*, a new species of Smilacaceae from limestone areas bordering Guizhou and Guangxi, China. *Plants* 11: 1032.
<https://doi.org/10.3390/plants11081032>
- Gagnepain, F. (1934) Liliacées. In: Lecomte, H. (Ed.) *Flore générale de l'Indo-Chine*, vol. 6. Masson, Paris, pp. 753–815.

- Gagnepain, F. (1934) Deux Liliacées nouvelles. *Bulletin de la Société Botanique de France* 81: 619.
<https://doi.org/10.1080/00378941.1934.10834005>
- Heckroth, H.P., Moong, J., Janka, H.I., Fiala, B., Chung, A.Y.C. & Maschwitz, U. (2004) *Smilax borneensis* (Smilacaceae), an unspecific climbing ant-plant from Borneo and myrmecophytic traits in other Asiatic *Smilax* species. *Sandakania* 14: 33–50.
- Hô, P. H. (2000) *Cay Co Viet Nam: an illustrated flora of Vietnam*, vol. 3. The author, Montreal, 1020 pp.
- Hooker, J.D. (1892) *Smilax*. In: Hooker, J.D. (Ed.) *The flora of British India*, vol 6, Reeve, London, pp. 302–314.
- Jin, X.J., Chen, Y., Lee J., Qi, Z.C., Liu, L.X., Li, Q. & Fu, C.X. (2016) A new species of *Smilax* (Smilacaceae) from Yunnan, China. *Phytotaxa* 275 (2): 159–167.
<https://doi.org/10.11646/phytotaxa.275.2.7>
- Kladwong, P. & Chantaranothai, P. (2020) Taxonomic notes on genus *Smilax* L. (Smilacaceae) from the mainland Southeast Asia. *Tropical Natural History* 20: 244–255.
- Kladwong, P. & Chantaranothai, P. (2022) Notes on *Smilax bockii* and *S. seisuiensis* (Smilacaceae), two newly recorded species from Laos and Thailand. *Thai Forest Bulletin (Botany)* 52: 153–160.
<https://doi.org/10.20531/tfb.2022.50.2.15>
- Kladwong, P. Chantaranothai, P. & Simpson, D.A. (2018) Two new names and five lectotypified taxa for the genus *Smilax* (Smilacaceae) and the transfer of *Smilax petiolatumidus* to the genus *Dioscorea* (Dioscoreaceae). *Thai Forest Bulletin (Botany)* 46: 44–57.
<https://doi.org/10.20531/tfb.2018.46.1.07>
- Koyama, T. (1975) Smilacaceae. In: Smitinand, T. & Larsen, K. (Eds.) *Flora of Thailand*, vol. 2(3). Applied Scientific Research Corporation of Thailand, Bangkok, pp. 211–250.
- Koyama, T. (1983) Smilacaceae. In: Leroy, Jean-F. (Ed.) *Flore du Cambodge, du Laos et du Viêt-Nam*, vol. 20. Muséum national d'histoire naturelle, Paris, pp. 69–124.
- Kunth, K.S. (1850) *Enumeratio plantarum*. Cottae, Stuttgart, 908 pp.
- Li, P., Qi, Z.C., Chen, S.C., Cameron, K.M. & Fu, C. (2011) *Smilax ligneoriparia* sp. nov.: a link between herbaceous and woody *Smilax* (Smilacaceae) based on morphology, karyotype and molecular phylogenetic data. *Taxon* 60: 1104–1112.
<https://doi.org/10.1002/tax.604013>
- Linnaeus, C. (1753) *Species plantarum*. Salvius, Stockholm, 1200 pp.
- Newman, M.F., Pullan, M., Ketphanh, S., Svengsuksa, B., Thomas, P., Sengdala, K., Lamxay, V. & Armstrong, K. (2017 onwards) *A checklist of the vascular plants of Lao PDR*. Available from: <https://padme.rbge.org.uk/laos/> (accessed 21 February 2022)
- Noyori, W., Komada, N., Souladeth, P. & Tagane, S. (2022) *Camchaya bolavenensis* (Asteraceae: Vernoniaceae), a new species from Bolaven Plateau, southern Laos. *Phytotaxa* 536 (1): 1–6.
<https://doi.org/10.11646/phytotaxa.536.1.1>
- Phonpaseuth, P., Souladeth, P., Souvannakhoummane, K., Vongthavone, T. & Tagane, S. (2021) Two new species of *Sonerila* Roxb. (Melastomataceae) from Laos. *European Journal of Taxonomy* 755: 136–148.
<https://doi.org/10.5852/ejt.2021.755.1403>
- POWO (2022) *Plants of the World online*. Facilitated by the Royal Botanic Gardens, Kew. Available from: <http://www.plantsoftheworldonline.org/> (accessed 1 March 2022)
- Profile, E. (2012) *Indo-Burma biodiversity hotspot*. Update Critical Ecosystem Partnership-Fund, Washington, 360 pp.
- Roxburgh, W. (1832) *Smilax*. In: Roxburgh, W. (Ed.) *Flora Indica*, vol. 3, Thacker, Serampore, pp. 791–796.
- Qi, Z.C., Cameron, K.M., Li, P., Zhao, Y.P., Chen, S.C., Chen, G.C. & Fu, C.X. (2013a) Phylogenetics, character evolution, and distribution patterns of the greenbriers, Smilacaceae (Liliales), a near-cosmopolitan family of monocots. *Botanical Journal of the Linnean Society* 173: 535–548.
<https://doi.org/10.1111/boj.12096>
- Qi, Z.C., Li, P. & Fu, C.X. (2013b) New combinations and a new name in *Smilax* for species of *Heterosmilax* in Eastern and Southeast Asian Smilacaceae (Liliales). *Phytotaxa* 117 (2): 58–60.
<https://dx.doi.org/10.11646/phytotaxa.117.2.4>
- Qi, Z.C., Shen, C., Han, Y.W., Wang, R.H., Liu, R.Z., Naiki, A. & Li, P. (2016) A new species of *Smilax* (Smilacaceae) from southern island of Kagoshima prefecture, Japan. *Phytotaxa* 269 (2): 120–130.
<http://dx.doi.org/10.11646/phytotaxa.269.2.7>
- Souladeth, P., Tagane, S., Newman, M.F. & Prajaksood, A. (2020) Two new species of *Eriocaulon* (Eriocaulaceae) from Laos. *Kew Bulletin* 75: 56.
<https://doi.org/10.1007/S12225-020-09909-0>
- Souvannakhoummane, K., Lanorsavanh, S., Tagane, S., Souladeth, P., Phonpaseuth, P., Pongamornkul, W. & Lamxay, V. (2021) Six new species and eight new records of Gesneriaceae from Laos. *Gardens' Bulletin Singapore* 73: 427–456.
[https://doi.org/10.26492/gbs73\(2\).2021-13](https://doi.org/10.26492/gbs73(2).2021-13)

- Sun, Z.Y., Wang, Y.H. Zhao, Y.P. & Fu, C.X. (2015) Molecular, chromosomal and morphological characters reveal a new diploid species in the *Smilax china* complex (Smilacaceae). *Phytotaxa* 212 (3): 199–212.
<https://doi.org/10.11646/phytotaxa.212.3.2>
- Tagane, S., Souladeth, P., Kongxaysavath, P., Rueangurea, S., Suddee, S., Suyama, Y., Suzuki, E. & Yahara T. (2020) Two new species and 18 new records for the flora of Laos. *Thai Forest Bulletin (Botany)* 49: 111–126.
<https://doi.org/10.20531/tfb.2021.49.1.14>
- Vongthavone, T., Tagane, S., Phonpaseuth, P., Souvannakhoummane, K. & Souladeth, P. (2021) Twelve new records of pteridophytes from Bolaven Plateau, sothern Laos. *Thai Forest Bulletin (Botany)* 49: 182–190.
<https://doi.org/10.20531/tfb.2021.49.2.05>
- Yamazaki, T., Souladeth, P. & Tagane S. (2021) *Strobilanthes bolavenensis*, a new species of Acanthaceae from Bolaven Plateau, southern Laos. *Phytotaxa* 513 (2): 152–158.
<https://doi.org/10.11646/phytotaxa.513.2.6>