

Two new species of Araliaceae from the limestone karst of Central Laos

DEUANTA KONGXAISAVATH^{1,5}, SHUICHIRO TAGANE^{2,6}, TAKENORI YAMAMOTO^{2,7}, PHONGPHAYBOUN PHONEPASEUTH^{3,8}, VAN-SON DANG^{4,9} & PHETLASY SOULADETH^{1,10*}

¹Faculty of Forest Science, National University of Laos, Dongdok Campus, Xaythany District, Vientiane Capital, Laos

²Kagoshima University Museum, Kagoshima University, 1-21-30, Korimoto, Kagoshima 890-0065, Japan

³Faculty of Environmental Sciences, National University of Laos, Dongdok Campus, Xaythany District, Vientiane Capital, Laos;
Department of Biological Sciences, Graduate School of Science, Tokyo Metropolitan University, 1-1 Minamiosawa, Hachioji, Tokyo 192-0397, Japan

⁴Institute of Life Sciences, Vietnam Academy of Science and Technology, 85 Tran Quoc Toan Street, District 3, Ho Chi Minh City, Vietnam

⁵✉ deuantar66789@gmail.com;  <https://orcid.org/0009-0000-4559-0647>

⁶✉ stagane29@gmail.com;  <https://orcid.org/0000-0002-1974-7329>

⁷✉ t.yamamoto0624@gmail.com;  <https://orcid.org/0000-0002-5487-6652>

⁸✉ p.phonepaseuth@nuol.edu.la;  <https://orcid.org/0000-0001-5514-777X>

⁹✉ dvsonitb@gmail.com;  <https://orcid.org/0000-0001-8681-4141>

¹⁰✉ p.souladeth@nuol.edu.la;  <https://orcid.org/0000-0001-8563-1773>

*Author for correspondence

Abstract

Two new species of Araliaceae, *Heptapleurum karsticola* and *Trevesia laotica*, are described from limestone karst in central Laos. *Heptapleurum karsticola* was found in Khammouane Province, and the second species, *T. laotica*, was found in Bolikhamxai and Khammouane Provinces. Information regarding the distribution, habitat, phenology, vernacular names, preliminary conservation assessment, DNA barcodes, and taxonomic notes compared to other closely related species is provided for each species.

Key words: Apiales, endemic species, flora, Indochina, phylogeny, taxonomy

Introduction

Araliaceae, commonly known as the ginseng family, comprise 46 genera and ca. 1,500 species that are widely distributed across the tropics and subtropics, with a few genera extending into the temperate zones (Wen *et al.* 2001, Plunkett *et al.* 2004, POWO 2025). Members of the family are mostly woody plants, with leaves ranging from simple to palmately or pinnately compound, and characterized by inflorescences usually in compound umbels (rarely racemes or heads), mostly 5-merous flowers with inferior ovaries, and drupaceous fruits containing 2–5 (rarely more) seeds (Philipson 1970, Yi *et al.* 2004). In Laos, 26 species and two varieties in 11 genera are enumerated to date (Newman *et al.* 2017–present).

During our field surveys in the limestone karst of central Laos, including Nam Kading National Protected Area (hereafter called Nam Kading NPA) in Bolikhamxai Province, Nam Sanam-Phou Pha Marn Provincial Protected Area (hereafter called Nam Sanam-Phou Pha Marn PPA) in Khammouane Province, we collected two unknown species of Araliaceae, belonging to *Heptapleurum* Gaertner (1791: 472), and *Trevesia* Visiani (1842: 262). After a careful examination based on taxonomic literature and herbarium specimens of each genus, as well as phylogenetic relationship, they appear to be hitherto undescribed taxa. Thus, we describe them as new species, *Heptapleurum karsticola* Kongx., Tak.Yamam. & Tagane and *Trevesia laotica* Kongx., Soulad. & Tagane as below.

Materials and Methods

Field surveys and morphological investigation

Nine field surveys were carried out from December 2016 to March 2025 to assess the plant diversity in the previously mentioned protected areas in the two provinces of Bolikhamxai and Khammouane. A total of nine specimens representing two unknown species of Araliaceae were carefully examined to confirm the validity of the new species, and the description of each species was based on these collections. We consulted related species using dry herbarium specimens at FOF, HN, KAG, VNM, as well as digitized images from several online herbaria databases, including the Muséum National d'Histoire Naturelle (<https://science.mnhn.fr/>), the Kew Data Portal (<https://data.kew.org/>), Natural History Museum Data Portal (<https://data.nhm.ac.uk/>), AAU Herbarium Database (<https://sciencemuseumne.dk/herbariet/>), and the Catalogue des herbier de Genève (<https://www.ville-ge.ch/musinfo/bd/cjb/chg/>). Also, we referred to relevant literature (Li 1942, Hoo & Tseng 1978, Shang 1997, Wen *et al.* 2001, 2003, 2007, Yi *et al.* 2004, Wen & Lowry 2006, Shang & Lowry 2007, Newman *et al.* 2007, 2017–present, Esser & Jebb 2010, 2019, Tagane *et al.* 2015, 2017, Esser 2018, de Kok 2022).

DNA extraction, sequencing and phylogenetic analysis

The total DNA was extracted from silica dried leaf following Doyle & Doyle (1987). The nuclear ribosomal internal transcribed spacer (ITS) region, primers (51NT and 26S1R) were amplified and sequenced according to published protocols following Clement *et al.* (2004). Purification, labeling and sequencing followed Souvannakhoummane *et al.* (2025).

For the phylogenetic analysis, we sampled taxa closely related to *Heptapleurum* and *Trevesia* within the Asian core Araliaceae (Wen *et al.* 2001) and the Asian Palmate group (Plunkett *et al.* 2004). The data set included six sequences of *Aralia* Linnaeus (1753: 273), *Schefflera* Forster & Forster (1776: 45), eight sequences of *Brassaiopsis* Decaisne & Planchon (1854: 106), three sequences of *Panax* Linnaeus (1753: 1059), two sequences of *Cussonia* Thunberg (1780: 210), *Dendropanax* Decaisne & Planchon (1854: 107), *Heteropanax* Seemann (1866: 297), *Osmoxylon* Miquel (1863: 5), *Raukaua* Seemann (1866: 352), *Tetrapanax* (Koch 1859a: 70) Koch (1859b: 371), and representatives of *Eleutherococcus* Maximowicz (1859: 132), *Harmsiopanax* Warb. (*in* Engler 1897: 166), *Oplopanax* (Torrey & Gray 1840: 648) Miquel (1863: 16), *Seemannaralia* Viguier (1906: 118), *Sinopanax* Li (1949: 231), 21 sequences of *Heptapleurum*, eight sequences of *Heptaphylla* and nine sequences of *Trevesia*. Most sequences were obtained from GenBank and derived from previous phylogenetic studies (Plunkett *et al.* 2004, 2019, Wen *et al.* 2007, Li & Wen 2013, 2014, Valcarcel *et al.* 2014). *Pittosporum illicioides* Makino (1900: 31) of Pittosporaceae was selected as the outgroup following Li & Wen (2014). In total of 72 ITS sequences, including two newly generated sequences, were analyzed. Details of each species, vouchers, country of origin and GenBank accession number are shown in Appendix Table S1.

The sequence alignment was performed with MAFFT (Katoh *et al.* 2019), followed by manual checking and editing in Aliview (Larsson 2014). A maximum likelihood tree of ITS was constructed and analyzed using the IQ-TREE web server (Nguyen *et al.* 2015) and employing one-thousand bootstrap pseudo replicates via the ultrafast bootstrap (Hoang *et al.* 2018). Tree was displayed and annotated using Interactive Tree of Life (iTOL) v5 (<https://itol.embl.de/>) (Letunic & Bork 2021).

Results and discussion

From 72 aligned sequences of ITS, which have 665 aligned positions and contain 272 variable characters, 183 of which are parsimony-informative. Maximum likelihood tree based on ITS showed that two new species are located within the clade of the Asian Palmate group of Araliaceae with strong bootstrap support (BS = 100) (Fig. 1).

The sample of *Heptapleurum* from Laos (Tagane *et al.* Z1201) falls within the strongly supported *Heptapleurum* clade (BS = 99). Within the clade, it is placed as sister to a clade including *H. petelotii* (Merrill 1924: 428) Plunkett & Lowry (*in* Lowry & Plunkett 2020: 162), *H. oblongum* (Craib 1930: 421) Plunkett & Lowry (*in* Lowry & Plunkett 2020: 160), *H. arboricola* Hayata (1916: 23), and *H. leucanthum* (Viguier 1909: 358) Y.F. Deng (*in* Ai 2018: 473) (BS = 96); together, Z1201 and this clade form the sister group to *H. bengalense* (Gamble 1919: 229) Esser (2021: 152) (BS = 100).

The sample of *Trevesia* from Laos (*Tagane et al. Z1168*) falls within the *Trevesia* clade with strong bootstrap support (BS=99). Within this clade, it is recovered as sister to a clade comprising *T. sphaerocarpa* Grushvitzky & Skvortsova (*in* Grushvitzky *et al.* 1984: 1025) and *T. palmata* (Roxb. ex Lindley 1825: t. 894) Visiani (1842: 262) (BS=88); together, these three species form the sister group to *T. sundaica* Miquel (1856: 137) (BS = 99).

Based on these results and their morphology, it is clear that the two species we discovered this time belong to *Heptapleurum* and *Trevesia*, respectively.




FIGURE 1. ML tree of Asian palmate group and related genera based on the nuclear ribosomal internal transcribed spacer (ITS) regions, including two new species from Laos showing in bold. Numbers on the branches indicate support values, only ≥ 50 are shown (1000 ultrafast bootstrap replicates).

Taxonomy

Heptapleurum karsticola Kongx., Tak.Yamam. & Tagane, *sp. nov.* (Figs. 2, 3)

TYPE:—LAOS. Khammouane Province, Khounkham District, the Rock Viewpoint at Phou Pha Marn, in limestone karst, 18.17689°N, 104.48409°E, 435 m elev., 19 March 2024, fl. & young fr., *S. Tagane*, *P. Souladeth*, *V.S. Dang*, *T. Yamamoto*, *K. Souvannakhoummane*, *N. Tanaka*, *D. Kongxaisavath*, *K. Phengmala*, *A. Sengthonh*, *Q.B. Nguyen* & *K. Takahashi* Z1201 (holotype: FOF [FOF0007163!], isotypes: KAG [KAG187600!], VNM [VNM00074078!]).

Heptapleurum karsticola is similar to *H. leucanthum* distributed in South-Central to Southeast China, Laos, Thailand, and Vietnam in having pleasant smell when crushed, glabrous leaves with 4–9 leaflets, secondary veins up to 6 pairs, leaflet laminae up to 9 × 2.8 cm, inflorescences with a short main axis of 0.3–2.7 cm long and 4 or 5 stamens, but distinguished by its leaf margin repand, sometimes sparsely serrate near apex (vs. entire in *H. leucanthum*), slightly glaucous lower leaf surface (vs. not glaucous), larger bracts (3–5 mm long vs. ca. 1 mm long), smaller umbels (0.7–1.5 cm in diam. vs. 2–2.5 cm in diam.), and shorter pedicels (2–3.5 mm long vs ca. 6 mm long) (Table 1).

TABLE 1. Morphological comparison between *Heptapleurum karsticola* and *H. leucanthum*. The measurements of the latter species are based on Shang & Lowry (2007) and Esser & Jebb (2019).

Character	<i>H. karsticola</i>	<i>H. leucanthum</i>
Habit	small shrubs	small shrubs to scandent
Leaf margin	margin slightly repand, sometimes sparsely serrate near apex	entire
Abaxial leaf surface	slightly glaucous	not glaucous
Length of petiolules (cm)	0.2–1.8	0.5–3(–5)
Length of bracts (mm)	3–5	ca. 1
Size of umbels in diam. (cm)	0.7–1.5	2–2.5
Length of pedicels (mm)	2.0–3.5	ca. 6

Small shrubs, up to 4 m tall, deciduous, all parts glabrous; bark smooth, brown to greyish brown, current year twigs green *in vivo*, up to 0.4 cm in diam., with prominent lenticels, old twigs brown to greyish brown, with pleasant smell when crushed. Leaves palmate, alternate; stipules adnate to base of petioles; petiole 3.5–9.5 cm long; leaflets (4–)5(–6), petiolules 0.2–1.8 cm long, laminae uniform in size, elliptic to narrowly elliptic, 3.5–8.5 × 1.1–2.8 cm, slightly glaucous on lower surface *in vivo*, base cuneate to attenuate, margin slightly repand, sometimes sparsely serrate near apex, apex acute to attenuate, midrib prominent on both surfaces, secondary veins 2–6 pairs. Inflorescences terminal, consisting of cymes of umbels, 2–5 umbels per inflorescence, 2.3–7.3 cm long, peduncle 0.3–1.2 cm long; primary axis 0.3–2.7 cm long, with 2–4 lateral branches; bracts broadly triangular, suborbicular to broadly elliptic, 3–5 × 3–7 mm, apex apiculate; umbels 0.7–1.5 cm in diam., with 6–12 flowers. Flowers bisexual, yellow *in vivo*; pedicel 2–3.5 mm long; bracteoles 2.3–4 × 1.2–2.7 mm; calyx cupules obconical, 1.5–2.1 × 1–2.5 mm, calyx lobes (3–)5, minute; petals (4 or)5, ovate-triangular, 1.1–1.8 × 0.7–1.2 mm, apex acute, margin entire, caducous; stamens (4 or)5, filaments 1.3–4 mm long, anthers 0.2–0.5 mm long; ovary (5 or)6-locular, style absent, stigma truncated conical, up to 0.2 mm long. Infructescences up to 2.5 cm long. Fruits on pedicels 3–6 mm long, ellipsoid-oblong, 3–7 mm long, 2–5 mm in diam., smooth, with persistent short stigma.

Additional specimens examined (paratypes):—Khammouane Province: Khounkham District, the Rock Viewpoint at Phou Pha Marn, 18.177497°N, 104.483794°E, 446 m elev., 2 July 2022, young fl. *Souladeth et al. PPM13* (FOF); same locality, 18.17756°N, 104.4848°E, 455 m elev., 19 March 2024, young fr. *Tagane et al. Z1204* (FOF [FOF0007164], KAG [KAG187604, KAG187605], VNM); same locality, same date, fl. & young fr., *Tagane et al. Z1210* (FOF [FOF0007162], KAG [KAG187612], VNM); same locality, 18.1763°N, 104.4842°E, 420 m elev., 9 March 2025, fl. & fr., *Tagane et al. Z2412* (FOF [FOF0009950]; KAG [KAG201134]; VNM).

Distribution:—Laos (Khammouane Province).

Habitat and Phenology:—On open limestone karst, at 420–455 m elev. Flowering and fruiting in March and July.

Etymology:—The specific epithet “*karsticola*” refers to its habitat (limestone karst) of the type locality.

Vernacular name:—Tarngh Pha Hom (ຕ້ງຢາງຫຼອມ; Lao), suggested here. “Tarngh” refers to the common name for Araliaceae in Lao in general. “Pha” refers to its habitat of a mountain cliff, and “Hom” refers to a good smell of this species.




FIGURE 2. *Heptapleurum karsticola* Kongx., Tak.Yamam. & Tagane. A. Habit. B. Flowering twigs. C. Portion of inflorescence. D. Flower. E & F. Bracts, abaxial (left) and adaxial side (right). G & H. Bracteole, abaxial (left) and adaxial side (right). I & J. Petals, abaxial (left) and adaxial side (right). K. Infructescence. L. Young fruit. M. Transverse section of a fruit, showing 6 locules. Photos A, B, E–J and L–M by D. Kongxaisavath., C & K by S. Tagane. Materials E–J from Tagane *et al.* Z1201 and L–M from Tagane *et al.* Z2412 (FOF). Scale bars C & K = 1 cm, D–M = 1 mm.



FIGURE 3. Holotype of *Heptapleurum karsticola* Kongx., Tak.Yamam. & Tagane. (*Tagane et al.* Z1201 (F0F0007163)).

GenBank accession no.:—PV821477 (ITS) (sequenced from *Tagane et al.* Z1201).

Preliminary conservation assessment:—Critically Endangered (CR). Currently, only a single population of *Heptapleurum karsticola* is found in a limited area of ca. 200 × 200 m at an open limestone karst along a walking trail within the Nam Sanam-Phou Pha Marn PPA in Khammouane Province, central Laos, where we observed a total of about 30 mature individuals. The similar habitat of the limestone karst is widely distributed in the Phou Hin Phoun NPA, the southern adjacent area of the PPA, and more populations and individuals can be expected to occur. However, some areas have suffered some degree of impact from activities such as limestone mining. Given its scarcity, it is assessed as Critically Endangered (CR) according to IUCN criterion B2a, b(iii), C2a(i, ii) and D (IUCN 2024).

Notes:—The new species is assigned to *Heptapleurum*, a monophyletic genus with 317 species currently accepted (Lowry & Plunkett 2020, POWO 2025) based on morphological characteristics, such as unarmed stems, palmately compound leaves, the leaflets with distinct, usually articulated petiolules, and valvate petals (Esser 2021).

***Trevesia laotica* Kongx., Soulad. & Tagane, *sp. nov.* (Figs. 4, 5)**

TYPE:—LAOS. Khammouane Province, Khounkham District, the Rock Viewpoint at Phou Pha Marn, in evergreen broad-leaved forest on limestone hills, 18.17646°N, 104.48848°E, 453 m elev., 8 March 2025, fl. & fr., *S. Tagane, P. Souladeth, V.S. Dang, T. Vongthavone, A. Sengthong, D. Kongxaisavath, K. Phengmala, T. Yamamoto, H. Toyama, Q.T. Pham & K. Yamazaki* Z2365 (holotype: FOF [FOF0009947!], flowers in spirit]; isotypes KAG [KAG201088!]; VNM [VNM00074074!]).

Trevesia laotica is morphologically similar to *T. palmata* but easily distinguished by its leaves (palmately compound in *T. laotica* vs. simple, deeply 5–9-lobed in *T. palmata*), longer filaments (8–9 mm long vs. 4–6 mm long), smaller calyx lobes (0.1–0.5 mm long vs. 1–2 mm), and longer styles on fruits (4–5.5 mm long vs. 2–4 mm long) (Table 2).

TABLE 2. Morphological comparison between *T. laotica* and *T. palmata*. The characters of the latter species are based on Shang & Lowry (2007).

Character	<i>T. laotica</i>	<i>T. palmata</i>
Height (m)	up to 4	up to 8
Leaf shape	palmately compound with 5–9 leaflets	palmately 5–9-lobed
Size of leaflet blades (cm)	7–41.5 × 1.3–10	60–90
Length of inflorescence (cm)	80	ca. 45
Length of peduncles (cm)	1.5	4–17
Length of calyx lobes (mm)	0.1–0.5	1–2
Number of stamens	7	7–12
Ovary	7-locular	7–12-locular
Diameter of fruits (cm)	2	1–1.8
Length of styles on fruits (mm)	4–5.5	2–4

Trees up to 4 m tall, evergreen. Stems with a few scattered prickles, bark grey to pale green; young branches sparsely to densely covered with ferruginous indumentum. Leaves palmately compound, with 5–9-leaflets; stipules connate to base of petiole, triangular, 0.6–1.1 cm long, sometimes bi-lobed to near the base; petioles 12.5–67.7 cm long, without spines; petiolules 0.3–2.5 cm long; leaflet blades lanceolate, oblong-lanceolate, narrowly oblong-elliptic, 7–41.5 × 1.3–10 cm, upper surface dark greyish brown, glabrous, lower surface yellowish brown, covered with indumentum when young, glabrous when mature, base narrowly attenuate, apex acuminate to caudate, margin serrate, midrib on each lobe prominent on both surfaces, lateral veins prominent on both surfaces. Inflorescences panicle of 1–4 umbels, terminal or axillary, 24 cm long (for solitary), up to 80 cm long (for 4-umbellate) pendulous; peduncle 1.5 cm long, covered with brown indumentum outside, glabrescent; bracts ovate-triangular, 0.6–1.1 × 0.3–0.4 cm, densely covered with brown indumentum outside, glabrous inside, apex acute, margin entire. Flowers 36–42 per umbels; pedicels 1.6–1.7 cm long, covered with brown indumentum, glabrescent; bracteoles narrowly triangular to linear, 1.8–3 mm long, apex cuspidate, densely covered with indumentum outside; calyx campanulate, 0.6–0.7 cm long, 0.5–0.6 cm in diam., sparsely covered with indumentum, calyx rim minutely 7-teethed, lobes 0.1–0.5 mm long; petals 7, often partly connate appearing 5, spread, triangular to ovate-triangular, 7–8 × 3–5.5 mm, glabrous on both surfaces, apex acute; stamens 7, filaments 8–9 mm long, anthers 2–3.8 × 1–2 mm; ovary 7-locular, with 1 or 2 ovules per carpel, style 2.2 mm long, united, conical, glabrous, stigma 7-lobed. Infructescences up to 87 cm long; axis sparsely to densely with



FIGURE 4. *Trevesia laotica* Kongx., Soulad. & Tagane. A. Habit. B. Fruiting branch. C. Trunk. D. Apical part of stem showing base of petioles and stipules. E. Sapling. F. Leaves. G. Portion of lower leaf surface. H. Inflorescence. I, J. Flowers. K. Fruits. L. Fruits dissected to reveal 7 developing seeds. M. Seeds. Photos A, H–J & M from Tagane *et al.* Z2365, B–D, F & G from Tagane *et al.* Z1168, E with no voucher, K & L from Tagane *et al.* Z798, taken by S. Tagane. Scale bars I–J & L–M = 1 cm.



FIGURE 5. Holotype of *Trevesia laotica* Kongx., Soulad. & Tagane (*Tagane et al.* Z2365 (FOF0009947)).

brown farinose stellate pubescence. Fruits globose to obovoid-globose, to 2 cm in diam., smooth; fruiting pedicel 1.7–3.5 cm long; style persistent, 4–5.5 mm long, stylopodium persistent. Seeds 4–7, ovoid-ellipsoid, flattened, $1.1\text{--}1.5 \times 0.7\text{--}1.0 \times 0.3\text{--}0.5$ cm.

Additional specimens examined (paratypes):—LAOS. Bolikhamxai Province: Pak Kading District, Ban Nabouay, 18.243472°N, 104.269306°E, 206 m elev., 7 November 2023, fl. buds, *Souladeth et al.* Z746 (FOF [FOF0005875], KAG [KAG187164], VNM [VNM00070350]); Pak Kading District, near Ban Naphong, Nam Kading National Protected Area, 18.22257°N, 104.38119°E, 602 m elev., 18 March 2024, fr., *Tagane et al.* Z1168 (FOF [FOF0007169, FOF0007170], KAG [KAG187570], VNM [VNM00073014]). Khammouane Province: Khounkham District, Tad Nam Sanam, 18.21963°N, 104.50306°E, 458 m elev., 15 March 2024, fr., *Tagane et al.* Z798 (FOF [FOF0005925], KAG [KAG187213, KAG187214], VNM [VNM00070840]).

Distribution:—Laos (Bolikhamxai and Khammouane Provinces).

Habitat and phenology:—In evergreen broad-leaved forest on limestone hill at 200–610 m elev. Flowering in December to March, and fruiting from March to April.

Etymology:—The specific epithet “*laotica*” refers to the country of the type locality.

Vernacular name:—Phak Narm Ton (ຜັກໜາມຕົ້ນ; Lao), suggested here. “Phak Narm” refers to plants with spines in Lao, and “Ton” refers to the tree habit of this species.

GenBank accession no.:—PV821478 (ITS) (sequenced from *Tagane et al.* Z1168).

Preliminary conservation assessment:—Endangered (EN). Currently, *Trevesia laotica* is known from four localities in limestone areas in Bolikhamxai and Khammouane Provinces of central Laos. We observed 10–30 mature individuals in each location. Based on our specimen records, the Area of Occupancy (AOO) is calculated as 12 km² and the Extent of Occupancy (EOO) is 208 km² using GeoCAT (Bachman *et al.* 2011). Although the locations are situated within the protected area of Nam Sanam-Phou Pha Marn PPA and Nam Kading NPA, some disturbances are observed, which might bring a negative impact on its survival. Given this situation, the species is being assessed as Endangered (EN) according to the IUCN criteria B1a & b(ii, iii), B2a & b(ii, iii), and D (IUCN 2024).

Notes:—The genus *Trevesia* was described by Visiani in 1842. Due to its unique characteristics, such as palmately lobed leaves and pedicels lacking articulations, it was segregated from *Gastonia* Comm. ex Lam. in Lamarck *et al.* (1788: 610). The most recent revision by Jebb (1998) recognized seven distinct species. Since then, one more species, *T. vietnamensis* J. Wen & P.K. Lôt in Wen *et al.* (2007), was described. Our new species is the 9th species in the genus.

The genus *Trevesia* is closely related to *Brassaiopsis* (Li & Wen 2016), and the two genera are sister but distinct in phylogenetic analysis based on ITS (Fig. 1). The two genera are traditionally distinguished by leaf shape (simple, palmately lobed in *Trevesia* vs. simple (entire or palmately lobed) or palmately compound in *Brassaiopsis*), the merosity of floral parts (at least 7-merous in *Trevesia* vs. 5-merous in *Brassaiopsis*), and the locular number of the ovary (6–16-locular vs. 2–5-locular) (Jebb 1998, Shang & Lowry 2007). However, the description of *T. vietnamensis*, in 2007, blurred the generic boundaries. This species is positioned in the *Trevesia* clade and sister to *T. lateospina* in the phylogenetic analysis based on nuclear (ITS) and ribosomal (*trnL-F* and *ndhF*) datasets, but shows *Brassaiopsis*-like characters, including palmately compound leaves and 4–5-merous flowers with 4–5 locular ovaries (Wen *et al.* 2007). Our new species possesses 7-merous flowers with 7-locular ovaries that are consistent with the traditional characteristics of *Trevesia*, and is the second species having palmately compound leaves. As Wen *et al.* (2007) proposed, it might be appropriate to merge *Brassaiopsis* with *Trevesia*. Further comprehensive morphological and phylogenetic studies based on a large number of specimens are needed.

Acknowledgements

The authors would like to thank the managers and staff of Nam Kading NPA in Bolikhamxai Province and Nam Sanam-Phou Pha Marn PPA in Khammouane Province, for permitting our botanical inventories in these conservation areas. We are especially grateful to Vongpheng Phommala (Provincial Agriculture and Forestry Office in Bolikhamxai), Thyraphon Vongthavone (ACD), Keoudone Souvannakhommmane, Kajonesuk Phengmala, Anousone Sengthong (NUoL), Kaito Yamazaki (KAG), Quoc Bao Nguyen and Quoc Trong Pham (VNM) for their assistance and support in specimen collection during fieldwork. Thanks to the Systematic Botany Laboratory, Tokyo Metropolitan University, for supporting our molecular study. This study was supported by the Nagao Natural Environment Foundation, Japan, and partially by the Lao Landscapes and Livelihoods Project under the Department of Forestry, Laos.

References

- Ai, T. (2018) *Medicinal Flora of China, Zhongguo Yaoyong Zhiwuzhi (Diqijuan)*. Peking University Medical Press, Beijing, 1344 pp. [in Chinese]
- Bachman, S., Moat, J., Hill, A.W., de la Torre, J. & Scott, B. (2011) *Supporting Red List threat assessments with GeoCAT: geospatial conservation assessment tool*. Available from: <https://geocat.iucnredlist.org/> (accessed 4 April 2025)
- Clement, W.L., Tebbitt, M.C., Forrest, L.L., Blair, J.E., Brouillet, L., Eriksson, T. & Swensen, S.M. (2004) Phylogenetic position and biogeography of *Hillebrandia sandwicensis* (Begoniaceae): a rare Hawaiian relict. *American Journal of Botany* 91: 905–917.
- Craib, W.G. (1930) Contributions to the Flora of Siam. Additamentum XXIX. *Bulletin of Miscellaneous Information Kew* 1930: 405–427.
- Decaisne, J. & Planchon, J.E. (1854) Esquisse d'une monographie des Araliacées. *Revue Horticole, Séries 4* 3: 104–109.
- de Kok, R.P.J. (2022) The Araliaceae of Peninsular Malaysia: five new species, two neotypifications and twenty-two lectotypifications in several genera. *Gardens' Bulletin Singapore* 74: 71–90.
[https://doi.org/10.26492/gbs74\(1\).2022-06](https://doi.org/10.26492/gbs74(1).2022-06)
- Doyle, J.J. & Doyle, J.L. (1987) A rapid isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* 19: 11–15.
- Engler, A. (1897) *Die Natürlichen Pflanzenfamilien. Nachträge zum II. bis IV, Teil 1*. Engelmann, Leipzig, pp. 1–380.
- Esser, H.-J. (2018) A new species of *Brassatopsis* (Araliaceae) from Thailand and lectotypifications of names for related taxa. *Thai Forest Bulletin (Botany)* 46: 129–133.
<https://doi.org/10.20531/tfb.2018.46.2.03>
- Esser, H.-J. (2021) *Schefflera* (Araliaceae) in Thailand becomes *Heptapleurum*. *Thai Forest Bulletin (Botany)* 49: 151–154.
<https://doi.org/10.20531/tfb.2021.49.1.19>
- Esser, H.-J. & Jebb, M.H.P. (2010) A new *Schefflera* and taxonomic notes on Araliaceae from Thailand. *Thai Forest Bulletin (Botany)* 38: 160–166.
- Esser, H.-J. & Jebb, M.H.P. (2019) Araliaceae. In: Chayamarit, K. & Balslev, H. (Eds.) *Flora of Thailand* 14 (2). Forest Herbarium, Department of National Parks, Wildlife and Plant Conservation, pp. 185–251.
- Forster, J.R. & Forster, G. (1776) *Characteres generum plantarum ed. 2*. B. White, B., Cadell, T. & Elmsley, P., London, 150 pp.
- Gaertner, J. (1791) *De fructibus et seminibus plantarum* 2 (3). G.H. Schramm, Tübingen, pp. 353–504.
- Gamble, J.S. (1919) XII. Decades Kewenses. Plantarum novarum in herbario Horti Regii conservatarum. Decades XCII. & XCIII. *Bulletin of Miscellaneous Information* 1919: 221–233.
- Grushvitzky, I.V., Skortsova, N.T., Arkhangelsky, D.B. & Chistyakova, L.D. (1984) The species of the genus *Trevesia* (Araliaceae) in the flora of Vietnam. *Botanicheskii Zhurnal* 69: 1019–1029.
- Hayata, B. (1916) *Icones Plantarum Formosanarum* 6. Government of Formosa, Taihoku, pp. 1–168, pl. 1–20.
- Hoang, D.T., Chernomor, O., Von Haeseler, A., Minh, B.Q. & Vinh, L.S. (2018) UFBoot2: improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* 35: 518–522.
<https://doi.org/10.1093/molbev/msx281>
- Hoo, G. & Tseng, C.J. (1978) Araliaceae. In: Editorial Committee of FRPS (Ed.) *Flora Reipublicae Popularis Sinicae* 54. Science Press, Beijing, pp. 1–188.
- IUCN (2024) *Guidelines for using the IUCN Red List categories and criteria, version 16*. Prepared by the IUCN Standards and Petitions Committee. Available from: <https://www.iucnredlist.org/documents/RedListGuidelines.pdf> (accessed 8 April 2025)
- Jebb, M.H.P. (1998) A revision of the genus *Trevesia* (Araliaceae). *Glasra* 3: 85–113.
- Katoh, K., Rozewicki, J. & Yamada, K.D. (2019) MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20: 1160–1166.
<https://doi.org/10.1093/bib/bbx108>
- Koch, K.H.E. (1859a) Ueber einige Araliaceen. *Wochenschrift für Gärtnerei und Pflanzenkunde* 2: 66–71.
- Koch, K.H.E. (1859b) Die Araliaceen im Allgemeinen und Aufzählung der in den Gärten kultivierten Arten. *Wochenschrift für Gärtnerei und Pflanzenkunde* 2: 371–372.
- Lamarck, J.B.A.P. de M. (1788) *Encyclopédie méthodique, Botanique* 2. Panckoucke, Paris, 774 pp.
- Larsson, A. (2014) AliView: a fast and lightweight alignment viewer and editor for large datasets. *Bioinformatics* 30: 3276–3278.
- Letunic, I. & Bork, P. (2021) Interactive Tree of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation, *Nucleic Acids Research* 49: W293–W296.
<https://doi.org/10.1093/nar/gkab301>
- Li, H.L. (1942) The Araliaceae of China. *Sargentia* 2: 1–134.
- Li, H.L. (1949) A new genus of the Araliaceae. *Journal of the Arnold Arboretum* 30: 231–232.

- Li, R. & Wen, J. (2013) Phylogeny and biogeography of *Dendropanax* (Araliaceae), a genus disjunct between tropical/subtropical Asia and the Neotropics. *Systematic Botany* 38: 536–551.
<https://doi.org/10.2307/25065918>
- Li, R. & Wen, J. (2014) Phylogeny and biogeography of Asian *Schefflera* (Araliaceae) based on nuclear and plastid DNA sequence data. *Journal of Systematics and Evolution* 52: 397–539.
<https://doi.org/10.1111/jse.12052>
- Li, R. & Wen, J. (2016) Phylogeny and diversification of Chinese Araliaceae based on nuclear and plastid DNA sequence data. *Journal of Systematics and Evolution* 54 (4): 453–467.
<https://doi.org/10.1111/jse.12196>
- Lindley, J. (1825) *Gastonia palmata*. *The Botanical Register* 11: t. 894.
- Linnaeus, C. (1753) *Species Plantarum*. L. Salvius, Stockholm, 1200 pp.
- Lowry II, P.P. & Plunkett, G.M. (2020) Resurrection of the genus *Heptapleurum* for the Asian clade of species previously included in *Schefflera* (Araliaceae). *Novon* 28: 143–170.
<https://doi.org/10.3417/2020612>
- Makino, T. (1900) *Plantae Japonenses novae vel minus cognitae*. *The Botanical Magazine Tokyo* 14: 31–34.
- Maximowicz, C.J. (1859) *Primitiae florum Amurensis*. Kaiserliche Akademie der Wissenschaften, St. Petersburg, 504 pp.
- Merrill, E.D. (1924) New species of plants from Indo-China. *University of California Publications in Botany* 10: 423–430.
- Miquel, F.A.W. (1856) Araliacearum indicarum genera et species aliquot novae. *Bonplandia* 4: 137–139.
- Miquel, F.A.W. (1863) Araliaceae novae, adiecta aliarum specierum praesertim indicarum revisione. *Annales Musei Botanici Lugduno-Batavi* 1: 1–27.
- Newman, M.F., Ketphanh, S., Svengsuksa, B., Thomas, P., Sengdala, K., Lamxay, V. & Armstrong, K. (2007) *A checklist of the vascular plants of Lao PDR*. Royal Botanic Garden, Edinburgh, 400 pp.
- Newman, M.F., Pullan, M., Souladeth, P., Ketphanh, S., Svengsuksa, B., Thomas, P., Sengdala, K., Lamxay, V. & Armstrong, K. (2017–present) *A checklist of the vascular plants of Lao PDR*. Online database available at: <https://padme.rbge.org.uk/laos/> (accessed 2 April 2025)
- Nguyen, L.T., Schmidt, H.A., von Haeseler, A. & Minh, B.Q. (2015) IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* 32: 268–274.
<https://doi.org/10.1093/molbev/msu300>
- Philipson, W.R. (1970) Constant and variable features of the Araliaceae. *Botanical Journal of the Linnean Society* 63: 87–100.
- Plunkett, G.M., Lowry II, P.P., Fiaschi, P., Frodin, D.G. & Nicolas, A.N. (2019) Phylogeny, biogeography, and morphological evolution among and within the Neotropical and Asian clades of *Schefflera* (Araliaceae). *Taxon* 68: 1149–1385.
<https://doi.org/10.1002/tax.12177>
- Plunkett, G.M., Wen, J. & Lowry II, P.P. (2004) Intrafamilial classifications and characters in Araliaceae: Insights from the phylogenetic analysis of nuclear (ITS) and plastid (*trnL-trnF*) sequences data. *Plant Systematics and Evolution* 245: 1–39.
<https://doi.org/10.1007/s00606-003-0101-3>
- POWO (2025) *Plants of the World Online*. Facilitated by the Royal Botanic Gardens, Kew. Available from: <https://powo.science.kew.org/> (accessed 8 April 2025)
- Seemann, B. (1866) Revision of the natural order Hederaceae. *Journal of Botany, British and Foreign* 4: 293–299; 352–356.
- Shang, C.B. (1984) Le genre *Schefflera* (Araliacées) en Chine et Indochine. *Candollea* 39: 453–486.
<https://doi.org/10.5169/SEALS-879923>
- Shang, C.B. (1997) Espèces nouvelles d'Araliaceae asiatiques. *Adansonia, sér. 3* 19: 77–83.
- Shang, C.B. & Lowry II, P.P. (2007) Araliaceae. In: Wu, Z.Y., Raven, P.H. & Hong, D.Y. (Eds.) *Flora of China*. Science Press, Beijing and Missouri Botanical Garden Press, St. Louis, pp. 435–491.
- Souvannakhoummane, K., Phonepaseuth, P., Souladeth, P., Lanorsavanh, S., Tagane, S. & Tanaka, N. (2025) Studies on the Gesneriaceae in Laos II: *Lysionotus calcicola*, a new species from limestone karsts in North and Central Laos, supported by morphological and molecular evidence. *Nordic Journal of Botany* 2025: e04620.
<https://doi.org/10.1002/njb.04620>
- Tagane, S., Rueangruea, S., Suddee, S., Nagamasu, H., Naiki, A. & Toyama, H. (2017) *Heteropanax thaiensis*, a new species and *Schefflera wrayi* (Araliaceae), a new record to Thailand. *Thai Forest Bulletin (Botany)* 45: 99–104.
<https://doi.org/10.20531/tfb.2017.45.2.03>
- Tagane, S., Toyama, H., Chhang, P., Nagamasu, H. & Yahara, T. (2015) Flora of Bokor National Park, Cambodia I: Thirteen new species and one change in status. *Acta Phytotaxonomica et Geobotanica* 66: 95–135.
<http://dx.doi.org/10.18942/apg.KJ00010001424>
- Thunberg, C.P. (1780) *Cussonia*, novum plantae genus e promontorio bonae spei africes. *Nova Acta Regiae Societatis Scientiarum*

Upsaliensis 3: 210–212.

- Torrey, J. & Gray, A. (1840) *A Flora of North America*. Wiley & Putnam, New York etc., pp. 545–711.
- Valcarcel, V., Fiz-Palacios, O. & Wen, J. (2014) The origin of the early differentiation of *Ivies* (*Hedera* L.) and the radiation of the Asian Palmate group (Araliaceae). *Molecular Phylogenetics and Evolution* 70: 492–503.
<https://doi.org/10.1016/j.ympev.2013.10.016>
- Viguier, R. (1906) Recherches anatomiques sur la classification des Araliacées. *Annales des Sciences Naturelles, Botanique* 9 (4): 1–210.
- Viguier, R. (1909) Nouvelles recherches sur les Araliacées. *Annales des Sciences Naturelles, Botanique* 9 (9): 305–405.
- Visiani, R. de. (1842) Sopra La *Gastonia palmata* de W. Roxburgh proposta qual tipo di un nuovo genere nella famiglia delle Araliacee. *Memorie della Reale Accademia delle Scienze di Torino* 2 (4): 257–266.
- Wen, J., Lee, C., Lowry II, P.P. & Hiep, N.T. (2003) Inclusion of the Vietnamese endemic genus *Grushvitzkya* in *Brassaiopsis* (Araliaceae): evidence from nuclear ribosomal ITS and chloroplast *ndhF* sequences. *Botanical Journal of the Linnean Society* 142: 455–463.
<https://doi.org/10.1046/j.1095-8339.2003.00188.x>
- Wen, J. & Lowry II, P.P. (2006) New species and new combination in *Brassaiopsis* (Araliaceae) from Vietnam and southern China. *Adansonia sér.* 3 28: 181–190.
- Wen, J., Lôt, P.K., Hiep, N.T., Regalado, J.J., Averyanov, L.V. & Lee, C. (2007) An unusual new species of *Trevesia* from Vietnam and its implications on generic delimitation in Araliaceae. *Taxon* 56: 1261–1268.
<https://doi.org/10.2307/25065918>
- Wen, J., Plunkett, G.M., Mitchell, A.D. & Wagstaff, S.J. (2001) The evolution of Araliaceae: A phylogenetic analysis based on ITS sequences of nuclear ribosomal DNA. *Systematic Botany* 26: 144–167.
- Yi, T.S., Lowry II, P.P., Plunkett, G.M. & Wen, J. (2004) Chromosomal evolution in Araliaceae and close relatives. *Taxon* 53: 987–1005.

Appendix

TABLE S1. Voucher information and GenBank accession numbers for Araliaceae in this study

Taxa	Voucher	Country (Locality)	ITS	Names used on Sequence
<i>Aralia chinensis</i> L. ²	Wen 1136 (CS)	China (Hainan)	AF242256	
<i>Aralia finlaysoniana</i> (Wall. ex G. Don) Seem. ³	Wen 10571 (US)	China (Yunnan, Hekou)	GU054665	
<i>Aralia henryi</i> Harms ⁴	Wen 10640 (US)	China (Yunnan, Marlipo)	KC952338	
<i>Aralia racemosa</i> L. ³	Wen 7291 (US)	USA (Wisconsin, Menominee)	GU054615	
<i>Aralia stellata</i> (King) J. Wen ³	Maxwell 03-505 (US)	Thailand (Doichiang Dao, Doi Luang Valley)	GU054650	
<i>Aralia vietnamensis</i> Ha ³	Shui 81844 (US)	China (Yunnan, Lvchun)	GU054666	
<i>Brassaiopsis</i> cf. <i>ficifolia</i> Dunn ²	N.T. Vinh HAL 4581 (HN)	Vietnam (Thanh Hoa)	EF152169	
<i>Brassaiopsis ficifolia</i> Dunn	Wen 5933-14 (F)	Vietnam (Ha Giang)	AF551722	
<i>Brassaiopsis glomerulata</i> (Blume) Regel ²	Wen 5064 (F)	China (Yunnan)	AY256901	
<i>Brassaiopsis glomerulata</i> (Blume) Regel ⁵	Wen 5839 (US)	Vietnam (Lao Cai)	KF591483	
<i>Brassaiopsis glomerulata</i> (Blume) Regel ¹	Lowry 4839 (MO)	Vietnam	AY304789	
<i>Brassaiopsis grushvitzkyi</i> J. Wen, Lowry & H.T. Nguyen ²	Wen 5864-2 (F)	Vietnam (Ha Giang)	AF551728	
<i>Brassaiopsis hainla</i> (Buch.-Ham. ex D. Don) Seem. ¹	Wen 4905 (F)	Nepal (Godawari)	AY389028	
<i>Brassaiopsis hispida</i> Seem ²	Wen 5031(F)	China (Yunnan, Dulongjiang)	EF152170	
<i>Cussonia gamtoosensis</i> Strey ⁶	Philipson 5623 (MO)	South Africa	MK978526	
<i>Cussonia paniculata</i> Eckl. & Zeyh. ⁶	Philipson 5612 (MO)	South Africa	AY389032	
<i>Dendropanax sessiliflorus</i> (Standl. & A.C. Smith) A.C. Smith ²	Wen 7002 (F)	Costa Rica (Puntarenas Prov.)	EF152174	
<i>Dendropanax trifidus</i> (Thunb.) Makino ²	Wen 2461 (CS)	Japan (Tokyo)	AF242238	
<i>Eleutherococcus sessiliflorus</i> (Rupr. & Maxim.) S.Y. Hu ²	Wen 3131 (CS)	China (Jilin)	AF242227	
<i>Harmsioplanax ingens</i> Philipson	Wen 10749 (US)	Indonesia (Papua, Jayawijaja)	JX106275	
<i>Heptapleurum arboricola</i> Hayata	Plunkett 1958 (NY)	USA	MK978569	
<i>Heptapleurum arboricola</i> Hayata	Wen 9448 (US)	China (Taiwan, Nantou)	KC952364	
<i>Heptapleurum arboricola</i> Hayata	Wen 6295 (US)	China (Yunnan, Baoshan)	GU054626	<i>Schefflera arboricola</i> (Hayata) Merr. ³
<i>Heptapleurum bengalense</i> (Gamble) Esser	Wen 10531 (US)	China (Yunnan, Gejiu)	KC952345	<i>Schefflera bengalensis</i> Gamble ⁴
<i>Heptapleurum brevipedicellatum</i> (Harms) Lowry & G. M. Plunkett	Lowry 4931 (MO)	Vietnam	MK978574	<i>Schefflera brevipedicellata</i> Harms ⁶
<i>Heptapleurum brevipedicellatum</i> (Harms) Lowry & G. M. Plunkett	Lowry 6550 (MO)	China (Yunnan)	MK978575	<i>Schefflera brevipedicellata</i> Harms ⁶

.....continued on the next page

TABLE S1. (Continued)

Taxa	Voucher	Country (Locality)	ITS	Names used on Sequence
<i>Heptapleurum brevipedicellatum</i> (Harms) Lowry & G. M. Plunkett	Plunkett 1998 (NY)	China (Yunnan)	MK978573	<i>Schefflera brevipedicellata</i> Harms ⁶
<i>Heptapleurum delavayi</i> Franch.	Wen 5021 (F)	China (Sichuan)	EF152176	<i>Schefflera delavayi</i> (Franch.) Harms ²
<i>Heptapleurum heptaphyllum</i> (L.) Y. F. Deng	Wen 6165 (US)	Vietnam (Ninh Binh, Bavi)	GU054641	<i>Schefflera heptaphylla</i> (L.) Frodin ³
<i>Heptapleurum heptaphyllum</i> (L.) Y. F. Deng	Lowry 4941 (MO)	Vietnam	AY955459	<i>Schefflera heptaphylla</i> (L.) Frodin ⁶
<i>Heptapleurum heptaphyllum</i> (L.) Y. F. Deng	Wen 8452 (F)	China (Yunnan)	MK978612	<i>Schefflera heptaphylla</i> (L.) Frodin ⁶
<i>Heptapleurum heptaphyllum</i> (L.) Y. F. Deng	Plunkett 2050 (NY)	Vietnam	MK978611	<i>Schefflera heptaphylla</i> (L.) Frodin ⁶
<i>Heptapleurum karsticola</i> Kongx., Tak.Yamam. & Tagane, sp. nov.	Tagane <i>et al.</i> Z1201 (FOF)	Laos (Khammouane, Khounkham)	PV821477	
<i>Heptapleurum leucanthum</i> (R.Vig.) Y.F.Deng	Wen 7369 (US)	Thailand (Chiang Mai, cult.)	KC952367	<i>Schefflera leucantha</i> R. Vig. ⁴
<i>Heptapleurum leucanthum</i> (R.Vig.) Y.F.Deng	Wen 5028 (F)	China (Yunnan, Baoshan)	AY389060	<i>Schefflera yunnanensis</i> H. L. Li ¹
<i>Heptapleurum oblongum</i> (Craib) G. M. Plunkett & Lowry ⁶	Plunkett 2036 (NY)	Vietnam	MK978632	
<i>Heptapleurum pauciflorum</i> (R. Vig.) Y. F.Deng	Wen 10594 (US)	China (Yunnan, Hekou)	KC952342	<i>Schefflera pauciflora</i> R. Vig. ⁴
<i>Heptapleurum petelotii</i> (Merr.) G. M. Plunkett & Lowry	Wen 10918 (US)	Vietnam (Ninh Binh, Cuc Phuong)	KC952380	<i>Schefflera petelotii</i> Merr. ⁴
<i>Heptapleurum vidalianum</i> (C. B. Shang) G. M. Plunkett & Lowry	Wen 10948 (US)	Vietnam (GuangNam, Ba Na)	KC952377	<i>Schefflera vidaliana</i> C. B. Shang
<i>Schefflera candelabrum</i> Baill. ¹	Lowry 4726 (MO)	New Caledonia	AF229728	
<i>Heptapleurum wardii</i> (C. Marquand & Airy Shaw) G. M. Plunkett & Lowry	Wen 9224 (US)	China (Xizang, Linzhi)	KC952332	<i>Schefflera wardii</i> Marquand & Airy Shaw ⁴
<i>Heptapleurum</i> sp.	Shui 71007 (US)	China (Yunnan, Mile)	KC952370	<i>Schefflera</i> sp. ⁴
<i>Heteropanax fragrans</i> (Roxb.) Seem. ²	Wen 2213 (CS)	China (Guangdong)	AF242242	
<i>Heteropanax fragrans</i> (Roxb.) Seem. ⁴	Pathak 4442 (US)	India (Arunachal Pradesh)	KC952400	
<i>Oplopanax elatus</i> (Nakai) Nakai ³	Wen 5418-11 (US)	China (Jilin, Wusong)	GU054662	
<i>Osmoxylon novoguineense</i> (Scheff.) Becc ⁶	Lorence 8157 (PTBG)	Hawaii (cultivated)	AF229726	
<i>Osmoxylon orientale</i> (Guillaumin) B.C.Stone ⁶	Plunkett 1858 (NY)	Vanuatu	MK978543	
<i>Panax bipinnatifidus</i> Seem.	Li 714	China (Yunnan, Gongshan)	KU246152	
<i>Panax trifolius</i> L. ³	Wen 10099 (US)	USA (Maryland, Baltimore)	GU054701	
<i>Panax wangianus</i> S. C. Sun	Wen 12167	China	KU246151	

.....continued on the next page

TABLE S1. (Continued)

Taxa	Voucher	Country (Locality)	ITS	Names used on Sequence
<i>Raukaua anomalus</i> (Hook.) A.D.Mitch. ⁶	Mitchell CHR-527931 (CHR)	New Zealand	MK978544	
<i>Raukaua edgerleyi</i> (Kirk) Seem. ⁶	Plunkett 2191 (NY)	New Zealand	MK978545	
<i>Schefflera digitata</i> J. R. Forst. & G. Forst. ¹	Mitchell s.n. (CHR 485642)	cultivated, Christchurch, New Zealand	U63188	
<i>Schefflera pseudocandelabrum</i> R. Vig. ¹	Lowry 4795 (MO)	New Caledonia	AY389054	
<i>Schefflera vieillardii</i> Baill.	Lowry 4747 (MO)	New Caledonia	AY389059	
<i>Schefflera vitiensis</i> (A.Gray) Seem. ³	Lowry 5497 (MO)	Fiji	AY955489	
<i>Schefflera vitiensis</i> (A.Gray) Seem. ⁶	Plunkett 2245 (NY)	Fiji	MK978682	
<i>Seemannaralia gerrardii</i> (Seem.) R. Vig. ⁶	Philipson 5471 (MO)	South Africa	AY389062	
<i>Sinopanax formosanus</i> (Hayata) H. L. Li ³	Wen 9390 (US)	China (Taiwan, Kaohsiung)	GU054628	
<i>Tetrapanax papyrifer</i> (Hook.) K. Koch ³	Wen 10135 (US)	Indonesia (West Java)	GU054663	
<i>Tetrapanax papyrifer</i> (Hook.) K. Koch ⁴	Wen 9431 (US)	China (Taiwan, Renai)	KC952336	
<i>Trevesia lateospina</i> Jebb ²	Wen 7480 (F)	Thailand (Chiangmai)	EF152172	
<i>Trevesia lateospina</i> Jebb ³	Wen 7480-1 (US)	Thailand (Lampang, Muang Bahn)	GU054664	
<i>Trevesia laotica</i> Kongx., Soulad. & Tagane, sp. nov.	Tagane <i>et al.</i> Z1168 (FOF)	Laos (Bolikhamxai, Nam Kading NPA)	PV821478	
<i>Trevesia palmata</i> (Roxb. Ex Lindl.) Vis. ⁴	Wen 8460 (US)	China (Yunnan, Hekou)	KC952333	
<i>Trevesia palmata</i> (Roxb. exLindl.) Vis. ¹	Hao 901 (F)	China (Guangdong, Guangzhou)	AF242247	
<i>Trevesia palmata</i> (Roxb. Ex Lindl.) Vis.	Wen 5669 (US)	China (Yunnan)	KF591488	
<i>Trevesia palmata</i> (Roxb. Ex Lindl.) Vis.	Plunkett 1329 (WS)	cultivated, New York Botanical Garden	MK978685	
<i>Trevesia sphaerocarpa</i> Grushv. & Skvortsova ²	N.T. Vinh HAL 4582 (HN)	Vietnam (Thanh Hoa)	EF152173	
<i>Trevesia sundaica</i> Miq ²	Wen 6222 (F)	Indonesia	AF551732	
<i>Trevesia vietnamensis</i> J. Wen & P.K. Lôc ²	N.T. Vinh HAL4579 (HN)	Vietnam (Thanh Hoa)	EF152171	
Outgroup				
<i>Pittosporum illicioides</i> Makino ⁴	Wen 8486	China (Zhejiang, Anji)	JX106283	

Note: ¹Sequences of taxa from Plunkett *et al.* (2004); ²Wen *et al.* (2007); ³Li & Wen (2013); ⁴Li & Wen (2014); ⁵Valcarcel *et al.* (2014);

⁶Plunkett *et al.* (2019); bold taxa are new sequences generated from the present study.