



<https://doi.org/10.11646/phytotaxa.716.1.8>

**Erratum: Muto, M., Chiba, T., Tuji, A., Horie, Y. & Fujibayashi, M. (2025) A new littoral diatom genus, *Lyreneis* gen. nov. (Bacillariophyceae, Lyrellales, Lyrellaceae), from southwestern Japan” *Phytotaxa* 711 (1): 1–27.**

Please note the Introduction section/seventh line from the bottom:

sMorphological classification of diatom frustules is one of the most common methods used in diatom research but requires a high level of expertise and is sometimes difficult to complete (Round *et al.* 1990).

**Should read:**

Morphological classification of diatom frustules is one of the most common methods used in diatom research but requires a high level of expertise and is sometimes difficult to complete (Round *et al.* 1990).

Please note the Acknowledgement section/fourth line from the bottom:

We are grateful to Professor Zhi-Qiang Zhang, managing editor and founder of *Phytotaxa*, for permitting the use of several LM images (Figs. 48–50 and Supplemental materials 13 reproduced from Al-Handal *et al.* 2018).

**Should read:**

We are grateful to Professor Zhi-Qiang Zhang, managing editor and founder of *Phytotaxa*, for permitting the use of several LM images (Figs. 48–50 and Appendix 13 reproduced from Al-Handal *et al.* 2018).