



***Bomarea rinconii* (Alstroemeriaceae), a new species from the Talamanca Mountains in Chiriqui Province, Panama**

DANIEL ADOLFO CÁCERES GONZÁLEZ

0427 Estafeta Universitaria, David, Chiriquí, Panamá; UCH Herbarium of the Universidad Autónoma de Chiriquí, David Chiriquí, Panama and Herbarium Senckenbergianum, Frankfurt/Main (FR), Germany; Email: consultoria.caceres@gmail.com

Introduction

Bomarea Mirbel (1804: 71; Alstroemeriaceae) includes 122 species (Govaerts 2013) and is divided into four subgenera (Hofreiter & Tillich 2002): *Baccata* (five species), *Bomarea* (ca. seventy species), *Sphaerine* (twelve species) and *Wichuraea* (eighteen species). *Bomarea* is distributed from Mexico and the Greater Antilles to southern South America (Gereau 1994). The centre of distribution of the genus is along the Andes from Colombia to Bolivia (Hofreiter 2008).

The diversity of *Bomarea* species scientifically documented for Panama has not increased in the last two decades: from five species (two endemic) in the *Flora of Panama* of Woodson, Schery and collaborators (Killip 1945) to 12 species (four endemic) in *Flora Mesoamericana* (Gereau 1994) and 12 species (four endemic) in the recent *Catálogo de plantas vasculares de Panamá* (Correa *et al.* 2004). After the recent treatment of *Bomarea* subgen. *Bomarea* sect. *Multiflorae* by Hofreiter (2008), this number decreased to 11 species: *Bomarea acuminata* Baker (1887: 212) is a synonym of *B. andreana* Baker (1882: 205), and *B. caldasii* (Kunth, 1816: 283) Ascherson & Graebner (1906: 431) is synonymous with *B. multiflora* (Linneaus f., 1782: 207) Mirbel (1804: 72), but *B. costaricensis* Kränzlin (1913: 4) is newly reported for Panama. Of these 11 species, four are endemic to Panama and Costa Rica, and only *B. caudatisepala* Gereau (1989: 599) and *B. bracteolata* Gereau (1989: 601) are endemic to Panama (Hofreiter 2008).

No new taxa of *Bomarea* have been described from Panama since Gereau (1989). However, some areas of the country have not been sufficiently collected because difficult orography and landscape, principally in far western (Cordillera de Talamanca) and far eastern Panama.

During a botanical excursion made in 2006, a specimen of *Bomarea* from a cloud forest was collected in the Reserva Forestal Fortuna (Cordillera de Talamanca) in Chiriqui Province. Subsequently, herbarium collections of *Bomarea* from 12 herbaria (B, CR, FR, INB, MO, NY, PMA, SCZ, SEL, UCH, USJ and WU; abbreviations from Thiers 2013) were studied. The specimen did not match any of the known taxa of the genus and hence is described and illustrated here as a new species.

***Bomarea rinconii* Cáceres González, sp. nov. (Fig. 1)**

Type:—PANAMA. Chiriqui Prov.: Distr. Gualaca, Reserva Forestal Fortuna, cloud forest, elevation 1800 m, 8° 46' 12.54" N, 82° 11' 44.02" W, 4 April 2006, Cáceres González 2988 (holotype: UCH!; isotypes: UCH!, PMA!).

This new species is similar to *Bomarea caudatisepala*, but it is characterized by leaves that are 2.6–3.0 times longer than wide (ovate or ovate-lanceolate), involucre bracts that are generally trifid and equal in length, every sepal having a short horn (1.0–1.5 mm long) located 1–2 mm below the apex, inner tepals (4.6–4.9 cm long), stigma trifid (every lobule 2.5–3.0 mm long), whereas *B. caudatisepala* has a horn, but it is longer (7–10 mm long) and located 2–4 mm below the apex, and the involucre bracts are entire.






FIGURE 1. *Bomarea rinconii* A. Habit. B. Dry inflorescence. C. Flower, front view with a short horn (upper right corner). D. Flower. E. Trifold involucral bract. All from Cáceres González 2988. Photographs by Daniel A. Cáceres González.

Plant twining, stem slender up to 3 m long, 3.5–4.0 mm in diameter, glabrous. Leaves ovate or ovate-lanceolate, 7.8–11.0 × 5–6 cm, from 1.5–2.1 times longer than wide. Both leaf surfaces glabrous, acuminate apex; petiole 1.4–1.8 cm long, glabrous. Involucral bracts 5, glabrous, generally trifid, 1.3–1.9 × 0.25 cm at base and 0.70 cm at apex, lobules 1–2 mm long. Inflorescence an umbel, pedicels up to 6, unbranched, ebracteolate 15.7–20.0 cm long, pubescent with short scattered hairs (erect to slightly curved), 0.1–0.2 mm long, brown. Flowers actinomorphic, calyx with glandular pubescence, 0.01–0.05 mm long and yellowish; outer tepals 3, free, 4.4–4.6 × 0.4–0.5 cm at base to 1.1–1.3 cm at apex, dark orange to reddish orange abaxially and orange-yellowish adaxially, every sepal with a short horn 1.0–1.5 mm long located 1–2 mm below the apex, green to blackish; inner tepals 3, free, unguiculate, 4.6–4.9 cm long (up to 2–5 mm longer than sepals), 1.4–1.7 mm wide at base to 2.1 cm wide on the apex, which is retuse-mucronate to lightly rounded, orange yellowish abaxially, yellowish adaxially in the lower half and orange at the apex, mottled purple-brown on the upper two-thirds. Stamens 6, free, filaments 3.0–4.5 cm long, straight, shorter than petals, anthers oblong 3.8–5.0 × 1.5–2.0 mm. Ovary inferior, globose, 0.5–0.7 cm long × 0.5 cm diameter, densely glandular-pubescent (cream to slightly brownish), axial placentation. Style 1.3–1.6 cm, glandular pubescent on the base, stigma trifid with every lobule 2.5–3.0 mm long. Fruits and seeds unknown.

Etymology:—*Bomarea rinconii* is named in honour of Prof. Rafael Rincón Gómez (1947–), from the Escuela de Biología in the Universidad Autónoma de Chiriquí (UNACHI), who is considered one of the most knowledgeable contemporary botanists of western Panama. He contributed vastly to the collections of the UCH herbarium and is highly esteemed for his generous nature of sharing his knowledge with colleagues and students. During his career, he has been a man with a great passion for botany and insatiable curiosity about the vascular plants of Panama; he has played an important role in the education of many professionals at University of Panama and now at UNACHI.

Distribution:—*Bomarea rinconii* is so far only known from the Reserva Forestal Fortuna, Chiriqui Province, in the Cordillera de Talamanca, Panama.

Habitat and Ecology:—The new species grows in primary cloud forest at an elevation around 1800 m. The common associates are *Guzmania scandens* H.E.Luther & W.J.Kress (Bromeliaceae), *Sobralia* species (Orchidaceae), *Anthurium* species (Araceae), *Satyria* species (Ericaceae). It was observed flowering from April to May (end of the dry season).

Conservation status:—*Bomarea rinconii* has been collected only once in Panama. At this site, few individuals have been observed, and, hence, it is assumed to be rare. However, only few areas on the Reserva Forestal Fortuna have been explored, and there are many endemic species of plants in this protected area. Therefore, in the conservation assessment presented here (following the guidelines of the IUCN 2001), *Bomarea rinconii* is categorized as Critically Endangered (CR), represented by only two distribution records and based on the criteria B2bi–ii.

Discussion:—*Bomarea rinconii* belongs to the former section *Caldasianae* with inner tepals exceeding outer and inflorescence an umbel (Killip 1935), but this section is now included in section *Multiflorae* (Hofreiter 2008). In Panama, section *Multiflorae*, according to Hofreiter (2008), is represented by seven species: *Bomarea acutifolia* (Link & Otto, 1828: 57) Herbert (1837: 112), *B. andreana*, *B. caudatisepala*, *B. costaricensis*, *B. chiriquina* Killip (1945: 15), *B. hirsuta* (Kunth, 1816: 285) Herbert (1837: 114) and *B. suberecta* Gereau (1989: 599). Four of the seven species are similar to *B. rinconii* (*B. caudatisepala*, *B. suberecta*, *B. andreana* and *B. costaricensis*).

In *Bomarea rinconii* every sepal has a short horn 1.0–1.5 mm long, located 1–2 mm below the apex. A horn is also present in *B. caudatisepala*, but it is longer, 7–10 mm, and located 2–4 mm below the apex. Another important feature is that in *B. rinconii* the involucral bracts are trifid, whereas in *B. caudatisepala* they are entire. Leaves in *B. rinconii* (5–6 cm) are wider than in *B. caudatisepala* (2.7–4.5), from 1.5 to 2.1 times longer than wide in the former and 2.6 to 3.0 times longer than wide in the latter.

Lobules of the trifid involucral bracts are equal in length in *Bomarea rinconii*, but in *B. suberecta* the middle (central) lobe is longer than the outer (lateral) ones, as is evident from the holotype (*D'Arcy 6587*, MO!). In *B. rinconii* one of the involucral bracts is leaf-like, and in *B. suberecta* all involucral bracts are

equal. *Bomarea rinconii* has a twining habit, whereas *B. suberecta* is suberect to erect. *Bomarea rinconii* can be easily distinguished from *B. andreana* and *B. costaricensis* by the open inflorescence. *Bomarea rinconii* has a short horn, which is not present at all in *B. costaricensis* and *B. andreana*.

Acknowledgements

I thank Katharina Schulte, Franziska Merkel, Stefan Dressler and Mark Chase for language editing and comments; Autoridad Nacional del Ambiente (ANAM) and Enel Fortuna S. A. for the collecting permit; curators of the herbaria B, CR, FR, INB, MO, NY, PMA, SCZ, SEL, UCH, USJ and WU for access to their collections; and Consultoría Ambiental Cáceres for facilities.

References

- Ascherson, P.F.A & Graebner, K.O.R.P.P (1906) *Synopsis der Mitteleuropäischen Flora. Dritter Band. Monocotyledones (Liliiflorae [Liliaceae; Amaryllidaceae; Dioscoreaceae; Iridaceae], Scitamineae, Microspermae [Orchidaceae])*. Engelmann, Leipzig, 934 pp.
- Baker, J.G. (1882) On a collection of bomareas made by M.E. André in New Granada and Ecuador. *Journal of Botany, British and Foreign* 20: 201–206.
- Baker, J.G. (1887) Plantae Lehmannianae in Guatemala, Costa Rica et Columbia collectae. Liliaceae, Haemodoraceae, Amaryllidaceae, Dioscoreaceae, Iridaceae. *Botanische Jahrbücher für Systematik Pflanzengeschichte und Pflanzengeographie* 8: 208–215.
- Correa, M.D., Galdames, C. & de Stapf, M.S. (2004) *Catálogo de plantas vasculares de Panamá*. Editora Novo Art, Bogotá, 599 pp.
- Gereau, R.E. (1989) Three new species of *Bomarea* (Alstroemeriaceae) from Mesoamerica. *Annals of the Missouri Botanical Garden* 76: 598–601.
<http://dx.doi.org/10.2307/2399503>
- Gereau, R.E. (1994) *Alstroemeriaceae*. In: Davidse, G., Sousa, S.M. & Chater, O.A. (eds.) *Flora Mesoamericana, Alismataceae a Cyperaceae* 6. Universidad Nacional Autónoma de México, México, pp. 48–51.
- Govaerts, R. (2013) *Bomarea*. In: *World checklist of selected plant families*. Royal Botanic Gardens, Kew. Available from: <http://apps.kew.org/wcsp/qsearch.do> (accessed 17 April 2013).
- Herbert, W. (1837) *Amaryllidaceae*. Ridgway & Sons, London, 428 pp.
- Hofreiter, A. & Tillich, H.J. (2002) The delimitation, ecology, distribution and infrageneric subdivision of *Bomarea* Mirbel (Alstroemeriaceae). *Feddes Repertorium* 113: 528–544.
- Hofreiter, A. (2008) A revision of *Bomarea* subgenus *Bomarea* s.str. section *Multiflorae* (Alstroemeriaceae). *Systematic Botany* 33: 661–684.
<http://dx.doi.org/10.1600/036364408786500172>
- IUCN (2001) *The IUCN Red List categories and criteria*, version 3.1. IUCN Species Survival Commission. Gland, Switzerland and Cambridge, U.K. Available from: <http://www.iucnredlist.org/technical-documents/categories-and-criteria/2001-categories-criteria> (accessed 10 February 2013).
- Killip, E.P. (1935) New species of *Bomarea* from the Andes. *Journal of the Washington Academy of Science* 25: 370–377.
- Killip, E.P. (1945) *Bomarea*. In: Woodson, R.E. & Schery, R.W. (eds.) *Flora of Panama. Annals of the Missouri Botanical Garden* 32: 14–17.
- Kränzlin, F.W.L. (1913) Amaryllidaceae quaedam novae v. criticae. *Botanische Jahrbücher für Systematik Pflanzengeschichte und Pflanzengeographie* 50 (Beibl. 112): 1–7.
- Kunth, K.S. (1816) Amaryllidaceae. In: Humboldt, F.W.H.A. von, Bonpland, A.J.A. & Kunth, K.S. *Nova genera et species plantarum I*. Librairie Graeco-Latino-Germanicae, Paris, pp. 278–285.
- Link, J.H.F. & Otto, C.F. (1828) Amaryllidaceae. In: *Icones plantarum rariorum horti regii botanici berlinensis cum descriptionibus et colendi ratione* 5: 57–58.
- Linnaeus, C. von filius (1782) *Supplementum plantarum systematis vegetabilium editionis decimae tertiae, generum plantarum editiones sextae, et specierum plantarum editionis secundae*. Orphanotrophi, Braunschweig, 468 pp.
- Mirbel, C.F.B. (1804) Les bomares, *Bomarea*. In: Buffon, G.L.L. (ed.) *Histoire naturelle, générale et particulière des plantes* 9: 71–72.
- Thiers, B. (2013) *Index herbariorum: a global directory of public herbaria and associated staff*. New York Garden's Virtual Herbarium. Available from: <http://sweetgum.nybg.org/ih/herbarium.php?irn=174420> (accessed 15 February 2013).