



Two new names in *Clematis* (Ranunculaceae)

DANIEL FALCK¹ & SAMULI LEHTONEN²

¹*Paraisten kaupunki, Finland*

²*Herbarium, FI-20014 University of Turku, Finland, Email: samile@utu.fi*

Molecular studies (Miikeda *et al.* 2006, Xie *et al.* 2011) have established that the genus *Naravelia* Adanson (1763: 460) *nom. cons.* is nested within the genus *Clematis* Linnaeus (1753: 543). Most of the names treated under *Naravelia* have combinations in *Clematis* already. However, *N. siamensis* Craib and *N. pauciflora* H.Eichler have not been transferred to *Clematis*, apparently because of the existing homonyms. Here, we provide new names for them.

Clematis aureolata D.Falck & Lehtonen, *nom. nov.* Replaced name:—*Naravelia siamensis* Craib (1915: 419–420).

Type:—THAILAND Chiangmai Doi Sutep, in scrub jungle, 330 m, 9 November 1911, *Kerr 1527b* (lectotype K! [sheet no. 000498799], designated here; isolectotypes BM! [sheet no. 000559550], E! [sheet no. 00386280]).

Etymology:—The existence of name *Clematis siamensis* Drummond & Craib (1915: 420–421) blocks the direct transfer of *N. siamensis* to *Clematis*. The new name refers to the halo-like arrangement of the petaloid staminodes. The original collection indicated by Craib (1915) includes three duplicates, none of which were designated as holotype. Following the ICN art. 9.2. (McNeill *et al.* 2012), we designate the Kew specimen as lectotype.

Clematis horripilata D.Falck & Lehtonen, *nom. nov.* Replaced name:—*Naravelia pauciflora* Eichler (1958: 57).

Type:—PHILIPPINES Luzon Rizal Prov. Montalban, May 1904, *Loher 6775* (holotype K [sheet no. 000634886]).

Etymology:—The existence of name *Clematis pauciflora* Nuttall ex Torrey & Gray (1838: 9) blocks the direct transfer of *N. pauciflora* to *Clematis*. The new name refers to the general hairiness of this species.

References

- Adanson, M. (1763) *Familles des plantes*. II Partie. Vincent, Paris. 640 pp.
- Craib, W.G. (1915) Contributions to the flora of Siam. *Bulletin of Miscellaneous Information / Royal Botanic Gardens, Kew* 10: 419–433.
<http://dx.doi.org/10.2307/4113296>
- Eichler, H. (1958) Revision der Ranunculaceen Malesiens. *Bibliotheca Botanica* 124: 1–110.
- Linnaeus, C. (1753) *Species plantarum, exhibentes plantas rite cognitatas ad genera relatas, cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum systema sexuale digestas*. Vol. 1. Laurentii Salvii, Holmiae. 560 pp.
<http://dx.doi.org/10.5962/bhl.title.669>
- McNeill, J., Barrie, F.R., Buck, W.R., Demoulin, V., Greuter, W., Hawksworth, D.L., Herendeen, P.S., Knapp, S., Marhold, K., Prado, J., Prud'homme van Reine, W.F., Smith, G.F., Wiersema, J.H. & Turland, N.J. (2012) International Code of Nomenclature for algae, fungi and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011. *Regnum Vegetabile* 154: 1–240.
- Miikeda, O., Kita, K., Handa, T. & Yukawa, T. (2006) Phylogenetic relationships of *Clematis* (Ranunculaceae) based on chloroplast and nuclear DNA sequences. *Botanical Journal of the Linnean Society* 152: 153–168.
<http://dx.doi.org/10.1111/j.1095-8339.2006.00551.x>
- Torrey, J. & Gray, A. (1838) *A Flora of North America: containing abridged descriptions of all the known indigenous and naturalized plants growing north of Mexico; arranged according to the natural system*. Wiley & Putnam, New York. 711 pp.
<http://dx.doi.org/10.5962/bhl.title.9466>
- Xie, L., Wen, J. & Li, L.Q. (2011) Phylogenetic analyses of *Clematis* (Ranunculaceae) based on sequences of nuclear ribosomal ITS and three plastid regions. *Systematic Botany* 36: 907–921.
<http://dx.doi.org/10.1600/036364411x604921>