



## A revised delimitation of the rattan genus *Calamus* (Arecaceae)

WILLIAM J. BAKER

Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AB, United Kingdom, [w.baker@kew.org](mailto:w.baker@kew.org)

### Abstract

All available phylogenetic evidence indicates that the rattan genus *Calamus*, the largest of all palm (Arecaceae) genera, is non-monophyletic and that the four remaining genera of subtribe Calaminae (Calameae: Calamoideae), *Ceratolobus*, *Daemonorops*, *Pogonotium* and *Retispatha*, are nested within it. This issue has not yet been adequately addressed in palm classifications, with recent authors preferring to wait for further phylogenetic evidence before revising the limits of the genera. Here, an alternative solution is proposed that is both pragmatic and phylogenetically robust. An expanded *Calamus* is recognised into which *Ceratolobus*, *Daemonorops*, *Pogonotium* and *Retispatha* are subsumed. This broad generic concept, which includes ca. 520 species, has practical advantages as it is more clearly defined by morphological and anatomical characters, and resolves potential biases introduced to recent eco-evolutionary research on palms by the non-monophyly of critical genera. Future phylogenetic research may yet provide an alternative means of delimiting these genera, but the broad sense *Calamus* proposed here is a justifiable alternative that can be adopted immediately. Nomenclatural synopses transferring currently accepted species of *Ceratolobus*, *Daemonorops* and *Pogonotium* to *Calamus* are provided, including 70 new combinations and 12 replacement names.

**Key words:** lianas, nomenclature, Palmae, palms, rattans, South-East Asia, taxonomy

### Introduction

The rattan genus *Calamus* Linnaeus (1753: 325) is the largest of all palm genera, including over 400 species distributed in the Asia-Pacific region, with one species in Africa. In the current classification of palms (Dransfield *et al.* 2008), *Calamus* is placed in subtribe Calaminae (Calameae: Calamoideae) along with four other rattan genera, *Ceratolobus* Blume (in Schultes & Schultes 1830: lxxx), *Daemonorops* Blume (in Schultes & Schultes 1830: 1333), *Pogonotium* Dransfield (1980a: 763) and *Retispatha* Dransfield (1980b: 529). Subtribe Calaminae is a strongly supported monophyletic group, whereas all available molecular and morphological phylogenetic evidence indicates that *Calamus* is not, and that the four other genera in the subtribe are variously nested within it (Baker *et al.* 2000a, Baker *et al.* 2000b, c, Baker *et al.* 2009). These studies also indicate that *Daemonorops*, the second largest genus of Calaminae with ca. 110 species, may not be monophyletic. In the current classification of palms, Dransfield *et al.* (2008) highlighted these systematic issues, but followed Baker *et al.* (2000b) in maintaining the prevailing generic limits, while acknowledging that further phylogenetic research would lead to changes in the future.

While published phylogenetic studies lack sufficient sampling, resolution and support within Calaminae to permit the delimitation of multiple, monophyletic genera within the subtribe, an alternative taxonomic solution is available that is both phylogenetically robust and pragmatic for users of palm taxonomy. An expanded genus *Calamus*, within which *Ceratolobus*, *Daemonorops*, *Pogonotium* and *Retispatha* are subsumed, can be recognised and justified by strong support for its monophyly in all published studies (Baker *et al.* 1999, Baker *et al.* 2000a, Baker *et al.* 2000b, c, Baker *et al.* 2009). In fact, steps along this path have already been taken. In preparation for the publication of the current palm classification (Dransfield *et al.* 2005, 2008), Baker and Dransfield (2008) reduced the genus *Calospatha* (1911: 232), which was accepted in the previous classification of palms (Uhl & Dransfield 1987), into *Calamus*. More recently, Henderson and Floda (2015) placed *Retispatha* in synonymy with *Calamus*, based on published research (Baker *et al.* 1999, Baker *et al.* 2000a, Baker *et al.* 2000b, c) and their own unpublished molecular and morphological phylogenetic studies, which they report to be consistent with the works of Baker *et al.* They chose not to make further changes in the Calaminae, preferring to wait until future analyses are completed. However, the phylogenetic topologies published by Baker *et al.* (1999, 2000a, b, c) dictate that if *Retispatha* is reduced into *Calamus*, then the remaining

genera must be subsumed in the same way. The sinking of one genus, but not the others is an incomplete solution to the *Calamus* problem.

Here, a revised classification is proposed in which *Ceratolobus*, *Daemonorops*, *Pogonotium* and *Retispatha* are placed in synonymy with *Calamus*. This is presented as a stable, alternative and pragmatic taxonomic solution for this problematic group; *Calamus sensu lato* is likely to be robust to new findings and unlikely to require further amendments. Although this solution further enlarges the largest palm genus to ca. 520 species, there is no guarantee that a future phylogenetic classification that recognises more genera will be any more manageable. Indeed, available evidence indicates that while some major clades within Calaminae may be resolved that can be recognised as genera (Baker *et al.* 2000b), a proliferation of genera may be required to account for all lineages and to maintain monophyly of all groups. Far from improving the situation, this could result in a classification of Calaminae that is more complex and difficult to use. In such circumstances, a broader taxonomic solution is preferable, as has been adopted in many similar cases for other plant groups (e.g. Gardiner 2012, Gardiner *et al.* 2013). Importantly, *Calamus* is the subject of ongoing taxonomic monographic work (e.g. Baker & Dransfield 2014, Henderson & Floda 2015) and the revised delimitation of *Calamus* proposed here is a tenable solution that can be adopted now for this purpose. It also resolves issues in recent evolutionary studies in which inherent assumptions of generic monophyly have been violated in the case of the Calaminae (e.g. Couvreur *et al.* 2011, Kissling *et al.* 2012, Baker & Couvreur 2013a, b, Couvreur *et al.* 2014), potentially leading to biased results.

The revised classification of *Calamus* offers practical advantages over the current delimitation. While the five genera recognised by Dransfield *et al.* (2008) have a number of reliable distinguishing morphological features, inexpert users of palm taxonomy may find these difficult to interpret. Moreover, *Calamus sensu lato* Dransfield *et al.* (2008) is highly polymorphic and lacks unequivocal synapomorphies. In contrast, *Calamus sensu lato* is very well defined and more readily identified, even when sterile. The species are dioecious, pleoanthic rattans which bear a conspicuous knee-like swelling on the leaf sheath below the petiole base. The inflorescences are adnate to both the internode and leaf sheath above the axil of origin. Like all Calamoideae, flowers are borne on the rachilla in sympodial pairs (dyads) or derivatives thereof; in pistillate plants, the pair comprises a functional pistillate flower and a sterile staminate flower, whereas in staminate plants the dyad is reduced to a solitary staminate flower. Infrequently, exceptions to these features occur in some species scattered across *Calamus sensu lato*, such as the knee being absent in a number of stemless species and, exceptionally, climbing species (e.g. *Calamus dumetosus* (Beccari) Henderson & Floda [2015: 59]). There is also a small number of hapaxanthic species and some other species present alternative floral cluster arrangements. Taken together, however, these characters allow ready discrimination of *Calamus sensu lato* from the other rattan species in subfamily Calamoideae. The expanded concept of *Calamus* is also consistent with evidence from root (Seubert 1996) and leaf (Tomlinson *et al.* 2011) anatomical studies. Furthermore, *Calamus sensu lato* appears to be unique among climbing palm groups in its stem vascular anatomy, particularly with respect to xylem construction (Tomlinson *et al.* 2001, Fisher *et al.* 2002, Tomlinson & Spangler 2002, Tomlinson *et al.* 2011).

The expanded concept of *Calamus* introduces some redundancy to the hierarchy of palm classification (Dransfield *et al.* 2008) as *Calamus sensu lato* and Calaminae contain equivalent grouping information, though there are precedents for this elsewhere (e.g. Korthalsiinae, Pigafettinae). Future phylogenetic research may yet lead to an alternative classification within the subtribe, although this might be more effectively implemented as an infrageneric classification within *Calamus sensu lato*. By adopting this pragmatic taxonomic solution, rattan researchers will no longer be undermined by systematic problems and will be better placed to further the understanding of rattan diversity, its origins, evolution, properties, conservation and utilisation.

## Taxonomic treatment

Generic nomenclature and complete nomenclatural synopses for the transfer of *Daemonorops*, *Ceratolobus* and *Pogonotium* species to *Calamus* are provided below, the transfer of *Retispatha* having already been completed by Henderson and Floda (2015). Species level taxonomy is based on the latest accounts of these genera obtained from the World Checklist of Areaceae (Govaerts *et al.* 2014), updated as necessary. For the remaining species in *Calamus*, see Govaerts *et al.* (2014).

***Calamus*** L., Sp. Pl.: 325 (1753)

*Rotanga* Boehm., Defin. Gen. Pl.: 395 (1760).

*Rotang* Adans., Fam. Pl. 2: 24 (1763).

*Daemonorops* Blume in J.A.Schultes & J.H.Schultes, Syst. Veg. 7: 1333 (1830), *synon. nov.*

*Ceratolobus* Blume in J.A.Schultes & J.H.Schultes, Syst. Veg. 7: lxxx (1830), *synon. nov.*

*Palmijuncus* Rumph. ex Kuntze, Revis. Gen. Pl. 2: 731 (1891), *nom. illeg.*

*Zalacella* Becc., Ann. Roy. Bot. Gard. (Calcutta) 11(1): 496 (1908).

*Calospatha* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 232 (1911).

*Cornera* Furtado, Gard. Bull. Singapore 14: 518 (1955).

*Schizospatha* Furtado, Gard. Bull. Singapore 14: 525 (1955).

*Pogonotium* J.Dransf., Kew Bull. 34: 763 (1980), *synon. nov.*

*Retispatha* J.Dransf., Kew Bull. 34: 529 (1980).

### **Synopsis of *Daemonorops* species transferred to *Calamus***

***Calamus acamptostachys*** (Becc) W.J.Baker, *comb. nov.*

*Daemonorops acamptostachys* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 209 (1911).

***Calamus acehensis*** (Rustiami) W.J.Baker, *comb. nov.*

*Daemonorops acehensis* Rustiami, Gard. Bull. Singapore 54: 202 (2002).

***Calamus affinis*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops affinis* Becc., Leaf. Philipp. Bot. 8: 3042 (1919).

***Calamus angustifolius*** Griff., Calcutta J. Nat. Hist. 5: 89 (1845).

*Daemonorops angustifolia* (Griff.) Mart., Hist. Nat. Palm. 3: 327 (1853).

*Calamus hygrophilus* Griff., Palms Brit. E. Ind.: t. 213 C (1850). *Daemonorops hygrophila* (Griff.) Mart., Hist. Nat. Palm. 3: 328 (1853).

*Palmijuncus hygrophilus* (Griff.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

*Daemonorops carcharodon* Ridl., Mat. Fl. Malay. Penins. 2: 178 (1907).

*Daemonorops angustispatha* Furtado, Gard. Bull. Straits Settle. 9: 161 (1937).

***Calamus asteracanthus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops asteracantha* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 227 (1911).

***Calamus ater*** (J.Dransf.) W.J.Baker, *comb. nov.*

*Daemonorops atra* J.Dransf., Bot. J. Linn. Soc. 81: 15 (1980).

***Calamus aureus*** (Renuka & Vijayak) W.J.Baker, *comb. nov.*

*Daemonorops aurea* Renuka & Vijayak., Rheedeia 4: 122 (1994).

***Calamus banggiensis*** (J.Dransf.) W.J.Baker, *comb. nov.*

*Daemonorops banggiensis* J.Dransf., Kew Bull. 36: 813 (1982).

***Calamus begunii*** (Burret) W.J.Baker, *comb. nov.*

*Daemonorops begunii* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 11: 204 (1931).

***Calamus binnendijkii*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops binnendijkii* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 67 (1911).

***Calamus brachypodus*** W.J.Baker, *nom. nov.*

*Daemonorops brachystachys* Furtado, Gard. Bull. Straits Settlement. 8: 344 (1935), non *Calamus brachystachys* Becc., Rec. Bot. Surv. India 2: 215 (1902).

**Note:**—A replacement name is required as the name *Calamus brachystachys* Becc. has already been used.

***Calamus brevicaulis*** (A.J.Hend. & N.Q.Dung) W.J.Baker, *comb. nov.*

*Daemonorops brevicaulis* A.J.Hend. & N.Q.Dung, Phytotaxa 8: 28 (2010).

***Calamus calapparius*** Mart., Hist. Nat. Palm. 3: 209 (1838).

*Daemonorops calapparia* (Mart.) Blume, Rumphia 3: 7 (1847), *Palmijuncus calapparius* (Mart.) Kuntze, Revis. Gen. Pl. 2: 731 (1891).

*Calamus amboinensis* Miq., Verh. Kon. Akad. Wetensch., Afd. Natuurk. 11(5): 20 (1868). *Palmijuncus amboinensis* (Miq.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

***Calamus calicarpus*** Griff., Calcutta J. Nat. Hist. 5: 92 (1845).

*Daemonorops calicarpa* (Griff.) Mart., Hist. Nat. Palm. 3: 326 (1853), *Palmijuncus calicarpus* (Griff.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

*Calamus petiolaris* Griff., Calcutta J. Nat. Hist. 5: 93 (1845). *Daemonorops petiolaris* (Griff.) Mart., Hist. Nat. Palm. 3: 326 (1853).

*Palmijuncus petiolaris* (Griff.) Kuntze, Revis. Gen. Pl. 2: 732 (1891).

*Daemonorops microthamnus* Becc., Rec. Bot. Surv. India 2: 221 (1902).

***Calamus clemensianus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops clemensiana* Becc., Philipp. J. Sci., C 4: 636 (1909).

***Calamus collariferus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops collarifera* Becc., Rec. Bot. Surv. India 2: 227 (1902).

*Daemonorops acanthobola* Becc., Rec. Bot. Surv. India 2: 228 (1902).

***Calamus confusus*** (Furtado) W.J.Baker, *comb. nov.*

*Daemonorops confusa* Furtado, Gard. Bull. Straits Settlement. 8: 347 (1935).

***Calamus conspectus*** W.J.Baker, *nom. nov.*

*Daemonorops spectabilis* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 228 (1911), non *Calamus spectabilis* Blume, Rumphia 3: 55 (1847).

**Note:**—A replacement name is required as the name *Calamus spectabilis* Blume has already been used.

***Calamus crinitus*** (Blume) Miq., Nieuwe Verh. Eerste Kl. Kon. Ned. Inst. Wetensch. Amsterdam, ser. 3, 3: 6 (1850).

*Daemonorops crinita* Blume, Rumphia 2: viii (1838). *Palmijuncus crinitus* (Blume) Kuntze, Revis. Gen. Pl. 2: 733 (1891). *Rotang crinitus* (Blume) Baill., Hist. Pl. 13: 300 (1895).

*Calamus manicatus* Teijsm. & Binn. ex Miq., Fl. Ned. Ind. 3: 135 (1855). *Palmijuncus manicatus* (Teijsm. & Binn. ex Miq.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

***Calamus cristatus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops cristata* Becc., Nelle Forest. Borneo: 608 (1902).

*Daemonorops diversispina* Becc., Rec. Bot. Surv. India 2: 229 (1902).

***Calamus curranii*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops curranii* Becc., Philipp. J. Sci., C 2: 238 (1907).

***Calamus depressiusculus*** Miq. ex H.Wendl. in O.C.E.de Kerchove de Denterghem, Palmiers: 236 (1878).

*Palmijuncus depressiusculus* (Miq. ex H.Wendl.) Kuntze, Revis. Gen. Pl. 2: 733 (1891), *Daemonorops depressiuscula* (Miq. ex H.Wendl.) Becc., Rec. Bot. Surv. India 2: 226 (1902).

***Calamus didymophyllus*** (Becc.) Ridl., J. Straits Branch Roy. Asiat. Soc. 30: 221 (1897).

*Daemonorops didymophylla* Becc. in J.D.Hooker, Fl. Brit. India 6: 468 (1893).

*Calamus cochleatus* Miq., Verh. Kon. Akad. Wetensch., Afd. Natuurk. 11(5): 29 (1868), nom. nud. *Palmijuncus cochleatus* (Miq.) Kuntze, Revis. Gen. Pl. 2: 733 (1891), nom. inval.

*Daemonorops cochleata* Teijsm. & Binn. ex Miq., Verh. Kon. Akad. Wetensch., Afd. Natuurk. 11(5): 29 (1868), nom. nud.

*Daemonorops mattanensis* Becc., Nelle Forest. Borneo: 608 (1902).

*Daemonorops motleyi* Becc., Rec. Bot. Surv. India 2: 224 (1902).

***Calamus draco*** Willd., Sp. Pl. 2: 203 (1799).

*Daemonorops draco* (Willd.) Blume, Rumphia 2: viii (1838), *Palmijuncus draco* (Willd.) Kuntze, Revis. Gen. Pl. 2: 732 (1891).

*Calamus draconis* Oken, Allg. Naturgesch. 3(1): 648 (1841).

*Daemonorops propinqua* Becc. in J.D.Hooker, Fl. Brit. India 6: 467 (1893).

***Calamus dracunculus*** (Ridl.) W.J.Baker, *comb. nov.*

*Daemonorops dracunculus* Ridl., Bull. Misc. Inform. Kew 1926: 91 (1926).

***Calamus elongatus*** (Blume) Miq., Nieuwe Verh. Eerste Kl. Kon. Ned. Inst. Wetensch. Amsterdam, ser. 3, 3: 6 (1850).

*Daemonorops elongata* Blume, Rumphia 2: iii (1838), *Palmijuncus elongatus* (Blume) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

***Calamus eugenii*** W.J.Baker, *nom. nov.*

*Daemonorops poilanei* J.Dransf., Kew Bull. 56: 663 (2001), non *Calamus poilanei* Conrard, Notul. Syst. (Paris) 7: 28 (1938).

**Note:**—A replacement name is required as the name *Calamus poilanei* Conrard has already been used. Poilane's given name, Eugene, is used in the replacement epithet.

***Calamus fissus*** (Blume) Miq., Nieuwe Verh. Eerste Kl. Kon. Ned. Inst. Wetensch. Amsterdam, ser. 3, 3: 6 (1850).

*Daemonorops fissa* Blume, Rumphia 3: 17 (1847). *Palmijuncus fissus* (Blume) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

*Daemonorops cinnamomea* Schaedler, Hamburger Garten-Blumenzeitung 31: 161 (1875).

*Daemonorops fissa* var. *cinnamomea* Becc., Rec. Bot. Surv. India 2: 219 (1902), nom. nud.

*Daemonorops hallieriana* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 218 (1911).

***Calamus fissilis*** A.J.Hend., N.K.Ban & N.Q.Dung, Palms 52: 192 (2008).

*Daemonorops fissilis* (A.J.Hend., N.K.Ban & N.Q.Dung) A.J.Hend., Phytotaxa 8: 27 (2010).

***Calamus flexilis*** W.J.Baker, *nom. nov.*

*Daemonorops gracilis* Becc., Leaflet. Philipp. Bot. 8: 3044 (1919), non *Calamus gracilis* Roxb., Fl. Ind. ed. 1832, 3: 781 (1832).

**Note:**—A replacement name is required as the name *Calamus gracilis* Roxb. has already been used.

***Calamus forbesii*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops forbesii* Becc., Rec. Bot. Surv. India 2: 227 (1902).

***Calamus formicarius*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops formicaria* Becc., Nelle Forest. Borneo: 608 (1902).

***Calamus geniculatus*** Griff., Calcutta J. Nat. Hist. 5: 67 (1845).

*Daemonorops geniculata* (Griff.) Mart., Hist. Nat. Palm. 3: 329 (1853), *Palmijuncus geniculatus* (Griff.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

***Calamus gracilipes*** Miq., Verh. Kon. Akad. Wetensch., Afd. Natuurk. 11(5): 28 (1868).

*Palmijuncus gracilipes* (Miq.) Kuntze, Revis. Gen. Pl. 2: 733 (1891). *Daemonorops gracilipes* (Miq.) Becc., Rec. Bot. Surv. India 2: 225 (1902).

*Daemonorops accedens* var. *brevispatha* Blume, Rumphia 3: 13 (1838).

*Daemonorops longipes* Miq., Fl. Ned. Ind., Eerste Bijv.: 592 (1861), nom. illeg.

***Calamus grandis*** Griff., Calcutta J. Nat. Hist. 5: 84 (1845).

*Daemonorops grandis* (Griff.) Mart., Hist. Nat. Palm. 3: 327 (1853). *Palmijuncus grandis* (Griff.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

*Calamus intermedius* Griff., Calcutta J. Nat. Hist. 5: 86 (1845). *Daemonorops intermedia* (Griff.) Mart., Hist. Nat. Palm. 3: 327 (1853).

*Palmijuncus intermedius* (Griff.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

*Calamus acanthopis* Griff., Palms Brit. E. Ind.: t. 216 (1850).

*Daemonorops kirtong* Griff., Palms Brit. E. Ind.: 102 (1850).

*Daemonorops malaccensis* Mart., Hist. Nat. Palm. 3: 327 (1853). *Palmijuncus malaccensis* (Mart.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

*Acanthophoenix grandis* André, Rev. Hort. (Paris) 67: 250 (1895).

*Daemonorops grandis* var. *megacarpus* Furtado, Gard. Bull. Singapore 14: 67 (1953).

*Daemonorops laciniata* Furtado, Gard. Bull. Singapore 14: 75 (1953).

***Calamus hirsutus*** (Blume) Miq., Verh. Kon. Akad. Wetensch., Afd. Natuurk. 11(5): 28 (1868).

*Daemonorops hirsuta* Blume, Rumphia 2: t. 135 (1843). *Palmijuncus hirsutus* (Blume) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

*Calamus hystrix* Griff., Calcutta J. Nat. Hist. 5: 70 (1845). *Daemonorops hystrix* (Griff.) Mart., Hist. Nat. Palm. 3: 328 (1853). *Palmijuncus*

*hystrix* (Griff.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

*Daemonorops hystrix* var. *exulans* Becc., Rec. Bot. Surv. India 2: 224 (1902).

*Daemonorops hystrix* var. *minor* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 138 (1911).

***Calamus horridus*** (Burret) W.J.Baker, *comb. nov.*

*Daemonorops horrida* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 15: 193 (1940).

***Calamus ingens*** (J.Dransf.) W.J.Baker, *comb. nov.*

*Daemonorops ingens* J.Dransf., Bot. J. Linn. Soc. 81: 20 (1980).

***Calamus jenkinsianus*** Griff., Calcutta J. Nat. Hist. 5: 81 (1845).

*Daemonorops jenkinsiana* (Griff.) Mart., Hist. Nat. Palm. 3: 327 (1853), *Palmijuncus jenkinsianus* (Griff.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

*Calamus nutantiflorus* Griff., Calcutta J. Nat. Hist. 5: 79 (1845). *Daemonorops nutantiflora* (Griff.) Mart., Hist. Nat. Palm. 3: 326 (1853).

*Palmijuncus nutantiflorus* (Griff.) Kuntze, Revis. Gen. Pl. 2: 732 (1891).

*Calamus margaritae* Hance, J. Bot. 12: 266 (1874). *Palmijuncus margaritae* (Hance) Kuntze, Revis. Gen. Pl. 2: 733 (1891). *Daemonorops*

*margaritae* (Hance) Becc., Rec. Bot. Surv. India 2: 220 (1902).

*Daemonorops pierreana* Becc., Rec. Bot. Surv. India 2: 220 (1902).

*Daemonorops schmidtiana* Becc., Bot. Tidsskr. 29: 98 (1909).

*Daemonorops jenkinsiana* var. *tenasserimica* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 44 (1911).

***Calamus johndransfieldii*** W.J.Baker, *nom. nov.*

*Daemonorops dransfieldii* Rustiami, Gard. Bull. Singapore 54: 199 (2002), non *Calamus dransfieldii* Renuka, Kew Bull. 42: 433 (1987).

**Note:**—A replacement name is required as the name *Calamus dransfieldii* Renuka has already been used. Dransfield's given name, John, is incorporated into the replacement epithet.

***Calamus komsaryi*** (Maturb., J.Dransf. & Mogeia) W.J.Baker, *comb. nov.*

*Daemonorops komsaryi* Maturb., J.Dransf. & Mogeia, Phytotaxa 195: 297 (2015)

***Calamus korthalsii*** (Blume) Miq., Nieuwe Verh. Eerste Kl. Kon. Ned. Inst. Wetensch. Amsterdam, ser. 3, 3: 6 (1850).

*Daemonorops korthalsii* Blume, Rumphia 3: 23 (1847), *Palmijuncus korthalsii* (Blume) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

***Calamus kunstleri*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops kunstleri* Becc. in J.D.Hooker, Fl. Brit. India 6: 469 (1893).

*Daemonorops vagans* Becc. in J.D.Hooker, Fl. Brit. India 6: 469 (1893).

***Calamus kurzianus*** (Hook.f. ex Becc.) W.J.Baker, *comb. nov.*

*Daemonorops kurziana* Hook.f. ex Becc. in J.D.Hooker, Fl. Brit. India 6: 463 (1893).

***Calamus lamprolepis*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops lamprolepis* Becc., Rec. Bot. Surv. India 2: 223 (1902).

***Calamus leptopus*** Griff., Calcutta J. Nat. Hist. 5: 73 (1845).

*Daemonorops leptopus* (Griff.) Mart., Hist. Nat. Palm. 3: 329 (1853), *Palmijuncus leptopus* (Griff.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

*Daemonorops congesta* Ridl., Mat. Fl. Malay. Penins. 2: 179 (1907).

***Calamus lewisianus*** Griff., Calcutta J. Nat. Hist. 5: 87 (1845).

*Daemonorops lewisiana* (Griff.) Mart., Hist. Nat. Palm. 3: 327 (1853), *Palmijuncus lewisianus* (Griff.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

*Daemonorops pseudosepal* Becc. in J.D.Hooker, Fl. Brit. India 6: 465 (1893).

*Daemonorops tabacina* Becc. in J.D.Hooker, Fl. Brit. India 6: 466 (1893).

*Daemonorops bakauensis* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 220 (1911).

*Daemonorops curtisii* Furtado, Gard. Bull. Straits Settlem. 9: 164 (1937).

***Calamus loherianus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops loheriana* Becc., Philipp. J. Sci., C 4: 637 (1909).

***Calamus longibracteatus*** W.J.Baker, *nom. nov.*

*Daemonorops longispatha* Becc., Rec. Bot. Surv. India 2: 230 (1902), non *Calamus longispathus* Ridl., Mat. Fl. Malay. Penins. 2: 209 (1907).

**Note:**—A replacement name is required as the name *Calamus longispathus* Ridl. has already been used.

***Calamus longipes*** Griff., Calcutta J. Nat. Hist. 5: 68 (1845).

*Daemonorops longipes* (Griff.) Mart., Hist. Nat. Palm. 3: 329 (1853), *Rotang longipes* (Griff.) Baill., Hist. Pl. 13: 300 (1895).

*Calamus ramosissimus* Griff., Calcutta J. Nat. Hist. 5: 78 (1845), *Daemonorops ramosissima* (Griff.) Mart., Hist. Nat. Palm. 3: 330 (1853).

*Palmijuncus ramosissimus* (Griff.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

*Daemonorops stricta* Blume, Rumphia 3: 19 (1847), *Calamus strictus* (Blume) Miq., Verh. Kon. Akad. Wetensch., Afd. Natuurk. 11(5): 28 (1868).

*Daemonorops virescens* Becc. in G.H.Perkins & al., Fragm. Fl. Philipp. 1: 47 (1904).

*Daemonorops sabensis* Becc. ex Gibbs, J. Linn. Soc., Bot. 42: 169 (1914).

*Daemonorops calothyrsa* Furtado, Gard. Bull. Straits Settlem. 8: 345 (1935).

*Daemonorops longipedunculata* Furtado, Gard. Bull. Straits Settlem. 8: 353 (1935).

***Calamus longispinosus*** (Burret) W.J.Baker, *comb. nov.*

*Daemonorops longispinosa* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 15: 196 (1940).

***Calamus longistipes*** (Burret) W.J.Baker, *comb. nov.*

*Daemonorops longistipes* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 15: 798 (1943).

*Daemonorops pleioclada* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 15: 797 (1943).

***Calamus macrophyllus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops macrophylla* Becc. in J.D.Hooker, Fl. Brit. India 6: 470 (1893).

***Calamus macropterus*** Miq., Verh. Kon. Akad. Wetensch., Afd. Natuurk. 11(5): 19 (1868).

*Palmijuncus macropterus* (Miq.) Kuntze, Revis. Gen. Pl. 2: 733 (1891), *Daemonorops macroptera* (Miq.) Becc., Rec. Bot. Surv. India 2: 223 (1902).

***Calamus maculatus*** (J.Dransf.) W.J.Baker, *comb. nov.*

*Daemonorops maculata* J.Dransf., Kew Bull. 45: 76 (1990).

***Calamus manii*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops manii* Becc. in J.D.Hooker, Fl. Brit. India 6: 463 (1893).

***Calamus megalocarpus*** (Burret) W.J.Baker, *comb. nov.*

*Daemonorops megalocarpa* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 15: 194 (1940).

***Calamus melanochaetes*** (Blume) Miq., Verh. Kon. Akad. Wetensch., Afd. Natuurk. 11(5): 28 (1868).

*Daemonorops melanochaetes* Blume in J.J.Roemer & J.A.Schultes, Syst. Veg. 7: 1333 (1830).

*Daemonorops ornata* W.Bull, Gard. Chron., n.s., 3: 523 (1875).

*Daemonorops melanochaetes* var. *depressiglobus* Teijsm. & Binn. ex Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 50 (1911).

*Daemonorops melanochaetes* var. *macrocarpus* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 49 (1911).

*Daemonorops melanochaetes* var. *macrocymbus* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 49 (1911).

*Daemonorops melanochaetes* var. *microcarpus* Teijsm. & Binn. ex Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 48 (1911).

*Daemonorops melanochaetes* var. *padangensis* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 49 (1911).

*Daemonorops aruensis* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 50 (1911), *synon. nov.*

*Daemonorops javanica* Furtado, Gard. Bull. Straits Settlem. 9: 170 (1937).

**Note:**—*Daemonorops aruensis* Beccari (1911: 50) was described from a collection made by Moseley on the Challenger expedition. The type material at Kew indicates some uncertainty in its origins, stating that it was “apparently” collected in the Aru Islands in 1874. The presence of a species from *Daemonorops* section *Daemonorops* in that area seems unlikely, as other members of this section are largely restricted to the west of Wallace’s Line. The material corresponds closely to this more widespread west Malesian species, as acknowledged in the protologue and in annotations on the type by Beccari, and is thus synonymised here.

***Calamus micracanthus*** Griff., Calcutta J. Nat. Hist. 5: 62 (1845).

*Palmijuncus micracanthus* (Griff.) Kuntze, Revis. Gen. Pl. 2: 733 (1891). *Daemonorops micracantha* (Griff.) Becc. in J.D.Hooker, Fl. Brit. India 6: 467 (1893). *Rotang micracanthus* (Griff.) Baill., Hist. Pl. 13: 299 (1895).

*Daemonorops draconcella* Becc., Nelle Forest. Borneo: 608 (1902).

***Calamus microstachys*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops microstachys* Becc., Rec. Bot. Surv. India 2: 225 (1902).

***Calamus mirabilis*** Mart., Hist. Nat. Palm. 3: 213 (1838).

*Daemonorops mirabilis* (Mart.) Mart., Hist. Nat. Palm. 3: 329 (1853). *Palmijuncus mirabilis* (Mart.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

***Calamus mogleanus*** (Rustiami) W.J.Baker, *comb. nov.*

*Daemonorops mogleana* Rustiami, Reinwardtia 13: 25 (2009).

**Note:**—This new combination should not be confused with *Calamus mogleae* J.Dransf., Kew Bull. 55: 717 (2000).

***Calamus mollis*** Blanco, Fl. Filip.: 264 (1837).

*Palmijuncus mollis* (Blanco) Kuntze, Revis. Gen. Pl. 2: 733 (1891). *Daemonorops mollis* (Blanco) Merr., Sp. Blancoan.: 86 (1918).

*Daemonorops fusca* Mart., Hist. Nat. Palm. 3: 331 (1853).

*Daemonorops gaudichaudii* Mart., Hist. Nat. Palm. 3: 331 (1853). *Calamus gaudichaudii* (Mart.) H.Wendl. in O.C.E.de Kerchove de Denterghem, Palmiers: 236 (1878). *Palmijuncus gaudichaudii* (Mart.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

***Calamus mollispinus*** (J.Dransf.) W.J.Baker, *comb. nov.*

*Daemonorops mollispina* J.Dransf., Kew Bull. 56: 662 (2001).



***Calamus monticola*** Griff., *Calcutta J. Nat. Hist.* 5: 90 (1845).

*Daemonorops monticola* (Griff.) Mart., *Hist. Nat. Palm.* 3: 328 (1853). *Palmijuncus monticola* (Griff.) Kuntze, *Revis. Gen. Pl.* 2: 733 (1891).

***Calamus niger*** Willd., *Sp. Pl.* 2: 203 (1799).

*Daemonorops nigra* (Willd.) Blume, *Rumphia* 3: 5 (1847). *Palmijuncus niger* (Willd.) Kuntze, *Revis. Gen. Pl.* 2: 732 (1891). *Rotang niger* (Willd.) Baill., *Hist. Pl.* 13: 299 (1895).

***Calamus nuichuaensis*** A.J.Hend., N.K.Ban & N.Q.Dung, *Palms* 52: 196 (2008).

*Daemonorops nuichuaensis* (A.J.Hend., N.K.Ban & N.Q.Dung) A.J.Hend., *Phytotaxa* 8: 27 (2010).

***Calamus oblatum*** (J.Dransf.) W.J.Baker, *comb. nov.*

*Daemonorops oblata* J.Dransf., *Bot. J. Linn. Soc.* 81: 18 (1980).

***Calamus oblongus*** Reinw. ex Blume in J.J.Roemer & J.A.Schultes, *Syst. Veg.* 7: 1323 (1830).

*Daemonorops oblonga* (Reinw. ex Blume) Blume, *Rumphia* 3: 25 (1847).

*Calamus platyacanthos* Mart., *Hist. Nat. Palm.* 3: 206 (1838). *Daemonorops platyacanthos* (Mart.) Mart., *Hist. Nat. Palm.* 3 (ed. 2): 204 (1845). *Palmijuncus platyacanthos* (Mart.) Kuntze, *Revis. Gen. Pl.* 2: 732 (1891).

***Calamus ochrolepis*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops ochrolepis* Becc. in G.H.Perkins & al., *Fragm. Fl. Philipp.* 1: 47 (1904).

***Calamus ocreatus*** (A.J.Hend. & N.Q.Dung) W.J.Baker, *comb. nov.*

*Daemonorops ocreata* A.J.Hend. & N.Q.Dung, *Phytotaxa* 8: 29 (2010).

***Calamus oligolepis*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops oligolepis* Becc., *Leafl. Philipp. Bot.* 8: 3035 (1919).

***Calamus oligophyllus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops oligophylla* Becc. in J.D.Hooker, *Fl. Brit. India* 6: 467 (1893).

***Calamus oxycoccus*** W.J.Baker, *nom. nov.*

*Daemonorops oxycarpa* Becc., *Nelle Forest. Borneo*: 607 (1902), non *Calamus oxycarpus* Becc., *Ann. Roy. Bot. Gard. (Calcutta)* 11 (App.): 138 (1913).

**Note:**—A replacement name is required as the name *Calamus oxycarpus* Becc. has already been used.

***Calamus pachystris*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops pachystris* Becc., *Ann. Roy. Bot. Gard. (Calcutta)* 12(1): 217 (1911).

***Calamus palembanicus*** (Blume) Miq., *Verh. Kon. Akad. Wetensch., Afd. Natuurk.* 11(5): 29 (1868).

*Daemonorops palembanica* Blume, *Rumphia* 3: 20 (1847). *Palmijuncus palembanicus* (Blume) Kuntze, *Revis. Gen. Pl.* 2: 732 (1891). *Rotang palembanicus* (Blume) Baill., *Hist. Pl.* 13: 300 (1895).

***Calamus pannosus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops pannosa* Becc., *Leafl. Philipp. Bot.* 8: 3033 (1919).

***Calamus parvifructus*** W.J.Baker, *nom. nov.*

*Daemonorops microcarpa* Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 15: 195 (1940), non *Calamus microcarpus* Becc., *Rec. Bot. Surv. India* 2: 213 (1902).

**Note:**—A replacement name is required as the name *Calamus microcarpus* Becc. has already been used.

***Calamus pedicellaris*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops pedicellaris* Becc., Leafl. Philipp. Bot. 8: 3040 (1919).

***Calamus periacanthus*** (Miq.) Miq., Verh. Kon. Akad. Wetensch., Afd. Natuurk. 11(5): 28 (1868).

*Daemonorops periacantha* Miq., Fl. Ned. Ind., Eerste Bijv.: 593 (1861). *Palmijuncus periacanthus* (Miq.) Kuntze, Revis. Gen. Pl. 2: 732 (1891). *Rotang periacanthus* (Miq.) Baill., Hist. Pl. 13: 300 (1895).

*Daemonorops dissitophylla* Becc., Nelle Forest. Borneo: 608 (1902).

*Daemonorops florida* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 230 (1911).

***Calamus plagiocyclus*** (Burret) W.J.Baker, *comb. nov.*

*Daemonorops plagiocycla* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 15: 194 (1940).

***Calamus politus*** (Fernando) W.J.Baker, *comb. nov.*

*Daemonorops polita* Fernando, Gard. Bull. Singapore 41: 56 (1988 publ. 1989).

***Calamus pumilus*** (Van Valk.) W.J.Baker, *comb. nov.*

*Daemonorops pumila* Van Valk., Blumea 40: 465 (1995).

***Calamus rarispinosus*** (Renuka & Vijayak.) W.J.Baker, *comb. nov.*

*Daemonorops rarispinosa* Renuka & Vijayak., Rheedeia 4: 125 (1994).

***Calamus riedelianus*** Miq., Verh. Kon. Akad. Wetensch., Afd. Natuurk. 11(5): 18 (1868).

*Palmijuncus riedelianus* (Miq.) Kuntze, Revis. Gen. Pl. 2: 733 (1891). *Daemonorops riedeliana* (Miq.) Becc., Rec. Bot. Surv. India 2: 226 (1902).

***Calamus ruber*** Reinw. ex Mart., Hist. Nat. Palm. 3: 209 (1838).

*Daemonorops rubra* (Reinw. ex Mart.) Blume, Rumphia 3: 6 (1847). *Palmijuncus ruber* (Reinw. ex Mart.) Kuntze, Revis. Gen. Pl. 2: 732 (1891).

*Daemonorops accedens* Blume, Rumphia 2: viii (1838). *Calamus accedens* (Blume) Miq., Verh. Kon. Akad. Wetensch., Afd. Natuurk. 11(5): 28 (1868). *Palmijuncus accedens* (Blume) Kuntze, Revis. Gen. Pl. 2: 734 (1891). *Rotang accedens* (Blume) Baill., Hist. Pl. 13: 300 (1895).

***Calamus rutilis*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops rutilis* Becc., Rec. Bot. Surv. India 2: 230 (1902).

***Calamus rutilis* var. *rutilis***

*Daemonorops rutilis* var. *rutilis*

***Calamus rutilis* var. *acaulescens*** (J.Dransf.) W.J.Baker, *comb. nov.*

*Daemonorops rutilis* var. *acaulescens* J.Dransf., Bot. J. Linn. Soc. 81: 23 (1980).

***Calamus sabut*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops sabut* Becc. in J.D.Hooker, Fl. Brit. India 6: 469 (1893).

*Daemonorops annulata* Becc., Rec. Bot. Surv. India 2: 227 (1902).

*Daemonorops pseudomirabilis* Becc., Rec. Bot. Surv. India 2: 226 (1902).

*Daemonorops turbinata* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 225 (1911).

***Calamus sarasinorum*** (Warb. ex Becc.) W.J.Baker, *comb. nov.*

*Daemonorops sarasinorum* Warb. ex Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 100 (1911).

***Calamus scapigerus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops scapigera* Becc., Rec. Bot. Surv. India 2: 228 (1902).

*Daemonorops lasiospatha* Furtado, Gard. Bull. Straits Settlem. 8: 351 (1935).

*Calamus bifacialis* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 15: 809 (1943).

***Calamus schlechteri*** (Burret) W.J.Baker, *comb. nov.*

*Daemonorops schlechteri* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 15: 799 (1943).

***Calamus sedisspirituum*** (Rustiami, J.Dransf. & Fernando) W.J.Baker, *comb. nov.*

*Daemonorops sedisspirituum* Rustiami, J.Dransf. & Fernando, Kew Bull. 69, 9531: 3

***Calamus sekundurensis*** (Rustiami & Zumaidar) W.J.Baker, *comb. nov.*

*Daemonorops sekundurensis* Rustiami & Zumaidar, Floribunda 2: 198 (2005).

***Calamus sepal*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops sepal* Becc. in J.D.Hooker, Fl. Brit. India 6: 465 (1893).

*Daemonorops imbellis* Becc., Rec. Bot. Surv. India 2: 220 (1902).

*Daemonorops aciculata* Ridl., Mat. Fl. Malay. Penins. 2: 176 (1907).

*Daemonorops scortechinii* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 81 (1911).

*Daemonorops kiahii* Furtado, Gard. Bull. Singapore 14: 73 (1953).

*Daemonorops nurii* Furtado, Gard. Bull. Singapore 14: 85 (1953).

***Calamus serpentinus*** (J.Dransf.) W.J.Baker, *comb. nov.*

*Daemonorops serpentina* J.Dransf., Kew Bull. 36: 810 (1982).

***Calamus siberutensis*** (Rustiami) W.J.Baker, *comb. nov.*

*Daemonorops siberutensis* Rustiami, Kew Bull. 57: 729 (2002).

***Calamus singalanus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops singalana* Becc., Rec. Bot. Surv. India 2: 219 (1902).

***Calamus sparsiflorus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops sparsiflora* Becc., Rec. Bot. Surv. India 2: 224 (1902).

***Calamus stenophyllus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops stenophylla* Becc., Rec. Bot. Surv. India 2: 220 (1902).

***Calamus takanensis*** (Rustiami) W.J.Baker, *comb. nov.*

*Daemonorops takanensis* Rustiami, Reinwardtia 13: 26 (2009).

***Calamus treubianus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops treubiana* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 75 (1911).

***Calamus trichrous*** (Miq.) Miq., Verh. Kon. Akad. Wetensch., Afd. Natuurk. 11(5): 28 (1868).

*Daemonorops trichroa* Miq., Fl. Ned. Ind., Eerste Bijv.: 952 (1861). *Palmijuncus trichrous* (Miq.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

***Calamus unijugus*** (J.Dransf.) W.J.Baker, *comb. nov.*

*Daemonorops unijuga* J.Dransf., Bot. J. Linn. Soc. 81: 24 (1980).

***Calamus urdanetanus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops urdanetana* Becc., Leaf. Philipp. Bot. 8: 3038 (1919).

***Calamus uschdraweitianus*** (Burret) W.J.Baker, *comb. nov.*

*Daemonorops uschdraweitiana* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 15: 191 (1940).

***Calamus validus*** W.J.Baker, *nom. nov.*

*Daemonorops robusta* Warb. ex Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(1): 101 (1911), non *Calamus robustus* L.Linden & Rodigas, Ill. Hort. 40: 19 (1893).

**Note:**—A replacement name is required as the name *Calamus robustus* L.Linden & Rodigas has already been used.

***Calamus verticillaris*** Griff., Calcutta J. Nat. Hist. 5: 63 (1845).

*Daemonorops verticillaris* (Griff.) Mart., Hist. Nat. Palm. 3: 329 (1853). *Palmijuncus verticillaris* (Griff.) Kuntze, Revis. Gen. Pl. 2: 733 (1891).

*Daemonorops setigera* Ridl., Fl. Malay Penins. 5: 45 (1925).

*Daemonorops stipitata* Furtado, Gard. Bull. Singapore 14: 142 (1953).

*Daemonorops verticillaris* var. *straminea* Furtado, Gard. Bull. Singapore 14: 147 (1953).

***Calamus wrightmyoensis*** (Renuka & Vijayak.) W.J.Baker, *comb. nov.*

*Daemonorops wrightmyoensis* Renuka & Vijayak., Rheede 4: 125 (1994).

### **Synopsis of *Ceratolobus* species transferred to *Calamus***

***Calamus ceratolobus*** W.J.Baker, *nom. nov.*

*Ceratolobus discolor* Becc., Malesia 3: 63 (1886), non *Calamus discolor* Mart., Hist. Nat. Palm. 3: 212 (1838).

*Ceratolobus hallierianus* Becc. ex K.Heyne, Nutt. Pl. Ned.-Ind.: 93 (1913).

**Note:**—A replacement name is required as the name *Calamus discolor* Mart. has already been used.

***Calamus concolor*** (Blume) W.J.Baker, *comb. nov.*

*Ceratolobus concolor* Blume, Rumphia 2: 165 (1843). *Ceratogynum concolor* Pritz., Icon. Bot. Index: 245 (1854), orth. var.

***Calamus georgei*** W.J.Baker, *nom. nov.*

*Ceratolobus kingianus* Becc. & Hook.f. in J.D.Hooker, Fl. Brit. India 6: 477 (1893), non *Calamus kingianus* Becc., Ann. Roy. Bot. Gard. (Calcutta) 11(1): 197 (1908).

**Note:**—A replacement name is required as the name *Calamus kingianus* Becc. & Hook.f. has already been used. King's given name, George, is used in the replacement epithet.

***Calamus glaucescens*** D.Dietr., Syn. Pl. 2: 1064 (1840).

*Ceratolobus glaucescens* Blume in J.J.Roemer & J.A.Schultes, Syst. Veg. 7: 1334 (1830), non *Calamus glaucescens* Blume, Rumphia 3: 65 (1847), nom. illeg.

**Note:**—The name *Calamus glaucescens* Blume is illegitimate as a later homonym of *Calamus glaucescens* D.Dietr. and is a synonym of *Calamus caesius* Blume, Rumphia 3: 57 (1847).

***Calamus pseudoconcolor*** (J.Dransf.) W.J.Baker, *comb. nov.*

*Ceratolobus pseudoconcolor* J.Dransf., Kew Bull. 34: 19 (1979).

***Calamus subangulatus*** Miq., Fl. Ned. Ind., Eerste Bijv.: 594 (1861).

*Palmijuncus subangulatus* (Miq.) Kuntze, Revis. Gen. Pl. 2: 734 (1891). *Ceratolobus subangulatus* (Miq.) Becc., Ann. Roy. Bot. Gard. (Calcutta) 11(App.): iii (1913). *Ceratolobus laevigatus* var. *subangulatus* (Miq.) Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(2): 16 (1918).

*Ceratolobus laevigatus* var. *angustifolius* Becc. in J.D.Hooker, Fl. Brit. India 6: 477 (1893).

*Ceratolobus laevigatus* var. *borneensis* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(2): 16 (1918).

*Ceratolobus laevigatus* var. *divaricatus* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(2): 16 (1918).

*Ceratolobus laevigatus* var. *major* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(2): 16 (1918).

*Ceratolobus laevigatus* var. *regularis* Becc., Ann. Roy. Bot. Gard. (Calcutta) 12(2): 16 (1918).

## **Synopsis of Pogonotium species transferred to Calamus**

***Calamus moorei*** (J.Dransf.) W.J.Baker, *comb. nov.*

*Pogonotium moorei* J.Dransf., *Principes* 26: 174 (1982).

***Calamus pogonotium*** W.J.Baker, *nom. nov.*

*Pogonotium divaricatum* J.Dransf., *Kew Bull.* 34: 766 (1980), non *Calamus divaricatus* Becc., *Ann. Roy. Bot. Gard. (Calcutta)* 11(App.): 10 (1913).

**Note:**—A replacement name is required as the name *Calamus divaricatus* Becc. has already been used.

***Calamus ursinus*** (Becc.) W.J.Baker, *comb. nov.*

*Daemonorops ursina* Becc., *Rec. Bot. Surv. India* 2: 222 (1902). *Pogonotium ursinum* (Becc.) J.Dransf., *Kew Bull.* 34: 763 (1980).

## **Acknowledgements**

I am extremely grateful to Thomas Couvreur, John Dransfield, Wolf Eiserhardt, Lauren Gardiner, Rafaël Govaerts, Fred Stauffer and Tim Utteridge for invaluable discussions and thoughtful reviews of earlier drafts of this paper.

## **References**

- Baker, W.J. & Dransfield, J. (2008) *Calospatha* subsumed in *Calamus* (Arecaceae: Calamoideae). *Kew Bulletin* 63: 161–162.  
<http://dx.doi.org/10.1007/s12225-007-9007-5>
- Baker, W.J. & Couvreur, T.L.P. (2013a) Global biogeography and diversification of palms sheds light on the evolution of tropical lineages. I. Historical biogeography. *Journal of Biogeography* 40: 274–285.  
<http://dx.doi.org/10.1111/j.1365-2699.2012.02795.x>
- Baker, W.J. & Couvreur, T.L.P. (2013b) Global biogeography and diversification of palms sheds light on the evolution of tropical lineages. II. Diversification history and origin of regional assemblages. *Journal of Biogeography* 40: 286–298.  
<http://dx.doi.org/10.1111/j.1365-2699.2012.02794.x>
- Baker, W.J. & Dransfield, J. (2014) New rattans from New Guinea (*Calamus*, Arecaceae). *Phytotaxa* 163: 181–215.  
<http://dx.doi.org/10.11646/phytotaxa.163.4.1>
- Baker, W.J., Dransfield, J. & Hedderson, T.A. (2000a) Phylogeny, character evolution, and a new classification of the calamoid palms. *Systematic Botany* 25: 297–322.  
<http://dx.doi.org/10.2307/2666644>
- Baker, W.J., Hedderson, T.A. & Dransfield, J. (2000b) Molecular phylogenetics of *Calamus* (Palmae) and related rattan genera based on 5S nrDNA spacer sequence data. *Molecular Phylogenetics and Evolution* 14: 218–231.  
<http://dx.doi.org/10.1006/mpev.1999.0697>
- Baker, W.J., Hedderson, T.A. & Dransfield, J. (2000c) Molecular phylogenetics of subfamily Calamoideae (Palmae) based on nrDNA ITS and cpDNA rps16 intron sequence data. *Molecular Phylogenetics and Evolution* 14: 195–217.  
<http://dx.doi.org/10.1006/mpev.1999.0696>
- Baker, W.J., Dransfield, J., Harley, M.M. & Bruneau, A. (1999) Morphology and cladistic analysis of subfamily Calamoideae (Palmae). In: Henderson, A. & Borchsenius, F. (Eds.) *Evolution and classification of palms. Memoirs of the New York Botanical Garden* 83, pp. 307–324.
- Baker, W.J., Savolainen, V., Asmussen-Lange, C.B., Chase, M.W., Dransfield, J., Forest, F., Harley, M.M., Uhl, N.W. & Wilkinson, M. (2009) Complete generic-level phylogenetic analyses of palms (Arecaceae) with comparisons of supertree and supermatrix approaches. *Systematic Biology* 58: 240–256.  
<http://dx.doi.org/10.1093/sysbio/syp021>
- Beccari, O. (1911) Asiatic palms – Lepidocaryeae. Part 2. The species of *Daemonorops*. *Annals of the Royal Botanic Garden, Calcutta* 12: 1–237.
- Couvreur, T.L.P., Forest, F. & Baker, W.J. (2011) Origin and global diversification patterns of tropical rain forests: inferences from a complete genus-level phylogeny of palms. *BMC Biology* 9: 44.

<http://dx.doi.org/10.1186/1741-7007-9-44>

- Couvreur, T.L.P., Kissling, W.D., Condamine, F.L., Svenning, J.-C., Rowe, N.P. & Baker, W.J. (2014) Global diversification of a tropical plant growth form: environmental correlates and historical contingencies in climbing palms. *Frontiers in Genetics* 5: 452.  
<http://dx.doi.org/10.3389/fgene.2014.00452>
- Dransfield, J. (1980a) *Pogonotium* (Palmae: Lepidocaryoideae), a new genus related to *Daemonorops*. *Kew Bulletin* 34: 761–768.  
<http://dx.doi.org/10.2307/4119069>
- Dransfield, J. (1980b) *Retispatha*, a new Bornean rattan genus (Palmae: Lepidocaryoideae). *Kew Bulletin* 34: 529–536.
- Dransfield, J., Uhl, N.W., Asmussen, C.B., Baker, W.J., Harley, M.M. & Lewis, C.E. (2005) A new phylogenetic classification of the palm family, Arecaceae. *Kew Bulletin* 60: 559–569.
- Dransfield, J., Uhl, N.W., Asmussen, C.B., Baker, W.J., Harley, M.M. & Lewis, C.E. (2008) *Genera Palmarum - the evolution and classification of palms*. Royal Botanic Gardens, Kew, Richmond, 732 pp.
- Fisher, J.B., Tan, H.T.W. & Toh, L.P.L. (2002) Xylem of rattans: Vessel dimensions in climbing palms. *American Journal of Botany* 89: 196–202.  
<http://dx.doi.org/10.3732/ajb.89.2.196>
- Gardiner, L.M. (2012) New combinations in the genus *Vanda* (Orchidaceae). *Phytotaxa* 61: 47–54.
- Gardiner, L.M., Kocyan, A., Motes, M., Roberts, D.L. & Emerson, B.C. (2013) Molecular phylogenetics of *Vanda* and related genera (Orchidaceae). *Botanical Journal of the Linnean Society* 173: 549–572.  
<http://dx.doi.org/10.1111/boj.12102>
- Govaerts, R., Dransfield, J., Zona, S., Hodel, D.R. & Henderson, A. (2014) World Checklist of Arecaceae. Facilitated by the Royal Botanic Gardens, Kew. Available from: <http://apps.kew.org/wcsp/>.
- Henderson, A. & Floda, D. (2015) *Retispatha* subsumed in *Calamus* (Arecaceae). *Phytotaxa* 192: 58–60.  
<http://dx.doi.org/10.11646/phytotaxa.192.1.8>
- Kissling, W.D., Eiserhardt, W.L., Baker, W.J., Borchsenius, F., Couvreur, T.L.P., Balslev, H. & Svenning, J.C. (2012) Cenozoic imprints on the phylogenetic structure of palm species assemblages worldwide. *Proceedings of the National Academy of Sciences of the United States of America* 109: 7379–7384.  
<http://dx.doi.org/10.1073/pnas.1120467109>
- Linnaeus, C. (1753) *Species plantarum*. vol. 1–2. Salvius, Holmiae, 1200 pp.
- Schultes, J.A. & Schultes, J.H. (1830) *Systema Vegetabilium*. Vol.7. J.G. Cottae, Stuttgart, 1816 pp.
- Seubert, E. (1996) Root anatomy of palms. II. Calamoideae. *Feddes Repertorium* 1–2: 43–59.
- Tomlinson, P.B. & Spangler, R. (2002) Developmental features of the discontinuous stem vascular system in the rattan palm *Calamus* (Arecaceae-Calamoideae-Calamineae). *American Journal of Botany* 89: 1128–1141.  
<http://dx.doi.org/10.3732/ajb.89.7.1128>
- Tomlinson, P.B., Horn, J.W. & Fisher, J.B. (2011) *The Anatomy of Palms*. Oxford University Press, Oxford, 251 pp.
- Tomlinson, P.B., Fisher, J.B., Spangler, R.E. & Richer, R.A. (2001) Stem vascular architecture in the rattan palm *Calamus* (Arecaceae-Calamoideae-Calamineae). *American Journal of Botany* 88: 797–809.  
<http://dx.doi.org/10.2307/2657032>
- Uhl, N.W. & Dransfield, J. (1987) *Genera Palmarum, a classification of palms based on the work of Harold E. Moore Jr.* L. H. Bailey Hortorium and the International Palm Society, Lawrence, Kansas, 610 pp.