



## A new combination in *Helosciadium* (Apiaceae) for the flora of North Africa

ADRIANO STINCA\* & RICCARDO MOTTI

Department of Agriculture, University of Naples Federico II, via Università 100, 80055 Portici (Naples), Italy; [adriano.stinca@unina.it](mailto:adriano.stinca@unina.it)  
\*author for correspondence

According to recent molecular studies (Hardway *et al.* 2004, Spalik & Downie 2006, Spalik *et al.* 2009), the genus *Helosciadium* Koch (1824: 125) was segregated from *Apium* Linnaeus (1753: 264). The separate position of *Helosciadium* is supported also by morphological characters and fruit anatomy (Ronse *et al.* 2010).

Five taxonomic units of the genus *Helosciadium* are reported until now for the Euro-Mediterranean area (Hand 2011): *H. bermejoi* (Llorens 1982: 28) Popper & M.F. Watson in Ronse *et al.* (2010: 12), *H. crassipes* Koch (1824: 126), *H. inundatum* (Linnaeus 1753: 253) Koch (1824: 126), *H. nodiflorum* (Linnaeus 1753: 251) Koch (1824: 126), and *H. repens* (Jacquin 1775: 34) Koch (1824: 126). However, *Apium nodiflorum* (L.) Lagasca y Segura (1821: 101) subsp. *mairei* Molina Abril & Sardinero (1996: 278) was described as a subspecies endemic to Morocco and Algeria. For this unit, we propose here a new combination under *Helosciadium*.

*Helosciadium nodiflorum* (L.) W.D.J.Koch subsp. *mairei* (Molina Abril & Sard.Rosc.) Stinca & Motti **comb. nov.**

Basionym: *Apium nodiflorum* (L.) Lag. subsp. *mairei* Molina Abril & Sardinero (1996: 278). Type:—MOROCCO. S<sup>i</sup>. Yahja du Rharb, in loco dictio Daitaicha, en el desagüe de una fuente y en los bordes de la charca junto a ella, 15 m, 11 June 1993, J.A. Molina Abril s.n. (holotype MAF142760; isotypes BC, COI, G00359754 [digital image]!, G00359755 [digital image]!, K, MA611416 [digital image]!, MA628093 [digital image]!, MPU, P00644239 [digital image]!, SEV).

### References

- Hand, R. (2011) *Helosciadium* W.D.J.Koch. *Euro+Med Plantbase*. Available from: <http://ww2.bgbm.org/EuroPlusMed/PTaxonDetail.asp?NameCache=Helosciadium&PTRefFk=7500000> (accessed 26 December 2014)
- Hardway, T., Spalik, K., Watson, M.F., Katz-Downie, D.S. & Downie, S.R. (2004) Circumscription of Apiaceae tribe Oenantheae. *South African Journal of Botany* 70: 393–406.  
[http://dx.doi.org/10.1016/S0254-6299\(15\)30222-2](http://dx.doi.org/10.1016/S0254-6299(15)30222-2)
- Jacquin, N.J. (1775) *Florae Austriacae* 3. Leopoldi Joannis Kaliwoda, Viennae, 61 pp. + 100 plates.
- Koch, W.D.J. (1824) *Generum Tribuumque Plantarum Umbelliferarum Nova Dispositio. Nova Acta Physico-Medica Academiae Caesareae Leopoldino-Carolinae Naturae Curiosorum* 12: 55–156.
- Lagasca y Segura, M. (1821) *Amenidades Naturales de las Españas* 1 (2). Ibarra, Madrid, 111 pp.
- Linnaeus, C. (1753) *Species Plantarum* 1. Laurentii Salvii, Holmiae, 560 pp.
- Llorens, L. (1982) Un nuevo endemismo de la isla de Menorca: *Apium bermejoi*. *Folia Botanica Miscellanea* 3: 27–33.
- Molina Abril, J.A. & Sardinero, S. (1996) Una nueva subespecie de *Apium*: *A. nodiflorum* subsp. *mairei* (Apiaceae). *Lagascalía* 18: 276–279.
- Ronse, A.C., Popper, Z.A., Preston, J.C. & Watson, M.F. (2010) Taxonomic revision of European *Apium* L. s.l.: *Helosciadium* W.D.J.Koch restored. *Plant Systematics and Evolution* 287: 1–17.  
<http://dx.doi.org/10.1007/s00606-010-0284-3>
- Spalik, K. & Downie, S.R. (2006) The evolutionary history of *Sium* sensu lato (Apiaceae) dispersal, vicariance, and domestication as inferred from ITS rDNA phylogeny. *American Journal of Botany* 93: 747–761.  
<http://dx.doi.org/10.3732/ajb.93.5.747>
- Spalik, K., Downie, S.R. & Watson, M.F. (2009) Generic delimitations within the *Sium* alliance (Apiaceae tribe Oenantheae) inferred from cpDNA *rps16-5'trnK<sup>(UUU)</sup>* and nrDNA ITS sequences. *Taxon* 58: 735–748.