



Phytotaxa 106 (1): 1–171 (2013)
www.mapress.com/phytotaxa/
Copyright © 2013 Magnolia Press

Monograph

ISSN 1179-3155 (print edition)
PHYTOTAXA
ISSN 1179-3163 (online edition)



<http://dx.doi.org/10.11646/phytotaxa.106.1.1>

PHYTOTAXA

106

Nomenclator botanicus for the neotropical genus *Miconia* (Melastomataceae: Miconieae)

RENATO GOLDENBERG¹, FRANK ALMEDA², MAYARA K. CADDAH³, ANGELA B. MARTINS³,
JULIA MEIRELLES³, FABIAN A. MICHELANGELI⁴ & MARKUS WEISS⁵

¹ Universidade Federal do Paraná, Departamento de Botânica, Centro Politécnico, Caixa Postal 19031, Curitiba, PR, 81531-970,
Brazil; renato.goldenberg@gmail.com

² Department of Botany, California Academy of Sciences, 55 Music Concourse Drive, Golden Gate Park, San Francisco, CA 94118,
USA; falmeda@calacademy.org

³ Universidade Estadual de Campinas, Pós-Graduação em Biologia Vegetal, Cidade Universitária Zeferino Vaz, 13083-970,
Campinas, São Paulo, Brazil.

⁴ The New York Botanical Garden, 2900 Southern Blvd, Bronx, NY 10458, USA.

⁵ Information Technology Center, Staatliche Naturwissenschaftliche Sammlungen Bayerns, Menzinger Straße 67, 80638 München,
Germany.



Magnolia Press
Auckland, New Zealand

Renato Goldenberg, Frank Almeda, Mayara K. Caddah, Angela B. Martins, Julia Meirelles,
Fabian A. Michelangeli & Markus Weiss

Nomenclator botanicus for the neotropical genus *Miconia* (Melastomataceae: Miconieae)

(*Phytotaxa* 106)

171 pp.; 30 cm.

6 June 2013

ISBN 978-1-77557-192-6 (paperback)

ISBN 978-1-77557-193-3 (Online edition)

FIRST PUBLISHED IN 2013 BY

Magnolia Press

P.O. Box 41-383

Auckland 1346

New Zealand

e-mail: magnolia@mapress.com

<http://www.mapress.com/phytotaxa/>

© 2013 Magnolia Press

All rights reserved.

No part of this publication may be reproduced, stored, transmitted or disseminated, in any form, or by any means, without prior written permission from the publisher, to whom all requests to reproduce copyright material should be directed in writing.

This authorization does not extend to any other kind of copying, by any means, in any form, and for any purpose other than private research use.

ISSN 1179-3155 (Print edition)

ISSN 1179-3163 (Online edition)

Table of contents

| | |
|---|-----|
| Abstract | 3 |
| Introduction | 3 |
| Acknowledgements | 7 |
| References | 7 |
| LIST 1. "Nomenclator botanicus" for all published names of <i>Miconia</i> | 11 |
| LIST 2. Species of <i>Miconia</i> organized by section | 129 |
| LIST 3. Species of <i>Miconia</i> by country, with respective vouchers..... | 140 |
| LIST 4. <i>Miconia</i> names synonymized under other genera | 168 |

Abstract

Miconia has 1057 species and is the largest genus of woody flowering plants with a distribution restricted to tropical America. These species occur from Mexico (79 spp) to Argentina (10) and Uruguay (1), and attain highest richness in Colombia (338 spp), Peru (302), Brazil (274), Ecuador (254) and Venezuela (212). More than half of the species belong to only three sections (*Miconia* with 245, *Cremanium* with 223, and *Amblyarrhena* with 212). The other sections are *Chaenopleura* (86 spp), *Glossocentrum* (82), *Tamonea* (71), *Octomeris* (59), *Jucunda* (23), *Laceraria* (19), *Chaenanthera* (17), *Hartigia* (7), and *Adenodesma* (6), while seven species have not been assigned to any section. In this nomenclator we attempt to enumerate all currently accepted species of *Miconia* with place of publication, information on types, synonymy, sectional placement, and distributional notes organized by country.

Key words: geographical distribution, megadiverse genus, nomenclature, taxonomy

Introduction

Miconia Ruiz & Pav. (1794: 60) is the largest genus of woody flowering plants with a distribution restricted to tropical America (Goldenberg 2000; Goldenberg et al. 2008). The genus includes just under one quarter of the total 5000 + species of Melastomataceae, which are largely centered in the tropical and subtropical regions of the world (Clausing & Renner 2001). As a generic name, *Miconia* has been conserved (Farr et al. 1979) against *Tamonea* Aublet (1775: 411), *Leonicenia* Scopoli (1777: 212), *Lieutatia* Buchoz (1779: 10), and *Zulatia* Necker (1790: 117; see Briquet 1906, Camp. et al. 1947, and subsequent codes). *Fothergilla* Aublet (1775:411) is an older name, but it is illegitimate as it was also preoccupied by *Fothergilla* Murray (1774: 418; see Rickett & Stafleu 1960, Williams 1963).

The last taxonomic revision of this genus was published in the late 19th century (Cogniaux 1891). That monumental account of the Melastomataceae treated only half of the species accepted here. Since then, *Miconia* has been treated mostly in regional Floras (Wurdack 1962, Wurdack 1973, Wurdack 1980, Wurdack et al. 1993, Almeda 2001, Almeda et al. 2007, Almeda 2009, Goldenberg 2009 and others) which concentrated on the delimitation and identification of large numbers of species. The very nature of these studies precluded any attempt to resolve problems involving generic circumscription and infrageneric classification. Consequently, Cogniaux (1891) was the last specialist who attempted to provide an infrageneric classification of *Miconia*. He divided the genus into 11 sections, based mostly on stamen morphology, but also on other features of the calyx and leaf morphology. Information on sectional placement of the 500 species described since Cogniaux's monograph has not always appeared in the protalogues of newly described species and in many cases no subsequent placement has been proposed. Distributional information for each species is also scattered despite the ongoing publication of regional floras. Each collecting expedition to the tropics results in the discovery of new taxa and expands the ranges of others. In this nomenclator we attempt to enumerate all currently accepted species of *Miconia* with place of publication, information on types, synonymy, sectional placement, and distributional notes organized by country. Detailed data on phylogeny, economic uses, ecology and taxonomic history of *Miconia* can be found in Goldenberg et al. (2008).

Miconia has been regarded as “non-natural” since the 19th century (see Judd & Skean 1991 and Goldenberg et al. 2008) and, more recently, has been shown to be polyphyletic based on morphological and molecular data sets (Michelangeli et al. 2004, Goldenberg et al. 2008). The relationships between *Miconia* and other genera in the tribe Miconieae are still unclear. There is a growing realization among students of the Miconieae about the eventual need to recognize a single mega-genus *Miconia* (Michelangeli et al., in prep.) to best reflect current understanding of phylogenetic relationships of this megadiverse clade (Ionta et al. 2012). Nevertheless, the alternative (i.e. splitting the large genera, like *Miconia*, *Leandra* Raddi and *Clidemia* D.Don) into several smaller ones cannot be completely ruled out, but this approach would be fraught with difficulties and inconsistencies. The sections of *Miconia* also seem to be non-monophyletic when treated as isolated entities, albeit groups of sections can be recognized in some clades (Goldenberg et al. 2008, Judd 2007).

Whatever new circumscription of the Miconieae is offered in the future, we find it timely and appropriate to publish a critical compendium on nomenclature, sectional placement and distribution of *Miconia* as it has traditionally been defined. This information will provide an indispensable tool for any forthcoming taxonomic realignments.

Methods

All data on nomenclature, sectional placement and distribution were organized in a Microsoft Access database, and later migrated to Diversity Workbench (Diversity Taxon Names 2011), an online application based at the Botanische Staatssammlung Munich, in Germany. The information on the nomenclatural aspects developed for *Miconia*, as well as for the rest of the family, is already available at <http://www.melastomataceae.net>.

The nomenclatural data assembled here are based on *Index Kewensis* and *Gray Herbarium Card Index* (both available at <http://www.ipni.org>), and Tropicos (<http://www.tropicos.org>), with later additions of recently described species. Authors and original publications were checked for each species and corrected when necessary. Validly published species and synonyms were extracted from the last monograph for the genus (Cogniaux 1891), then complemented with all new taxa and nomenclatural changes published subsequently through mid-2012 (see Goldenberg et al. 2010).

Information on the types is given only for accepted species. This is based on the original publications and checked whenever possible in several herbaria with important historical collections of Melastomataceae and in relevant bibliographic references (Howard & Kellogg 1986, Martin & Cremers 2007, Goldenberg & Martin 2008, Goldenberg et al. 2010, Michelangeli & Goldenberg 2012). We also made extensive use of web-based resources such as the Global Plants Initiative Project (GPI) (<http://plants.jstor.org>), and the databases of specific herbaria that are not completely available through GPI, such as BRS (<http://www.bergianska.se/>), G (<http://www.ville-ge.ch/cjb/>), and P (<http://coldb.mnhn.fr/colweb>). Some types that were not located are marked with a question mark. All lectotypes and neotypes noted here were designated in previous publications. We chose not to designate lectotypes or neotypes in this paper. Therefore, only syntypes are cited for several species.

Each valid species was assigned to a section, following the classification proposed by Cogniaux (1891), with the addition of section *Hartigia* (Miq.) Griseb. (Goldenberg 2000), which was combined by Cogniaux with section *Amblyarrhena* (Naudin) Triana ex Hook.f. The other sections are *Adenodesma* (Naudin) Cogn., *Cremanium* (D.Don) Triana ex Hook.f., *Chaenanthera* (Naudin) Cogn., *Chaenopleura* (Rich ex. DC.) Triana ex Hook.f., *Glossocentrum* (Crueger) Triana ex Hook.f., *Jucunda* (Cham.) Triana ex Hook.f., *Laceraria* (Naudin) Triana ex Hook.f., *Miconia* DC., *Octomeris* (Naudin) Triana ex Hook.f., and *Tamonea* Cogn. Sectional assignments follow Cogniaux (1891). We followed the judgment of original authors for all subsequently described species and where possible we made every attempt to place species that had been unassigned to section in the protogues. In a few cases we reassigned species to sections when the original was flagrantly wrong. Changes by subsequent authors were accepted whenever properly published (i.e. unpublished manuscripts and notes on herbarium sheets were not accepted).

The distribution of each species was established based on their occurrence in the western hemisphere countries, from Mexico to Argentina, including the Caribbean. We also list records for invasive plants found outside the Americas. Due to its large size, Brazil is divided into five regions following the Instituto Brasileiro de Geografia e Estatística (North, Northeast, West-Central, Southeast and South). Each distributional record was based on a flora or checklist compiled by family specialists (listed in the introduction, plus Duss 1896; Cogniaux 1913; Riley 1925; Fawcett & Rendle 1926; Standley 1931; Williams 1934; Gleason 1935, 1940, 1958; Standley & Record 1936; Macbride 1941; Johnston 1949; Liogier 1957; Standley & Williams 1963; Wurdack 1963, 1970, 1971a, 1971b, 1971c, 1976, 1993; Winkler 1965; Proctor 1972; Uribe-Uribe 1972; Molina 1975; Baumgratz 1980, 1982, 1984; Seymour 1980; Liogier & Martorell 1982; Bremer 1987; Howard 1989; Nicolson 1991; Baumgratz et al. 1995; Martins et al. 1996; Meyer 1996; Renner 1999; Liogier 2000; Berry et al. 2001; Goldenberg 2004; Ulloa-Ulloa et al. 2004; Ulloa-Ulloa & Neill 2005; Cotton & Michelangeli 2007; Funk et al. 2007; Csurhes 2008; Michelangeli & Cotton 2008; Neill & Ulloa-Ulloa 2011), and also confirmed by voucher specimens examined in several herbaria, mostly CAS, COL, F, MO, NY, UEC, UPCB and US. Our decision to accept distributional records was certainly subjective in some cases. We adopted a conservative approach. If we were not certain about a species report in a Flora or had doubts about herbarium specimen determinations, we typically opted not to accept such records. Each species record is linked to a voucher in one of the above cited herbaria or to a bibliographic reference citing it.

From this database, we generated the following lists: (1) a “nomenclator botanicus” for all the names of *Miconia*, including the actual species and synonyms, information on the types for the accepted species, sectional placement and distribution by country; (2) a list with all the species of *Miconia* organized by section; (3) a list with all species of *Miconia* by country, with respective vouchers; (4) a list with *Miconia* names synonymized under other genera.

Results and Discussion

As traditionally circumscribed, *Miconia* has 1057 species (List 1). About two thirds of the species belong to only three sections (*Miconia* with 245, *Cremanium* with 223, and *Amblyarrhena* with 212; List 2). The other sections, ordered by richness, are *Chaenoplectrona* (86 spp), *Glossocentrum* (82), *Tamonea* (71), *Octomeris* (59), *Jucunda* (23), *Laceraria* (19), *Chaenanthera* (17), *Hartigia* (7), and *Adenodesma* (6). Seven species have not been assigned to any section.

The genus occurs from Mexico (79 spp) to Argentina (10) and Uruguay (1). It attains highest richness in Colombia (338 spp), then Peru (302), Brazil (274), Ecuador (254), and Venezuela (212; List 3).

Species richness in the genus is concentrated along the Andes, but it is also relatively high in the Guayana Highlands, Southeastern Brazil, Greater Antilles and Central America (Table 1). The composition of the floras in these regions is rather distinct, when considering the sections. In Colombia, Ecuador, Peru, Bolivia and Andean Venezuela, the genus *Miconia* is strongly represented by species from the three large sections (*Miconia*, *Cremanium* and *Amblyarrhena*), with section *Miconia* present mostly below 1500 m and *Cremanium* and *Amblyarrhena* present mostly above 1800 m. In the Guianas, non-Andean Venezuela and Northern Brazil, the number of species from sections *Cremanium* and *Amblyarrhena* is much smaller, whereas the proportion of species from section *Miconia* is higher. In Southeastern and Southern Brazil, the richest section is *Glossocentrum*, whereas in the Greater Antilles it is section *Chaenoplectrona*.

While some sections seem to be concentrated in restricted areas, others are more widespread. *Octomeris* is more common in Central America, Colombia and Venezuela than in other countries, being absent in the Guianas and parts of Brazil. *Glossocentrum* and *Chaenanthera* are more common in Southeastern Brazil, but their species occur throughout Tropical America. *Amblyarrhena* is almost absent outside of the Andes from Colombia to Bolivia and also Central America. The distribution of *Cremanium* is similar, but with more species beyond its center of richness in the Andes. *Chaenoplectrona* is quite heterogeneous (Judd 2007), with a good portion of the species in the Greater Antilles (mostly Hispaniola and Cuba), and another portion in the Andes. The other sections are not clearly concentrated in any particular geographical region. Section *Miconia* is the largest one in almost all countries/regions, except for Southeastern and Southern Brazil, Greater Antilles and Ecuador. It has

more than 100 species in Northern Brazil alone (2/3 of the species there), and that might be explained either by the fact that this region is surrounded by the Andes, Guayana Highlands, and Central Brazil (deriving its species from different floras; Oliveira & Daly 1999) or because species of section *Miconia* apparently prefer lowland forests. Sections *Jucunda* and *Tamonea* are doubtfully distinct (Goldenberg 2000), and together are richer in Colombia, but equally diverse in Central America, Venezuela, Ecuador, Peru and Northern Brazil. Among the sections, endemism rates are apparently higher in sections *Chaenopleura*, *Amblyarrhena*, *Octomeris* and *Cremanium*, and much lower in *Miconia*, *Jucunda* and *Tamonea*, (Goldenberg 2000).

TABLE 1. Number of species of *Miconia* by country or region and by section. Ade: section *Adenodesma*; Amb: *Amblyarrhena*; Chn: *Chaenanthera*; Chp: *Chaenopleura*; Cre: *Cremanium*; Glo: *Glossocentrum*; Hrt: *Hartigia*; Juc: *Jucunda*; Lac: *Laceraria*; Mic: *Miconia*; Oct: *Octomeris*; Tam: *Tamonea*; Indet: section uncertain

| Countries/Regions | Total | ade | amb | cht | chp | cre | glo | hrt | juc | lac | mic | oct | tam | indet |
|---------------------------------|-------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-------|
| Mexico | 79 | 1 | 12 | 1 | - | 5 | 2 | 4 | 4 | 1 | 31 | 10 | 8 | - |
| Central America | 157 | 1 | 17 | 2 | 6 | 31 | 2 | 5 | 7 | 4 | 55 | 11 | 16 | - |
| Greater Antilles | 112 | 1 | 4 | 2 | 48 | 9 | 6 | 2 | 1 | - | 20 | 8 | 7 | 4 |
| Lesser Antilles | 21 | - | - | - | - | 5 | 3 | 1 | - | 2 | 5 | 2 | 3 | - |
| Colombia | 338 | 3 | 78 | 3 | 11 | 69 | 8 | 6 | 7 | 4 | 97 | 13 | 39 | - |
| Venezuela | 212 | 5 | 22 | 3 | 3 | 29 | 7 | 4 | 8 | 4 | 93 | 14 | 20 | - |
| Trinidad | 42 | 2 | - | 3 | - | - | 3 | 3 | 2 | 1 | 22 | - | 6 | - |
| Guianas | 104 | 4 | 3 | 4 | - | 3 | 5 | 1 | 10 | 7 | 58 | - | 9 | - |
| Ecuador | 254 | 2 | 73 | 3 | 14 | 52 | 7 | 2 | 2 | 7 | 62 | 7 | 23 | - |
| Peru | 302 | 4 | 61 | 3 | 18 | 61 | 11 | 2 | 7 | 9 | 98 | 4 | 24 | - |
| Bolivia | 141 | 1 | 15 | 2 | 3 | 27 | 9 | 1 | 7 | 5 | 59 | 2 | 10 | - |
| Brazil North | 162 | 5 | - | 4 | - | 1 | 11 | 2 | 14 | 6 | 102 | 1 | 16 | - |
| Brazil Northeast | 71 | 1 | - | 4 | - | 3 | 15 | 2 | 4 | 1 | 32 | 2 | 5 | 2 |
| Brazil Central West | 68 | 1 | - | 4 | - | 3 | 11 | 1 | 7 | - | 37 | - | 4 | - |
| Brazil Southeast | 129 | - | 3 | 11 | - | 8 | 56 | 1 | 5 | 1 | 34 | 4 | 5 | 1 |
| Brazil South | 40 | - | 1 | 3 | - | 4 | 22 | - | 2 | - | 7 | - | 1 | - |
| Argentina, Paraguay and Uruguay | 21 | - | - | 1 | - | 3 | 5 | - | 2 | - | 9 | - | 1 | - |
| Species totals | 1057 | 6 | 212 | 17 | 86 | 223 | 82 | 7 | 23 | 19 | 245 | 59 | 71 | 7 |

Recent molecular phylogenies of the Miconieae (Michelangeli et al. 2004, Goldenberg et al. 2008, Martin et al. 2008, Michelangeli et al. 2008) have shown that both the large genera and their sections seem not to be monophyletic. In the case of *Miconia*, we can identify at least three patterns that explain the non-monophyletic genus and sections (Goldenberg et al. 2008, Michelangeli et al. 2008; Michelangeli et al., in prep.). First, some groups traditionally placed in the genus *Miconia* seem to be basal within the tribe, like some species from section *Tamonea* (*M. mirabilis* and *M. dodecandra*), and section *Laceraria*. Second, there are several cases of species that have been assigned to a given section, but are actually more closely related to species from another section or even another genus (i.e. all the species of *Amblyarrhena* and *Octomeris* that are from Southeastern and Southern Brazil appear to be better placed elsewhere; Goldenberg et al. 2008). Finally, albeit the sections are not monophyletic, groups of sections seem to assemble monophyletic entities, at least on a large scale. Moreover, these large clades seem to be geographically coherent. Molecular phylogenies have shown, for instance, that the majority of the species from sections *Jucunda*, *Tamonea*, *Adenodesma*, *Miconia*, *Glossocentrum* and *Chaanthera* belong to clades that are strongly represented in extra-Andean South America. Sections *Amblyarrhena*, *Cremanium* and *Chaenopleura* are nested in a clade that is strongly centered

in the Andes, but also diverse in Central America and the Greater Antilles. Finally, species from *Octomeris* either belong to this last clade, or are nested among other genera in a large Caribbean clade.

The strong correlation between sectional placement of the different species of *Miconia* and biogeography, which can also be seen in phylogenetic analyses of the Miconieae (Goldenberg et al. 2008, Michelangeli et al., 2008) is not only seen on a large scale, but also in the type of environment and plant associations that they occupy. For example, not only are most species of *Miconia* section *Miconia* found in lowlands of South America (and most of these in the Amazonian basin), but within this area they are predominantly found in moist forests over acidic soils. In fact, even though different species of *Miconia* can be found throughout the Neotropics from sea level up to 4000 m, the vast majority of species of *Miconia* (and Miconieae) occupy humid and forested environments, being rare in seasonal forests and completely absent from dry environments in tropical America. This pattern of distribution of *Miconia* (and the Miconieae as a whole) provides a great system in which to test the effect of niche conservatism (see Harvey & Pagel 1991, Wiens 2004, Donoghue 2008, Crisp et al. 2009, Antonelli & Sanmartin 2011) and the colonization of new environments as these appear (i.e. the uplift of the Andes, changes in the hydrology of the Amazon basin, etc; Bell & Donoghue 2005, Hughes & Eastwood 2006, Antonelli et al. 2009) on the diversification of large neotropical clades. Similar patterns of geographic and environmental coherence have been recently shown for the neotropical members of the capsular-fruited tribe Melastomeae (Michelangeli et al. 2013).

As for nomenclatural and taxonomic work still to be done, there are many issues regarding types that are in need of attention. First there is a long list of species with types that have not been located, and others still needing lectotypifications. The latter were described based on syntypes or without a clear choice of a holotype among cited collections. Several holotypes were destroyed at Berlin during World War II. Among those described by Markgraf there are apparently no extant isotypes (or none has yet been located). Finally, most species described by Ruiz & Pavon, D. Don and J. Triana based on the former's Peruvian collections present typification problems in need of careful study. Because *Miconia* is a conserved generic name (Farr et al. 1979), it is somewhat disconcerting that the type specimen of *Miconia triplinervis* Ruiz & Pav., the generic type, has not been located. We are uncertain if a type even exists but the identity of this species is unquestionable and has never been a source of conflicting interpretations.

Acknowledgements

This work is dedicated to six botanists, ordered here chronologically: Alphonse L. de Candolle, Charles V. Naudin, José J. Triana, Alfred C. Cogniaux, Henry A. Gleason and John J. Wurdack who collectively named more than 2/3 of the species listed here. We particularly thank the late John J. Wurdack for advice and assistance, and for assembling an extraordinary collection at US, upon which much of this work has been based.

The authors thank Carmen Ulloa-Ulloa and Janelle Burke for reviewing the lists, CNPq (Proc. 302325/2009-3) and CAPES (BEX 9095/11-0) for funding R.G., NSF (DEB-0818399) for funding F.A.M., F.A., R.G., CNPq and CAPES (Brazil) for funding M.K.C. and J.M. Curators and staff at the US National Herbarium at Smithsonian Institution have also facilitated extensive use of this collection by most of the authors, and particularly Dr. Vicki Funk hosted RG for a sabbatical from July 2012 to July 2013. The Global Biodiversity Information Facility funded the Melastomataceae.net structure that houses the MelNames database, a project initiated and coordinated by Susanne Renner at Botanische Staatssammlung München.”

References

- Almeda, F. (2001) Melastomataceae. In: Stevens, W.D., Ulloa-Ulloa, C., Pool, A., & Montiel, O. M. (eds.), *Flora de Nicaragua*. Monographs in Systematic Botany from the Missouri Botanical Garden 85: 1339–1419.
Almeda, F. (2009) Melastomataceae. In: Davidse, G., Sousa-Sánchez, M., Knapp, S. & Chiang, F. (eds.) *Flora Mesoamericana* 4 (1). Universidad Nacional Autónoma de México, México, pp. 164–338.

- Almeda, F., Kriebel, R. & Umaña, G. (2007) Melastomataceae. In Hammel, B.E., Grayum, M.H., Herrera, C. & Zamora, N. (eds.) *Manual de Plantas de Costa Rica*. Monographs in Systematic Botany from the Missouri Botanical Garden 6: 394–574.
- Antonelli, A., Nylander, J.A.A., Persson, C. & Sanmartin, I. (2009) Tracing the impact of the Andean uplift on Neotropical plant evolution. *Proceedings of the National Academy of Sciences of the United States of America* 106: 9749–9754.
- Antonelli, A. & Sanmartín, I. (2011) Why are there so many plant species in the Neotropics? *Taxon* 60(2): 403–414.
- Aublet, J.B.C.F. (1775) *Histoire des plantes de la Guiane Françoise* 1. P.F. Didot jeune, London, Paris, 621 p.
- Baumgratz, J.F.A. (1980) Miconias do município do Rio de Janeiro. Seção *Miconia* DC. (Melastomataceae). *Rodriguésia* 32: 73–95.
- Baumgratz, J.F.A. (1982) Miconias do estado do Rio de Janeiro. Seção *Tamonea* (Aubl.) Cogn. (Melastomataceae). *Arquivos do Jardim Botânico do Rio de Janeiro* 26: 69–86.
- Baumgratz, J.F.A. (1984) Miconias do estado do Rio de Janeiro. Seção *Chaenanthera* Naud. (Melastomataceae). *Rodriguésia* 36: 45–58.
- Baumgratz J.F.A., D'El Rey Souza, M.L., Martins, A. B., Lughadha, E.N. & Woodgyer, E.M. (1995) Melastomataceae. In: Stannard, B.L. (ed.), *Flora of the Pico das Almas*. Royal Botanical Gardens, Kew, pp. 433–483.
- Bell, C.D. & Donoghue, M.J. (2005) Phylogeny and biogeography of Valerianaceae (Dipsacales) with special reference to the South American valerians. *Organisms Diversity & Evolution* 5: 147–159.
<http://dx.doi.org/10.1016/j.ode.2004.10.014>
- Berry, P.E. (2001) *Miconia*. In: J.A. Steyermark, P.E. Berry, K. Yatskivych & B.K. Holst (eds.) *Flora of the Venezuelan Guayana* 6. Missouri Botanical Garden, St. Louis, pp. 433–483.
- Bremer, K. (1987) Melastomataceae. In: Dassanayake, M.D. & Fosberg, F.R. (eds.), *A Revised Handbook to the Flora of Ceylon Vol. VI*. Amerind Publishing Company, New Delhi, pp. 157–240.
- Briquet, J. (1906) *International rules of botanical nomenclature adopted by the International Botanical Congress of Vienna, 1905*. Fisher, Jena, 99 p.
- Buchoz, P.J. (1779) *Plantes nouvellement découvertes recemment dénommées et classées*. Edited by the author, Paris, 160p.
- Camp, W.H., Rickett, H.W. & Weatherby, C.A. (1947) Appendix III: Nomina Generica Conservanda. *Brittonia* 6: 47–93.
- Clausing G. & Renner, S.S. (2001) Molecular Phylogenetics of Melastomataceae and Memecylaceae: implications for character evolution. *American Journal of Botany* 88: 486–498.
<http://dx.doi.org/10.2307/2657114>
- Cogniaux, C.A. (1891) Mélastomacées. In: De Candolle, A. & De Candolle, C. (eds.), *Monographiae Phanerogamarum* 7. G. Masson, Paris, pp. 1–1256.
- Cogniaux, C.A. (1913) Die von Dr. Th. Herzog auf seiner zweiten Reise durch Bolivien in den Jahren 1910 und 1911 gesammelten Pflanzen — Melastomataceae. *Mededeelingen van 's Rijks Herbarium Leiden* 19: 76—81.
- Cotton, E. & Michelangeli, F.A. (2007) Melastomataceae. In: Duno de Stefano, R., Aymard, G. & Huber, O. (eds.), *Catálogo anotado e ilustrado de la flora vascular de los llanos de Venezuela*, FUDENA, Fundación Empresas Polar & FIBV, Caracas, pp. 559–571.
- Crisp, M.D., Arroyo, M.T.K., Cook, L.G., Gandolfo, M.A., Jordan, G.J., McGlone, M.S., Weston, P.H., Westoby, M., Wilf, P. & Linder, H.P. (2009) Phylogenetic biome conservatism on a global scale. *Nature* 458: 754–790.
- Csurhes, S. (2008) Pest Risk Assessment - *Miconia calvescens*. Queensland Government, Department of Primary Industries and Fisheries, Brisbane, pp. 1–14
- Donoghue, M.J. (2008) A phylogenetic perspective on the distribution of plant diversity. *Proceedings of the National Academy of Sciences of the United States of America* 105: 11549–11555.
- Duss, R.P. (1896) Plantes de la Guadeloupe et de la Martinique. *Annales de L'Institut Colonial de Marseille* 3: 1–656.
- Fawcett, W. & Rendle, A.B. (1926) Melastomataceae. In: Fawcett, W. & Rendle, A.B. (eds.), *Flora of Jamaica* 5. The British Museum, London, pp. 353–405.
- Farr, E.R., Leussink, J.A. & Stafleu, F.A. (1979) Index Nominum Genericorum (Plantarum) Vol. II – *Eprolithus* – *Pelesia*. *Regnum Vegetabile* 101: 631–1276.
- Funk, V., Hollowell, T., Berry, P., Kelloff, C. & Alexander, S.N. (2007) Checklist of the plants of the Guiana Shield (Venezuela: Amazonas, Bolívar, Delta Amacuro; Guyana, Surinam, French Guiana). *Contributions from the United States National Herbarium* 55: 1–584.
- Gleason, H.A. (1935) Melastomataceae. In: Pulle, A. (ed.), *Flora of Suriname*. Koninklijk Koloniaal Instituut, Amsterdam, pp. 178–281.
- Gleason, H.A. (1940) Botany of the Maya area: miscellaneous papers. XX. The Melastomataceae of the Yucatan peninsula. *Publications of the Carnegie Institution of Washington* 522: 323–373.
- Gleason, H.A. (1958) Flora of Panama, part VII. Fascicle 3. Melastomataceae. *Annals of the Missouri Botanical Garden* 45: 203–304.
- Goldenberg, R. (2000) O gênero *Miconia* Ruiz & Pav. (Melastomataceae). I. Listagens analíticas, II. Revisão taxonômica

- da seção *Hypoxanthus* (Rich. ex DC.) Hook. F. Ph.D diss. Universidade Estadual de Campinas, Campinas, Brazil.
- Goldenberg, R. (2004) O gênero *Miconia* (Melastomataceae) no estado do Paraná, Brasil. *Acta Botanica Brasilica* 18: 927–947.
<http://dx.doi.org/10.1590/S0102-33062004000400024>
- Goldenberg, R. (2009) *Miconia Ruiz & Pav.* In: Wanderley, M.G.L., Shepherd, G.J., Melhem, T.S., Giulietti, A.M. & Martins, S.E. (eds.), *Flora fanerogâmica do estado de São Paulo 6*. Fapesp, São Paulo, pp. 73–103.
- Goldenberg, R., Caddah, M.K. & Martin, C.V. (2010) Taxonomic notes on South American *Miconia* (Melastomataceae). *II. Rodriguésia* 61: S23–S28.
- Goldenberg, R. & Martin, C.V. (2008) Taxonomic notes on South American *Miconia* (Melastomataceae). *Harvard Papers in Botany* 13: 223–227.
<http://dx.doi.org/10.3100/1043-4534-13.2.223>
- Goldenberg, R., Penneys, D., Almeda, F., Judd, W.S. & Michelangeli, F.A. (2008) Phylogeny of *Miconia* (Melastomataceae): patterns of stamen diversification in a megadiverse neotropical genus. *International Journal of Plant Sciences* 169: 963–979.
- Harvey, P.H. & Pagel, M.D. (1991). *The comparative method in evolutionary biology*. Oxford University Press, Oxford, 239p.
- Howard, R.A. (1989) Melastomataceae. In: *Flora of the Lesser Antilles, Dicotyledonae-Part 2*. Arnold Arboretum, Jamaica Plain, pp. 532–579.
- Howard, R.A. & Kellogg, E.A. (1986) Nomenclatural notes on *Miconia* (Melastomataceae). *Journal of the Arnold Arboretum* 67: 233–255.
- Hughes, C. & Eastwood, R. (2006) Island radiation on a continental scale: Exceptional rates of plant diversification after uplift of the Andes. *Proceedings of The National Academy of Sciences of The United States of America* 103: 10334–10339.
- Ionta, G.M., Judd, W.S., Skean, J.D. & McMullen, C.K. (2012) Two new species of *Miconia* sect. *Sagraea* (Melastomataceae) from the Macaya Biosphere Reserve, Haiti, and twelve relevant new species combinations. *Brittonia* 64: 61–72.
- Johnston, I.M. (1949) The botany of San Jose Island. *Sargentia* 8: 1–306.
- Judd, W.S. (2007). Revision of *Miconia* sect. *Chaenopleura* (Miconieae, Melastomataceae) in the Greater Antilles. *Systematic Botany Monographs* 81: 1–235.
- Judd, W. S. & Skean, J.D. (1991) Taxonomic studies in the Miconieae (Melastomataceae). IV. Generic realignments among terminal-flowered taxa. *Bulletin of the Florida State Museum, Biological Science* 36: 25–84.
- Liogier, A.H. [Alain, Brother] (1957) Flora de Cuba. Part. 3 – Melastomataceae. *Contribuciones Ocasionales del Museo de Historia Natural del Colegio "De La Salle"* 16: 9–62.
- Liogier, A.H. (2000) Melastomataceae. In: *La Flora de La Española 9*. Editora Corripio, Santo Domingo, pp. 1–81.
- Liogier, A.H. [Alain, Brother] & Martorell, L.F. (1982) *Flora of Puerto Rico and adjacent islands: a systematic synopsis*. Editorial de la Universidad de Puerto Rico, Rio Piedras, 342 pp.
- Macbride, J.F. (1941) Flora of Peru — Melastomataceae. *Publications of the Field Museum of Natural History, Botanical Series* 13: 249–523.
- Martin, C.V. & Cremers, G. (2007) Les Melastomataceae américaines décrites par C. Naudin. *Le Journal de Botanique* 37: 3–111.
- Martin, C.V., Little, D.P., Goldenberg, R., & Michelangeli, F.A. (2008) A phylogenetic evaluation of *Leandra* (Miconieae, Melastomataceae): a polyphyletic genus where the seeds tell the story, not the petals, *Cladistics* 24: 315–327.
- Martins A.B., Semir, J., Goldenberg, R. & Martins, E. (1996) O gênero *Miconia Ruiz & Pav.* (Melastomataceae) no estado de São Paulo. *Acta Botanica Brasilica* 10: 267–316.
- Meyer, J.Y. (1996) Status of *Miconia calvescens* (Melastomataceae), a dominant tree in the Society Islands (French Polynesia). *Pacific Science* 50: 66–76.
- Michelangeli, F.A. & Cotton, E. (2008) Melastomataceae. In: Hokche, R., Berry, P.E. & Huber, O. (eds.) *Nuevo Catálogo de la Flora Vascular de Venezuela*. Fundación Instituto Botánico de Venezuela, Caracas, pp. 466–484.
- Michelangeli, F.A. & Goldenberg, R. (2012) Replacement names and synonyms in *Miconia* (Melastomataceae). *Phytotaxa* 52: 29–32.
- Michelangeli, F.A., Penneys, D.S., Giza, J., Soltis, D., Hils, M.H. & Skean, J.D. (2004) A preliminary phylogeny of the tribe Miconieae (Melastomataceae) based on nrITS sequence data and its implications on inflorescence position. *Taxon* 53: 279–290.
- Michelangeli, F.A., Judd, W.S., Penneys, D.S., Skean, J.D., Bécquer-Granados, E.R., Goldenberg, R., Martin, C.V. (2008) Multiple events of dispersal and radiation of the tribe Miconieae (Melastomataceae) in the Caribbean. *Botanical Review* 74: 53–77.
<http://dx.doi.org/10.1007/s12229-008-9004-x>
- Michelangeli, F.A., Guimaraes, P.J.F., Penneys, D.S., Almeda, F. & Kriebel, R. (2013) Phylogenetic relationships and distribution of New World Melastomeae (Melastomataceae). *Botanical Journal of the Linnean Society* 171: 38–60

- Molina, A. (1975) Enumeracion de las plantas de Honduras. *Ceiba* 19: 1–118.
- Murray, J.A (1774) *Systema Vegetabilium: secundum classes, ordines, genera, species cum characteribus et differentiis 13*. Jo. Christ. Dietrich, Gottingen, 844p.
- Neill, D.A. & Ulloa-Ulloa, C. (2011) *Adiciones a la Flora del Ecuador: Segundo suplemento, 2005–2010*. Fundación Jatun Sacha, Quito, 202 pp.
- Necker, N.J. (1790) *Elementa Botanica*. Bosange, Paris, 460p.
- Nicolson, D.H. (1991) Flora of Dominica, Part 2: Dicotyledonae. *Smithsonian Contributions to Botany* 77: 1–274.
<http://dx.doi.org/10.5479/si.0081024X.77>
- Oliveira, A.A. & Daly, D. 1999. Geographic distribution of tree species occurring in the region of Manaus, Brazil: Implications for regional diversity and conservation. *Biodiversity and Conservation* 8(9): 1245–1259.
- Proctor, G.R. (1972) Melastomataceae. In: Adams, C. D. (ed.), *Flowering Plants of Jamaica*. University of the West Indies, Mona, pp. 529–549.
- Renner, S.S. (1999) Catalogue of the vascular plants of Ecuador (by Jørgensen, P.M. & León-Yáñez, S.) – Melastomataceae. *Monographs in Systematic Botany from the Missouri Botanical Garden* 75: 561–585.
- Rickett, H.W. & Stafleu, F.A. (1960). Nomina generica conservanda et rejicienda spermatophytorum IV (Continued). *Taxon* 9: 67–86.
<http://dx.doi.org/10.2307/1216832>
- Riley, L.A.M. (1925) Critical notes on Trinidad plants. *Kew Bulletin* 3: 133–141.
- Ruiz, H. & Pavón, J. (1794) *Florae peruvianaæ, et chilensis prodromus*. Imprenta de Sancha, Madrid, 153 p.
- Scopoli, G.A. (1777) *Introductio ad Historiam Naturalem*. 540p. Wolfgang Gerle, Prague, 540p.
- Seymour, F.C. (1980) A Check List of the Vascular Plants of Nicaragua based largely on collections in Nicaragua made by the author and companions 1968–1976. *Phytologia Memoirs* 1: 1–314.
- Standley, P.C. (1931) Flora of the Lancetilla Valley, Honduras. *Publications of the Field Museum of Natural History, Botanical Series* 10: 305–311.
- Standley, P.C. & Record, S.J. (1936) Forests and flora of British Honduras – *Miconia*. *Publications of the Field Museum of Natural History, Botanical Series* 12: 293–298.
- Standley, P.C. & Williams, L.O. (1963) Flora of Guatemala: Melastomaceae. *Fieldiana, Botany* 24: 407–525.
- Uribe-Uribe, L. (1972) Melastomataceae. In: Pinto-Escobar, P. (ed.), *Catalogo ilustrado de las plantas de Cundinamarca*. Universidad nacional, Bogotá, pp. 61–150.
- Ulloa-Ulloa, C., Zarucchi, J.L. & León, B. (2004) Diez años de adiciones a la flora del Perú: 1993–2003. *Arnaldoa Edición Especial*: 1–242.
- Ulloa-Ulloa, C. & Neill, D.A. (2005) *Cinco años de adiciones a la flora del Ecuador: 1999–2004*. UTP Loja, MBG Press & Funbotanica, Loja, 75 pp.
- Wiens, J.J. (2004) Speciation and ecology revisited: Phylogenetic niche conservatism and the origin of species. *Evolution* 58: 193–197.
<http://dx.doi.org/10.1111/j.0014-3820.2004.tb01586.x>
- Williams, L.O. (1963) Tropical American Plants, V. *Fieldiana, Botany* 29: 545–597.
- Williams, R.O. (1934) *Flora of Trinidad and Tobago*. Department of Agriculture, Port-of-Spain v.1, pp. 333–410.
- Winkler, S. (1965) Die Melastomataceae von El Salvador. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 83: 331–369.
- Wurdack, J.J. (1962) Melastomataceae of Santa Catarina. *Sellowia* 14: 109–217.
- Wurdack, J.J. (1963) Plantae Austro-Americanæ. XI (Colombia) – Melastomataceæ. *Rhodora* 65: 18–20.
- Wurdack, J.J. (1970) Erroneous data in Glaziou collections of Melastomataceæ. *Taxon* 19: 911–913.
<http://dx.doi.org/10.2307/1218309>
- Wurdack, J.J. (1971a) Notes on some cultivated species of *Miconia* (Melastomataceæ). *Baileya* 18: 17–19.
- Wurdack, J.J. (1971b) Geographic errors in Bonpland collections of Melastomataceæ. *Taxon* 20: 591–593.
<http://dx.doi.org/10.2307/1218260>
- Wurdack, J.J. (1971c) The Melastomataceæ collections of A.R. Ferreira. *Taxon* 20: 595–596.
<http://dx.doi.org/10.2307/1218261>
- Wurdack, J.J. (1973) Melastomataceæ. In: T. Lasser (ed.), *Flora de Venezuela* 8. Instituto Botanico, Caracas, pp. 1–819.
- Wurdack, J.J. (1976) Certamen Melastomataceis. XXV. *Phytologia* 35: 1–13.
- Wurdack, J.J. (1980) 138. Melastomataceæ. In: Harling, G. & Sparre, B. (eds.), *Flora of Ecuador* 13. University Göteborg and Riksmuseum, Stockholm, pp. 1–406.
- Wurdack, J.J. (1993) Melastomataceæ. In: Brako, L. & Zarucchi, J.L. (eds.), *Catalogue of the flowering plants and gymnosperms of Peru*. Missouri Botanical Garden, Saint Louis, pp. 672–707.
- Wurdack J.J., Renner, S.S. & Morley, T. (1993) Melastomataceæ. In: Görtz van Rijn, A.R.A. (ed.), *Flora of the Guianas* 13. Koeltz Scientific Books, Koenigstein, pp. 1–425.

LIST 1. “Nomenclator botanicus” for all published names of *Miconia*, including accepted species, synonyms, sectional placement, and distribution by country.

Miconia abbreviata Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 9: 1145 (1927). Type:—PERU. *Tessmann* 4548 (Holotype: B, destroyed; isotype: F).

Section: Miconia.

Distribution: Colombia, Venezuela, Guyana, Ecuador, Peru, Bolivia, Brazil North.

Miconia abeggii Urb. & Ekman, Ark. Bot. 22A (17): 35 (1929). Type:—HAITI. *Ekman H2790* (Holotype: S).

Section: Miconia.

Distribution: Haiti.

Miconia abyasmophila Wurdack, Mem. New York Bot. Gard. 10 (4): 39 (1961). Type:—VENEZUELA. *Maguire* 42206 (Holotype: NY; isotypes: F, K, S, US).

Section: Miconia.

Distribution: Venezuela.

Miconia acalephoides Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 207 (1850). *Acinodendron acalephoides* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Goudot s.n.* (Holotype: P; isotype: K).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia acanthocoryne Wurdack, Phytologia 5: 127 (1955). Type:—COLOMBIA. *von Sneidern* 5364 (Holotype: F; isotypes: F, MO).

Section: Cremanium.

Distribution: Colombia.

***Miconia acinodendron* (L.) Sweet**, Hort. Brit. (Sweet) 1: 159 (1826).

Basionym: *Melastoma acinodendrum* L., Sp. Pl. 1: 389 (1753). Type:—"AMERICA". *Hort. Cliff.* 162 (Lectotype: BM).

Melastoma plantaginifolium Mill., Gard. Dict. ed. 8 (1768). *Acinodendron plantaginifolium* (Mil.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Melastoma fragile L.f., Suppl. Pl. 236 (1781). *Acinodendron fragile* (L.f.) Kuntze, Revis. Gen. Pl. 2: 949 (1891).

Melastoma pleurocarpum E.Mey., Nova Acta Phys.-Med. 12: 795 (1825). *Miconia pleurocarpa* (E.Mey.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).

Melastoma serrulatum Spreng., Syst. Veg. 2: 300 (1825).

Clidemia berbiceana DC., Prodr. 3: 160 (1828). *Miconia berbiceana* (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 174 (1850). *Tschudya berbiceana* (DC.) Griseb., Abh. Königl. Ges. Wiss. Göttingen 9: 47 (1860).

Miconia eriopoda Benth., J. Bot. 2: 311 (1840).

Decaraphe fockeana Miq., Comm. Phytogr. 2: 84 (1840).

Decaraphe floribunda Miq., Linnaea 18: 280 (1844). *Miconia miquelianiana* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).

Miconia clidemiooides Steud., Flora 27: 724 (1844).

Section: Miconia.

Distribution: Martinique, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Peru, Brazil North.

***Miconia acuminata* (Steud.) Naudin**, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850).

Basionym: *Conostegia acuminata* Steud., Flora 27: 722 (1844). Type:—SURINAME. *Hostmann* 1266 (Holotype: P; isotype: MO).

Miconia axilliflora Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 179 (1850). *Acinodendron axilliflorum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Section: Jucunda.

Distribution: Suriname, French Guiana, Peru, Bolivia, Brazil North, Brazil Central-West.

Miconia acuminifera Triana, Trans. Linn. Soc. London 28: 105 (1871).

Acinodendron acuminiferum (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Triana* 3997
(Holotype: BM; isotypes: COL, K).

Section: Tamonea.

Distribution: Colombia.

Miconia acunae Borhidi, Abstracta Botanica (Budapest) 5: 23 (1977). Type:—CUBA. *Stuchlick* 27223 (Holotype: HAC; isotypes: BP, HAC).

Section: Glossocentrum.

Distribution: Cuba.

Miconia acutifolia Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 356 (1915). Type:—GUYANA. *Ule* 8694 (Holotype: B, destroyed; isotypes: G, K).

Section: Tamonea.

Distribution: Venezuela, Guyana, Brazil North.

Miconia acutipetala Sprague, Trans. & Proc. Bot. Soc. Edinb. 22: 432 (1905). Type:—COLOMBIA. *Sprague* s.n. (Holotype: K; isotypes: F, K, NY, US).

Section: Miconia.

Distribution: Colombia.

Miconia adenocalyx Urb. & Ekman, Ark. Bot. 21A (5): 41 (1927). Type:—HAITI. *Ekman* H2836 (Holotype: S; isotypes: NY, S, US).

Section: Chaenopleura.

Distribution: Haiti, Dominican Republic.

Miconia adinantha Wurdack, Phytologia 23: 487 (1972). Type:—PERU. *Dudley* 10384 (Holotype: US; isotypes: MO, NA).

Section: Miconia.

Distribution: Peru.

Miconia adrieni J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 180 (1929). Type:—PERU. *Macbride* 5077 (Holotype: F).

Section: Amblyarrhena.

Distribution: Peru.

Miconia aenigmatica Wurdack, Phytologia 5: 128 (1955). Type:—COLOMBIA. *Cuatrecasas* 21752 (Holotype: F; isotypes: F, NY, U, US, S).

Section: Cremanium.

Distribution: Colombia.

Miconia aequatorialis Wurdack, Phytologia 38: 297 (1978). Type:—ECUADOR. *Sparre* 18736 (Holotype: S; isotypes: NY, TEX, US).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia aeruginosa Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 135 (1850). *Acinodendron aeruginosum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Bonpland* s.n. (Holotype: P; isotype: P).

Miconia lindenii Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 135 (1850). *Acinodendron lindenii* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Section: Miconia.

Distribution: Mexico, Guatemala, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Guyana, Ecuador, Bolivia.

Miconia affinis DC., Prodr. 3: 187 (1828). *Acinodendron affine* (DC.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—FRENCH GUIANA. *Poiteau s.n.* (Holotype: G-DEL).

Miconia microcarpa DC., Prodr. 3: 189 (1828). *Acinodendron microcarpum* (DC.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Miconia planinervia Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 160 (1850). *Acinodendron planinerve* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Miconia cecidophora Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 166 (1850). *Acinodendron cecidophorum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Miconia pusilliflora Beurl., Kongl. Vetensk. Acad. Handl. 40: 130 (1854), *nom illeg.* *Miconia beurlingii* Triana, Trans. Linn. Soc. London 28: 107 (1871). *Acinodendron beurlingii* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Miconia cayumbensis Gleason, Bull. Torrey Bot. Club 66: 416 (1939).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Dominican Republic, Puerto Rico, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia aggregata Gleason, Bull. Torrey Bot. Club 52: 448 (1925). Type:—COLOMBIA. *Rusby 813* (Holotype: NY; isotype: US).

Miconia aggregata ssp. *australis* Wurdack, Phytologia 38: 295 (1978).

Section: Amblyarrhena.

Distribution: Colombia, Ecuador, Peru.

Miconia aguirrei L.Uribe, Caldasia 12 (56): 14 (1977). Type:—COLOMBIA. *Aguirre 428* (Holotype: COL; isotypes: COL, FMB, HUA, NY, US).

Section: Cremanium.

Distribution: Colombia.

Miconia aguitensis Gleason, Bull. Torrey Bot. Club 58: 426 (1931). Type:—VENEZUELA. *Tate 933* (Holotype: NY; isotype: US).

Section: Tamonea.

Distribution: Venezuela.

Miconia alainii Judd & Skean, Novon 4: 112 (1994). Type:—DOMINICAN REPUBLIC. *Liogier 25179* (Holotype: JBSD).

Section: Chaenopleura.

Distribution: Dominican Republic.

Miconia alata (Aubl.) DC., Prodr. 3: 184 (1828).

Basionym: *Melastoma alatum* Aubl., Hist. pl. Guiane 1: 410 (1775). *Clidemia alata* (Aublet) G.Don ex Loud., Hort. Brit. 1: 174 (1830). *Acinodendron alatum* (Aubl.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—FRENCH GUIANA. *Aublet s.n.* (Holotype: BM).

Miconia alata var. *amazonica* Schrank ex DC., Prodr. 3: 184 (1828).

Miconia pterophora Miq., Linnaea 18: 622 (1844).

Section: Miconia.

Distribution: Colombia, Venezuela, Guyana, Suriname, French Guiana, Brazil North, Brazil Central-West, Brazil Northeast.

Miconia alberti Gleason, Bull. Torrey Bot. Club 57: 70 (1930). Type:—COLOMBIA. *Killip 15896* (Holotype: NY; isotype: A).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia albicans (Sw.) Steud., Nomencl. Bot. 2: 139 (1841).

Basionym: *Melastoma albicans* Sw., Prodr. 70 (1788). *Miconia albicans* (Sw.) Triana, Trans. Linn. Soc. London 28: 116 (1871), nom. illeg.. *Acinodendron albicans* (Sw.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—JAMAICA. *Swartz s.n.* (Lectotype: S; isolectotype: BM).

Melastoma holosericeum Vahl, Eclog. Amer. 1: 42 (1797).

Melastoma velutinum Willd., Sp. Pl. 2: 584 (1799).

Miconia detergilobis DC., Prodr. 3: 181 (1828). *Melastoma detergibile* Schrank & Mart. ex DC., Prodr. 3: 181 (1828), pro syn.

Miconia holosericea var. *acuminata* Ser. ex DC., Prodr. 3: 182 (1828).

Miconia holosericea var. *oblongata* Ser. ex DC., Prodr. 3: 182 (1828).

Miconia holosericea var. *obtusiuscula* DC., Prodr. 3: 182 (1828).

Miconia heterochroa Miq., Linnaea 17: 621 (1844).

Miconia renggeri Steud., Flora 27: 723 (1844).

Miconia holosericea var. *montana* Crueg., Linnaea 20: 110 (1847).

Miconia montana Crueg. ex Triana, Trans. Linn. Soc. London 28: 116 (1871), pro syn.

Miconia rufescens Macfad. ex Griseb., Fl. Brit. W. Indian Is. 256 (1860).

Melastoma nitidum Pav. ex Triana, Trans. Linn. Soc. London 28: 116 (1871), pro syn.

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Cuba, Jamaica, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast, Brazil South, Paraguay.

Miconia albiviridis Urb. & Ekman, Ark. Bot. 20A (5): 26 (1926). Type:—HAITI. *Ekman H1400* (Holotype: S; isotypes: EHH, NY, US).

Section uncertain

Distribution: Haiti.

Miconia alborosea L.Uribe, Caldasia 10: 289 (1969). Type:—COLOMBIA. *Uribe-Uribe 6082* (Holotype: COL; isotypes: NY, US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia alborufescens Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 160 (1850). *Acinodendron alborufescens* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Claussen 154* (Holotype: P; isotypes: BR, NY).

Miconia arirambae Huber, Bull. Soc. Bot. Genève 6: 192 (1914).

Miconia cachimbensis Brade, Arch. Jard. Bot. Rio de Janeiro 16: 14 (1959).

Section: Glossocentrum.

Distribution: Colombia, Venezuela, Guyana, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia aligera Wurdack, Phytologia 23: 486 (1972). Type:—PERU. *Dudley 11261* (Holotype: US).

Section: Amblyarrhena.

Distribution: Peru.

Miconia aliquantula Wurdack, Brittonia 39: 161 (1987). *Tschudya pulverulenta* DC., Prodr. 3: 155 (1828). *Staphidiastrum pulverulentum* (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 334 (1851). *Oxymeris pulverulenta* (DC.) Triana, Trans. Linn. Soc. London 28: 96 (1871). *Leandra pulverulenta* (DC.) Cogn. in Martius, Fl. bras. 14 (4): 208 (1886). Type:—FRENCH GUIANA. *Richard s.n.* (Holotype: P).

Clidemia pusilliflora Hoehne, Anexos Mem. Inst. Butantan, Secç. Bot. 1 (5): 167 (1922).

Section: Laceraria.

Distribution: Venezuela, Suriname, French Guiana, Brazil North.

Miconia alloeotricha (Urb.) Judd, Penneys & Skean, Brittonia 56: 160 (2004).

Basionym: *Ossaea alloeotricha* Urb., Ark. Bot. 22A (17): 64 (1929). *Leandra alloeotricha* (Urb.) Judd & Skean, Bull. Florida State Mus., Biol. Sci. 36: 61 (1991). Type:—HAITI. *Ekman H7473* (Holotype: S; isotype: NY).

Section uncertain

Distribution: Haiti.

Miconia alpestris Cogn. in Smith, Bot. Gaz. 20: 288 (1895). Type:—GUATEMALA. *Heyde 3334* (Holotype: BR; isotypes: K, NY, US).

Section: Cremanium.

Distribution: Guatemala.

Miconia alpina Cogn., Bot. Jahrb. Syst. 42: 146 (1909). Type:—PERU. *Weberbauer 4953* (Syntype: B, destroyed; isosyntypes: BR, F, MOL).

Section: Chaenopleura.

Distribution: Peru.

Miconia alternans Naudin, Melast. Monogr. Descr. 720 (1853). *Acinodendron alternans* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—FRENCH GUIANA. *Martin s.n.* (Lectotype: P; isolectotypes: BR, P).

Section: Miconia.

Distribution: Colombia, Venezuela, Guyana, Suriname, French Guiana, Peru, Bolivia, Brazil North.

Miconia alternifolia (Griseb.) Alain, Contr. Ocas. Mus. Hist. Nat. Colegio "De La Salle" 14: 2 (1955).

Basionym: *Catachaenia alternifolia* Griseb., Abh. Konigl. Ges. Wiss. Gottingen 9: 55 (1860). *Acinodendron alternifolium* (Griseb.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—CUBA. *Wright 179 p.p.* (Holotype: GOET; isotypes: BM, GH, NY, S).

Catachaenia parviflora Griseb., Cat. Pl. Cub. 102 (1866). *Miconia grisebachii* Triana, Trans. Linn. Soc. London 28: 130 (1871).

Section: Chaenopleura.

Distribution: Cuba.

Miconia altissima Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 837 (1891). *Acinodendron altissimum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Glaziou 16909* (Holotype: BR; isotypes: C, F, G, K, NY, P, R, US).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia alypifolia Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 216 (1850). *Acinodendron alypifolium* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. *Bonpland s.n.* (Holotype: P; isotype: P).

Section: Amblyarrhena.

Distribution: Colombia, Ecuador, Peru.

Miconia amabilis Cogn. in Rusby, Bull. New York Bot. Gard. 4: 359 (1907). Type:—BOLIVIA. *Rusby 2649* (Holotype: NY; isotypes: BR, CORD).

Section: Amblyarrhena.

Distribution: Bolivia.

Miconia amacurensis Wurdack, Acta Bot. Venez. 2: 373 (1967). Type:—VENEZUELA. *Steyermark 87115* (Holotype: US; isotypes: CAS, NY, VEN).

Section: Jucunda.

Distribution: Venezuela, Guyana, French Guiana, Brazil North, Brazil Northeast.

Miconia amambayensis Kraenzl., Vierteljahrsschr. Naturf. Ges. Zürich 76: 156 (1931). Type:—PARAGUAY. *Hassler 11381* (Holotype: Z; isotypes: A, BM, G, K, LIL, MO, MPU, NY, P, US).

Section: Tamonea.

Distribution: Paraguay.

Miconia amazonica Triana, Trans. Linn. Soc. London 28: 103 (1871). *Acinodendron amazonicum* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. Poeppig 3086 (Holotype: B, destroyed; isotype: BR).
Section: Tamonea.
Distribution: Colombia, Ecuador, Peru, Brazil North.

Miconia amblyandra Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 206 (1850). *Acinodendron amblyandrum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. Funck 1408 (Holotype: P; isotypes: BR, G, MPU, S).
Section: Amblyarrhena.
Distribution: Colombia, Venezuela.

Miconia amilcariana Almeda & Dorr, Proc. Calif. Acad. Sci., Ser. 4 57: 549 (2006). Type:—VENEZUELA. *Stergios* 17582 (Holotype: PORT; isotypes: CAS, US).
Section: Cremanium.
Distribution: Venezuela.

Miconia amissa Wurdack, Phytologia 16: 177 (1968). *Graffenrieda stellipilis* Gleason, Amer. J. Bot. 19: 742 (1932).
Type:—BOLIVIA. Buchtien 7407 (Holotype: US).
Section: Miconia.
Distribution: Bolivia.

Miconia amnicola Wurdack, Phytologia 48: 243 (1981). Type:—PERU. Schunke 158 (Holotype: US).
Section: Miconia.
Distribution: Colombia, Ecuador, Peru, Brazil North.

Miconia amoena Triana, Trans. Linn. Soc. London 28: 115 (1871). *Acinodendron amoenum* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. Blanchet 1707 (Holotype: B, destroyed; isotype: BR).
Section: Miconia.
Distribution: Brazil Northeast, Brazil Southeast.

Miconia ampla Triana, Trans. Linn. Soc. London 28: 101 (1871). *Acinodendron amplum* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—TRINIDAD. Lockhardt s.n. (Holotype: K).

Miconia involucrata Donn.Sm., Bot. Gaz. 37: 209 (1904).

Miconia megaphylla Gleason, Bull. Torrey Bot. Club 59: 363 (1932).

Section: Jucunda.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Jamaica, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West.

Miconia amplexicaulis Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 136 (1850). *Acinodendron amplexicaule* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. Poeppig 1424 (Holotype: P; isotypes: BM, BR, P).

Miconia amplexicaulis var. *parvifolia* Cogn. in Martius, Fl. bras. 14 (4): 273: 14(4): 273 (1887).

Section: Miconia.

Distribution: Peru.

Miconia amplinodis Umaña & Almeda, Novon 5: 110 (1995). Type:—COSTA RICA. Umaña 495 (Holotype: CR; isotypes: CAS, COL, F, MEXU, MO, US, USJ).

Section: Cremanium.

Distribution: Costa Rica.

Miconia ancistrophora (C.Wright) Triana, Trans. Linn. Soc. London 28: 103 (1871).

Basionym: *Tetrazygia ancistrophora* C.Wright in Sauvalle, Anales Acad. Ci. Med. Habana 5: 465 (1869). *Acinodendron ancistrophorum* (C.Wright) Kuntze, Revis. Gen. Pl. 2: 950 (1891). *Tetrazygiopsis ancistrophora* (C.Wright) Borhidi, Acta Bot. Hung. 29: 187. 29: 187 (1983). Type:—CUBA. Wright s.n. (Syntypes: GH, K, NY).

Section Uncertain

Distribution: Cuba.

Miconia andersonii Fawc. & Rendle, J. Bot. 64: 104 (1926). Type:—JAMAICA [but probably from St. Vincent].

Anderson s.n. (Syntype: K).

Miconia caribaea Domin, Preslia 9: 3: tab. 1 (1930).

Section: Miconia.

Distribution: Saint Vincent, Grenada.

Miconia andreana Cogn., Bull. Acad. Belg., Ser. 3 14: 952 (1887). *Acinodendron andreadnum* (Cogn.) Kuntze, Revis.

Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *André* 2325 (Holotype: K; isotypes: BR, K).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia androsaemifolia Griseb., Cat. Pl. Cub. 100 (1866). *Acinodendron androsaemifolium* (Griseb.) Kuntze, Revis.

Gen. Pl. 2: 950 (1891). *Tamonea androsaemifolia* (Griseb.) Jenn., Ann. Carnegie Mus. 11: 209 (1917). Type:—CUBA.

Wright 2515 (Holotype: GOET; isotypes: BM, GH, K, MO, NY, S, US).

Section: Amblyarrhena.

Distribution: Cuba.

Miconia angelana R.Romero & R.Goldenb., Novon 9: 98 (1999). Type:—BRAZIL. *Romero* 3773 (Holotype: HUFU; isotypes: K, MO, RB, UEC, US).

Section: Cremanium.

Distribution: Brazil Southeast.

Miconia anisophylla Triana, Trans. Linn. Soc. London 28: 122 (1871). *Acinodendron anisophyllum* (Triana) Kuntze, Revis.

Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Triana* 4040 (Holotype: BM; isotypes: BR, COL, K, NY, US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia anisotricha (Schltdl.) Triana, Trans. Linn. Soc. London 28: 102 (1871).

Basionym: *Melastoma anisotrichon* Schltdl., Linnaea 13 (5): 427 (1839). *Clidemia anisotricha* (Schltdl.) Naudin, Ann.

Sci. Nat., Bot., ser 3 17: 376 (1851). *Acinodendron anisotrichum* (Schltdl.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Type:—MEXICO. *Schiede* s.n. (Holotype: HAL?).

Cremanium berghesianum S.Schauer, Linnaea 20: 735 (1847). *Miconia berghesiana* (S.Schauer) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 246 (1850).

Miconia erythrantha Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 213 (1850). *Acinodendron erythranthum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Clidemia urticifolia Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 354 (1851).

Clidemia monticola Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 341 (1851).

Section: Octomeris.

Distribution: Mexico.

Miconia annulata (Naudin) Triana, Trans. Linn. Soc. London 28: 121 (1871).

Basionym: *Miconia stipularis* var. *annulata* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 195 (1850). *Acinodendron annulatum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Linden* 818 (Holotype: P; isotypes: BR, GH, K, MPU, US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia antioquiensis Wurdack, Caldasia 7 (35): 331 (1957). Type:—COLOMBIA. *Uribe-Uribe* 2697 (Holotype: NY; isotype: COL).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia apiculata Urb. & Ekman, Ark. Bot. 22A (17): 48 (1929). Type:—HAITI. *Ekman H10641* (Holotype: S; isotypes: NY, US).

Miconia niedenzuana Urb. & Ekman, Ark. Bot. 22A (17): 45 (1929).

Section: Chaenopleura.

Distribution: Haiti.

Miconia apostachya (Bonpl.) DC., Prodr. 3: 183 (1828).

Basionym: *Melastoma apostachyum* Bonpl., Monogr. Melast. 1: 1 (1806/1807). *Acinodendron apostachyum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—VENEZUELA. *Bonpland 859* (Holotype: P).

Miconia macrostachya DC., Prodr. 3: 190 (1828). *Acinodendron macrostachyum* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Section: Miconia.

Distribution: Colombia, Venezuela, Guyana, Suriname, French Guiana, Brazil North, Brazil Central-West, Brazil Northeast.

Miconia aponeura Triana, Trans. Linn. Soc. London 28: 106 (1871). *Acinodendron aponeurum* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Triana 4014* (Holotype: BM; isotypes: COL, G, K, P).

Section: Tamonea.

Distribution: Costa Rica, Colombia.

Miconia appendiculata Triana, Trans. Linn. Soc. London 28: 131 (1871). *Acinodendron appendiculatum* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Triana 4059* (Holotype: BM).

Section: Miconia.

Distribution: Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador, Peru.

Miconia aprica Gleason, Bull. Torrey Bot. Club 58: 244 (1931). Type:—PERU. *Killip 24246* (Holotype: NY; isotypes: K, US).

Section: Cremanium.

Distribution: Peru.

Miconia araguensis Wurdack, Phytologia 22: 402 (1972). *Heterotrichum glandulosum* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 955 (1891). Type:—VENEZUELA. *Fendler 432* (Holotype: BR; isotypes: BM, BR, GH, K).

Heterotrichum glandulosum var. *angustifolium* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 955 (1891).

Miconia araguensis var. *angustifolia* (Cogn.) Wurdack, Phytologia 22: 402 (1972).

Heterotrichum glandulosum var. *parvifolium* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 955 (1891).

Section: Octomeris.

Distribution: Colombia, Venezuela.

Miconia arboricola Almeda, Proc. Calif. Acad. Sci., Ser. 4 43: 277 (1984). Type:—PANAMA. *Stevens 18266* (Holotype: CAS; isotypes: MO, PMA).

Section: Miconia.

Distribution: Costa Rica, Panama, Colombia, Ecuador.

Miconia arbutifolia Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 210 (1850). *Acinodendron arbutifolium* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—VENEZUELA. *Funck 739* (Holotype: P; isotypes: BR, P).

Section: Amblyarrhena.

Distribution: Venezuela.

Miconia archeri Wurdack, Phytologia 8: 166 (1962). Type:—COLOMBIA. *Archer 232* (Holotype: US).

Section: Tamonea.

Distribution: Colombia.

Miconia argentea (Sw.) DC., Prodr. 3: 182 (1828).

Basionym: *Melastoma argenteum* Sw., Fl. Ind. Occid. 2: 779 (1800). *Acinodendron argenteum* (Sw.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—"MOSQUITO SHORE" (NICARAGUA?). *Miller s.n.* (Holotype: BM).

Cremanium compressum Benth., Bot. Voy. Sulphur 96 (1844). *Miconia procera* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Colombia.

Miconia argyraea Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 838 (1891). *Acinodendron argyraeum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Glaziou 16905* (Holotype: BR; isotypes: C, F, G, K, P, R).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia argyrophylla DC., Prodr. 3: 181 (1828). *Melastoma argyrophyllum* Schrank & Mart. ex DC., Prodr. 3: 181 (1828), pro syn. *Acinodendron argyrophyllum* (DC.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M).

Miconia longistyla Steud., Flora 27: 724 (1844).

Miconia argyrophylla var. *attenuata* Cogn. in Martius, Fl. bras. 14 (4): 296 (1887).

Miconia argyrophylla ssp. *gracilis* Wurdack, Mem. New York Bot. Gard. 10 (4): 37 (1961).

Section: Miconia.

Distribution: Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Peru, Bolivia, Brazil North, Brazil Central-West.

Miconia aristata Gleason, Bull. Torrey Bot. Club 58: 427 (1931). Type:—VENEZUELA. *Tate 897* (Holotype: NY; isotypes: K, US).

Section: Miconia.

Distribution: Venezuela.

Miconia ascendens Wurdack, Brittonia 9: 106 (1957). Type:—ECUADOR. *Drew E-359* (Holotype: US; isotype: NY).

Section: Chaenopleura.

Distribution: Ecuador.

Miconia asclepiadea Triana, Trans. Linn. Soc. London 28: 121 (1871). *Acinodendron asclepiadeum* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Triana 4001* (Lectotype: BM; isolectotype: COL).

Miconia corazonica Gilli, Feddes Rep. 92: 682 (1981), *nom. illeg.*

Section: Amblyarrhena.

Distribution: Colombia, Peru, Ecuador.

Miconia aspergillaris (Bonpl.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 214 (1850).

Basionym: *Melastoma aspergillare* Bonpl., Monogr. Melast. 1: 44 (1809). *Cremanium aspergillare* (Bonpl.) DC., Prodr. 3: 193 (1828). *Acinodendron aspergillare* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—ECUADOR. *Bonpland s.n.* (Holotype: P).

Miconia mutisella Gleason, Phytologia 1: 47 (1933).

Miconia aspergillaris var. *gracilior* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 215 (1850).

Section: Amblyarrhena.

Distribution: Ecuador, Peru.

Miconia asperrima Triana, Trans. Linn. Soc. London 28: 124 (1871). *Acinodendron asperrimum* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Triana 4042* (Holotype: BM?; isotype: COL).

Miconia pennellii Gleason, Bull. Torrey Bot. Club 52: 448 (1925).

Section: Amblyarrhena.

Distribution: Colombia, Ecuador, Peru.

Miconia asplundii Wurdack, Phytologia 24: 202 (1972). Type:—ECUADOR. Asplund 9854 (Holotype: S).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia aspratilis Wurdack, Mem. New York Bot. Gard. 16: 25 (1967). Type:—ECUADOR. Camp E-704 (Holotype: US; isotype: NY).

Section: Cremanium.

Distribution: Ecuador.

Miconia astroplocama Donn. Sm., Bot. Gaz. 42: 295 (1906). Type:—COSTA RICA. Tonduz 11438 (Holotype: US; isotype: NY).

Sphaerogyne cinnamomea Linden, Belgique Hort. 15: 101 (1865). *Tococa neocinnamomea* Buchheim & Potztal in Encke, Pareys Blumeng. 2: 191 (1960).

Miconia macdanielii Wurdack, Phytologia 21: 119 (1971).

Section: Miconia.

Distribution: Nicaragua, Costa Rica, Panama, Colombia, Venezuela. Ecuador, Peru.

Miconia astrotricha (DC.) Triana, Trans. Linn. Soc. London 28: 113 (1871).

Basionym: *Clidemia astrotricha* DC., Prodr. 3: 160 (1828). *Staphidium astrotrichum* (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 324 (1851). *Acinodendron astrotrichum* (DC.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M; isotypes: BR, G-DC, M).

Melastoma agreste Schrank & Mart. ex DC., Prodr. 3: 160 (1828), *pro syn.*

Section: Miconia.

Distribution: Colombia, Brazil North.

Miconia atrofusca Cogn., Bot. Jahrb. Syst. 42: 142 (1909). Type:—PERU. Weberbauer 3726 (Holotype: B, destroyed; isotypes: BR, MOL).

Section: Cremanium.

Distribution: Peru.

Miconia atropilis Cogn. & Gleason, Brittonia 1: 114 (1932). Type:—COLOMBIA. Lehmann 6216 (Holotype: K?; isotype: BR).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia augusti Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 835 (1891). *Acinodendron augusti* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. Glaziou 16895 (Holotype: BR; isotypes: C, G, K, P, R).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia aulocalyx Mart. ex Triana, Trans. Linn. Soc. London 28: 115 (1871). *Acinodendron aulocalyx* (Mart. ex Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. Poeppig 2308 (Holotype: B, destroyed; isotypes: BM, BR, NY).

Section: Miconia.

Distribution: Ecuador, Peru, Brazil North.

Miconia aurea (D.Don) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850).

Basionym: *Chitonia aurea* D.Don, Mem. Wern. Nat. Hist. Soc. 4: 319 (1823). *Diplochita aurea* (D.Don) DC., Prodr. 3: 177 (1828). *Acinodendron aureum* (D.Don) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. Pavón s.n. (Holotype: BM ex Herb. Lamb.?; isotype: B, destroyed).

Miconia adenophora Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 182 (1850).

Miconia aurea var. *latifolia* Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 9: 1144 (1927).

Section: Tamonea.

Distribution: Colombia, Ecuador, Peru, Brazil North.

Miconia aureoides Cogn. in Martius, Fl. bras. 14 (4): 260 pl. 52 (1887). *Acinodendron aureoides* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. Spruce 4519 (Holotype: BR; isotypes: BM, BR, E, F, GH, K, NY).
Section: Tamonea.
Distribution: Colombia, Ecuador, Peru, Bolivia, Brazil North.

Miconia auritinoda Wurdack, Phytologia 64: 298 (1988). Type:—ECUADOR. Lutelyn 11007 (Holotype: NY; isotypes: CAS, QCA, US).
Section: Cremanium.
Distribution: Ecuador.

Miconia avia Wurdack, Acta Bot. Venez. 13: 144 (1978). Type:—VENEZUELA. Wurdack 2766 (Holotype: US; isotypes: F, MICH, MO, NY, VEN).
Section: Octomeris.
Distribution: Venezuela.

Miconia axinaeoides Gleason, Bull. Torrey Bot. Club 59: 368 (1932). Type:—BOLIVIA. Buchtien 1099 (Holotype: US; isotype: NY).
Section: Octomeris.
Distribution: Peru, Bolivia.

Miconia ayacuchensis Wurdack, Phytologia 23: 497 (1972). Type:—PERU. Dudley 11992 (Holotype: US).
Section: Cremanium.
Distribution: Peru.

Miconia aymardii Wurdack, Phytologia 69: 319 (1990). Type:—VENEZUELA. Aymard 4412 (Holotype: PORT; isotypes: US, VEN).
Section: Amblyarrhena.
Distribution: Venezuela.

Miconia bailloniana J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (4): 388 (1941). *Pteroclodon sprucei* Hook.f. in Bentham & Hooker f., Gen. Pl. 1: 763 (1867). *Muelleramra sprucei* (Hook.f.) Kuntze, Revis. Gen. Pl. 1: 247 (1891). Type:—PERU. Spruce 4312 (Holotype: K; isotypes: GH, NY).
Section: Amblyarrhena.
Distribution: Peru.

Miconia bangii Cogn. in Rusby, Mem. Torrey Bot. Club 3: 30 (1893). Type:—BOLIVIA. Bang 282 (Holotype: BR; isotypes: F, GH, M, NY, PH, US, WIS).
Section: Amblyarrhena.
Distribution: Peru, Bolivia.

Miconia baracoensis Urb., Symb. Antill. 9: 112 (1923). Type:—CUBA. Ekman 4221 (Holotype: S; isotypes: NY, US).
Section: Octomeris.
Distribution: Cuba.

Miconia barbeyana Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 872 (1891). *Acinodendron barbeyanum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. Pavón s.n. (Holotype: G-BOISS; isotype: BR).
Section: Amblyarrhena.
Distribution: Bolivia, Ecuador, Peru.

Miconia barbinervis (Benth.) Triana, Trans. Linn. Soc. London 28: 113 (1871).
Basionym: *Clidemia barbinervis* Benth., Bot. Voy. Sulphur 95 (1844). *Acinodendron barbinerve* (Benth.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. Sinclair s.n. (Syntype: K).

Miconia stellifera Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 142 (1850).

Clidemia chrysopogon Beurl., Kongl. Vetensk. Acad. Handl. 40: 127 (1854).

Section: Miconia.

Distribution: Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Trinidad & Tobago, Ecuador, Peru, Bolivia, Brazil North.

Miconia barbipilis Gleason, Phytologia 2: 303 (1947). Type:—ECUADOR. Steyermark 54308 (Holotype: F).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia barclayana Wurdack, Phytologia 39: 327 (1978). Type:—ECUADOR. Barclay 8936 (Holotype: US).

Section: Cremanium.

Distribution: Ecuador.

Miconia barkeri Urb. & Ekman, Ark. Bot. 21A (5): 41 (1927). Type:—HAITI. Ekman H5329 (Holotype: S; isotypes: C, EHH, NY, US).

Section: Chaenopleura.

Distribution: Haiti.

Miconia basileensis Urb. & Ekman, Ark. Bot. 22A (17): 44 (1929). Type:—HAITI. Ekman H9663 (Holotype: S; isotypes: NY, US).

Section: Chaenopleura.

Distribution: Haiti.

Miconia baumgratziana R.Goldenb. & C.V.Martin, Harvard Pap. Bot. 13: 223 (2008). *Miconia angustifolia* Cogn. in Martius, Fl. bras. 14 (4): 613: 613 (1888), nom. illeg. *Acinodendron angustifolium* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. Glaziou 16878 (Lectotype: BR; isolectotypes: K, P).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia bella Wurdack, Phytologia 38: 288 (1978). Type:—COLOMBIA. García-Barriga 16122 (Holotype: US; isotype: COL).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia beneolens Wurdack, Phytologia 13: 77 (1966). Type:—ECUADOR. Steyermark 54789 (Holotype: US; isotypes: F, NY).

Section: Amblyarrhena.

Distribution: Ecuador, Peru.

Miconia benoistii Wurdack, Phytologia 41: 4 (1978). Type:—ECUADOR. Benoit 2504 (Holotype: P).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia benthamiana Triana, Trans. Linn. Soc. London 28: 102 (1871). *Clidemia polyandra* Benth., Bot. Voy. Sulphur 95 (1844). *Octomeris polyandra* (Benth.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 382 (1851). *Staphidium polyandrum* (Benth.) Griseb. in Lechler, Berberid. Amer. Austral. 59 (1857). *Heterotrichum polyandrum* (Benth.) Gleason, Bull. Torrey Bot. Club 52: 378 (1925). *Acinodendron benthamianum* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. Sinclair s.n. (Holotype: K).

Clidemia tonduzii Gleason, Brittonia 3: 116 (1939).

Section: Octomeris.

Distribution: Costa Rica, Panama, Colombia, Ecuador, Peru.

Miconia bernardii Wurdack, Phytologia 22: 413 (1972). Type:—VENEZUELA. Bernardi 3209 (Holotype: NY; isotypes: BM, G-DEL, NY, US, VEN).

Section: Cremanium.

Distribution: Venezuela.

Miconia berryi Wurdack, Acta Bot. Venez. 13: 158 (1978). Type:—VENEZUELA. *Berry 132* (Holotype: US).

Section: Cremanium.

Distribution: Venezuela.

Miconia biacuta Cogn. in Rusby, Mem. Torrey Bot. Club 3: 31 (1893). Type:—BOLIVIA. *Bang 701* (Holotype: BR; isotypes: GH, K, MO, NY, PH, US, WIS).

Section: Cremanium.

Distribution: Bolivia.

Miconia biappendiculata (Naudin) L.Uribe, Caldasia 8: 326 (1960).

Basionym: *Miconia granulosa* var. *biappendiculata* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 218 (1850). Type:—COLOMBIA. *Bonpland s.n.* (Holotype: P).

Melastoma granulosum Bonpl., Monogr. Melast. 1: 25 (1807), *nom illeg.* *Cremanium granulosum* (Bonpl.) DC., Prodr. 3: 192 (1828). *Miconia granulosa* (Bonpl.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 218 (1850). *Acinodendron granulosum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Melastoma revoluta Dietr. ex Steud., Nomencl. Bot., ed. 2: 117 (1841).

Section: Cremanium.

Distribution: Colombia, Bolivia.

Miconia bifaria Urb. & Ekman, Ark. Bot. 23A (11): 25 (1931). Type:—DOMINICAN REPUBLIC. *Ekman H12265* (Holotype: S).

Section: Chaenopleura.

Distribution: Dominican Republic.

Miconia biformis Cogn., Meded. Rijks-Herb. 19: 80 (1913). Type:—BOLIVIA. *Herzog 1573* (Holotype: BR; isotype: JE).

Miconia biformis var. *brevifolia* Cogn., Meded. Rijks-Herb. 19: 81 (1913).

Section: Cremanium.

Distribution: Bolivia.

Miconia biglandulosa Gleason, Bull. Torrey Bot. Club 59: 367 (1932). Type:—PERU. *Killip 26933* (Holotype: NY; isotypes: F, US).

Section: Adenodesma.

Distribution: Colombia, Venezuela, Ecuador, Peru, Brazil North.

Miconia biglomerata (Bonpl.) DC., Prodr. 3: 183 (1828).

Basionym: *Melastoma biglomeratum* Bonpl., Monogr. Melast. 1: 34 (1807). *Acinodendron biglomeratum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—VENEZUELA. *Bonpland 1160 bis* (Holotype: P).

Clidemia campestris var. *pauciflora* Benth., J. Bot. 2: 308 (1840).

Section: Miconia.

Distribution: Venezuela.

Miconia bilopezii Wurdack, Phytologia 22: 412 (1972). Type:—VENEZUELA. *Figueiras 8688* (Holotype: US; isotype: MER).

Section: Cremanium.

Distribution: Venezuela.

Miconia bipatrialis Wurdack, Mem. New York Bot. Gard. 16: 19 (1967). Type:—ECUADOR: *Camp E-4631* (Holotype: US; isotypes: COL, F, GH, NY).

Miconia bipatrialis ssp. *peruviana* Wurdack, Mem. New York Bot. Gard. 16: 20 (1967).

Section: Amblyarrhena.

Distribution: Ecuador, Peru.

Miconia biperulifera Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 911 (1891). *Acinodendron biperuliferum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COSTA RICA. Pittier s.n. (Holotype: BR; isotype: US).
Miconia biperulifera var. *rigida* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 1192 (1891).
Section: Cremanium.
Distribution: Costa Rica, Panama.

Miconia bisulcata Urb., Repert. Spec. Nov. Regni Veg. 22: 228 (1926). Type:—CUBA. Ekman 14707 (Holotype: S; isotypes: GH, NY, US).
Section: Glossocentrum.
Distribution: Cuba.

Miconia blakeifolia Gleason, Bull. Torrey Bot. Club 72: 475 (1945). Type:—COLOMBIA. Cuatrecasas 15181 (Holotype: NY; isotypes: COL, F, US).
Section: Tamonea.
Distribution: Colombia.

Miconia bolivarensis Wurdack, Phytologia 26: 8 (1973). Type:—ECUADOR. Asplund 8328 (Holotype: S; isotypes: A, NY).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia boliviensis Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 901 (1891). *Acinodendron boliviense* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BOLIVIA. Mandon 643 (Syntype: P; isosyntypes: A, GH, P, S).
Miconia boliviensis var. *glabra* Cogn. in Rusby, Mem. Torrey Bot. Club 3: 30 (1893).
Section: Cremanium.
Distribution: Bolivia.

Miconia boomii Wurdack, Brittonia 40: 9 (1988). Type:—BOLIVIA. Boom 4478 (Holotype: NY; isotypes: F, GH, K, MO, U, US).
Section: Jucunda.
Distribution: Bolivia.

Miconia bordoncilloana Lozano & Morales-P., Revista Acad. Colomb. Ci. Exact. 23: 85 (1999). Type:—COLOMBIA. Muñoz 60 (Holotype: COL; isotype: COL).
Section: Cremanium.
Distribution: Colombia, Ecuador.

Miconia borjensis Wurdack, Mem. New York Bot. Gard. 10 (4): 40 (1961). Type:—VENEZUELA. Wurdack 39827 (Holotype: NY; isotypes: F, K, RB, S, U, US, VEN).
Section: Miconia.
Distribution: Colombia, Venezuela.

Miconia boxii Wurdack, Phytologia 20: 386 (1970). Type:—VENEZUELA. Box 3745 (Holotype: BM; isotypes: VEN).
Section: Amblyarrhena.
Distribution: Venezuela.

Miconia brachyanthera Triana, Trans. Linn. Soc. London 28: 129 (1871). *Acinodendron brachyantherum* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. Mathews 1726 (Holotype: K; isotypes: BR, K).
Section: Cremanium.
Distribution: Peru.

Miconia brachybotrya Triana, Trans. Linn. Soc. London 28: 111 (1871). *Acinodendron brachybotryum* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—VENEZUELA. Spruce 3383 (Syntypes: BM, BR, GH, K, NY, TCD).
Section: Miconia.
Distribution: Panama, Venezuela, Guyana, Ecuador, Peru.

Miconia brachycalyx Triana, Trans. Linn. Soc. London 28: 102 (1871). *Acinodendron brachycalyx* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Triana* 5955 (Holotype: BM; isotypes: BR, COL).
Section: Octomeris.
Distribution: Colombia, Ecuador.

Miconia brachygyna Gleason, Bull. Torrey Bot. Club 57: 72 (1930). Type:—COLOMBIA. *Killip* 15266 (Holotype: NY; isotypes: CAS, GH, K, US).
Section: Cremanium.
Distribution: Colombia, Venezuela.

Miconia bracteata (DC.) Triana, Trans. Linn. Soc. London 28: 111 (1871).
Basionym: *Clidemia bracteata* DC., Prodr. 3: 162 (1828). *Acinodendron bracteatum* (DC.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—FRENCH GUIANA. *Richard* s.n. (Holotype: P).
Clidemia trichodes DC., Prodr. 3: 161 (1828). *Miconia trichodes* (DC.) Triana, Trans. Linn. Soc. London 28: 112 (1871).
Acinodendron trichodes (DC.) Kuntze, Revis. Gen. Pl. 2: 953 (1891).

Melastoma triste Rich. ex DC., Prodr. 3: 162 (1828), *pro syn.* *Staphidium triste* (Rich. ex DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 320 (1851).

Clidemia cephalophora Steud. ex Triana, Trans. Linn. Soc. London 28: 111 (1871), *pro syn.*

Clidemia interrupta Poepp. ex Triana, Trans. Linn. Soc. London 28: 111 (1871), *pro syn.*

Miconia demerarensis Gleason, Bull. Torrey Bot. Club 75: 551 (1948).

Section: Miconia.

Distribution: Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Brazil North.

Miconia bracteolata (Bonpl.) DC., Prodr. 3: 182 (1828).
Basionym: *Melastoma bracteolatum* Bonpl., Monogr. Melast. 1: 46 (1809). *Acinodendron bracteolatum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Bonpland* 3366 (Holotype: P; isotypes: BM, G).

Section: Cremanium.

Distribution: Colombia, Ecuador, Peru.

Miconia brasiliensis (Spreng.) Triana, Trans. Linn. Soc. London 28: 118 (1871).
Basionym: *Melastoma brasiliense* Spreng., Syst. Veg. 2: 297 (1825). *Charianthus brasiliensis* (Spreng.) Cham. ex Triana, Trans. Linn. Soc. London 28: 118 (1871), *pro. syn.* *Acinodendron brasiliense* (Spreng.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Sello* 743 (Lectotype: P).

Miconia pauciflora Triana, Trans. Linn. Soc. London 28: 115 (1871). *Acinodendron pauciflorum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Miconia venustula Benth. ex Triana, Trans. Linn. Soc. London 28: 118 (1871), *pro syn.*

Section: Glossocentrum.

Distribution: Brazil Southeast, Brazil South.

Miconia brenesii Standl., Publ. Field Mus. Nat. Hist., Bot. Ser. 18 (3): 816 (1938). Type:—COSTA RICA. *Brenes* 21981 (Holotype: F; isotypes: CR, NY).
Section: Cremanium.
Distribution: Costa Rica.

Miconia breteleri Wurdack, Phytologia 22: 400 (1972). Type:—VENEZUELA. *Breteler* 4910 (Holotype: US; isotypes: F, K, L, M, MO, WAG, S, U, UC, US).

Section: Octomeris.

Distribution: Venezuela.

Miconia brevipes Benth., J. Bot. 2: 313 (1840). *Acinodendron brevipes* (Benth.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).
Type:—GUYANA. *Schomburgk I add. 125.S* (Holotype: K).

Miconia plebeia Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 170 (1850).

Miconia rubiginosa var. *kuhlmannii* Hoehne, Anexos Mem. Inst. Butantan, Secc. Bot. 1 (5): 138 (1922).

Section: Miconia.

Distribution: Venezuela, Guyana, Suriname, Brazil North.

Miconia brevis J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 188 (1929). Type:—PERU. *Weberbauer* 6334 (Holotype: F; isotype: US).
Section: Cremanium.
Distribution: Peru.

Miconia brevistylis Cogn., Bot. Jahrb. Syst. 42: 144 (1909). Type:—PERU. *Weberbauer* 3553 (Holotype: B, destroyed; isotypes: BR, MOL).
Section: Cremanium.
Distribution: Peru.

Miconia brevitheca Gleason, Brittonia 2: 322 (1937). Type:—ECUADOR. *Mexia* 6730 (Holotype: NY; isotypes: K, UC, US).
Section: Cremanium.
Distribution: Costa Rica, Panama, Colombia, Ecuador.

Miconia brittonii Cogn. in Britton, Bull. Torrey Bot. Club 17: 92 (1890). *Acinodendron brittonii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BOLIVIA. *Rusby* 2309 (Syntype: NY; isosyntypes: MICH, US).
Miconia brittonii var. *glabrata* Cogn., Bull. Torrey Bot. Club 23: 277 (1896).
Section: Cremanium.
Distribution: Bolivia.

Miconia brunnea DC., Prodr. 3: 184 (1828). *Melastoma brunneum* Mart. ex DC., Prodr. 3: 184 (1828), *pro syn.* *Acinodendron brunneum* (DC.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Martius* s.n. (Holotype: M; isotype: G-DC).
Section: Glossocentrum.
Distribution: Brazil Southeast, Brazil South.

Miconia bubalina (D.Don) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850).
Basionym: *Chitonia bubalina* D.Don, Mem. Wern. Nat. Hist. Soc. 4: 319 (1823). *Melastoma bubalinum* Pav. ex D.Don, Mem. Wern. Nat. Hist. Soc. 4: 319 (1823), *pro syn.* *Diplochita bubalina* (D.Don) DC., Prodr. 3: 177 (1828).
Acinodendron bubalinum (D.Don) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. *Pavón* s.n. (Holotype: BM).

Miconia belizensis Standl., Publ. Field Mus. Nat. Hist., Bot. Ser. 8 (1): 30 (1930).
Section: Tamonea.
Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Trinidad & Tobago, Guyana, Ecuador, Peru, Bolivia, Brazil North.

Miconia bucherae Alain, Contr. Ocas. Mus. Hist. Nat. Colegio "De La Salle" 14: 2 (1955). Type:—CUBA. *Bucher* 124 (Holotype: NY; isotype: HAC).
Section: Octomeris.
Distribution: Cuba.

Miconia bubblejoides Triana, Trans. Linn. Soc. London 28: 118 (1871). *Acinodendron bubblejoides* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Pohl* 99-1165 (Syntypes: BM, BR, F, K, M, NY).
Section: Glossocentrum.
Distribution: Brazil Northeast, Brazil Southeast, Brazil South.

Miconia bullata (Turcz.) Triana, Trans. Linn. Soc. London 28: 130 (1871).
Basionym: *Schizanthera bullata* Turcz., Bull. Soc. Nat. Mosc. 35 (4): 322 (1862). *Acinodendron bullatum* (Turcz.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. *Mathews* 873 (Holotype: K; isotypes: BM, E).
Miconia trichocaula J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 189 (1929).
Section: Chaenopleura.
Distribution: Ecuador, Peru.

Miconia burchellii Triana, Trans. Linn. Soc. London 28: 116 (1871). *Acinodendron burchellii* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Burchell* 7918 (Syntype: K).
Miconia burchellii var. *glabrata* Cogn. ex Glaz., Bull. Soc. Bot. France, Mem. 54 (3c): 288 (1905).
Section: Miconia.
Distribution: Brazil North, Brazil Central-West, Brazil Southeast.

Miconia buxifolia Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 208 (1850). *Acinodendron buxifolium* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Funck* 1316 (Holotype: P; isotypes: BM, BR, G, P).
Section: Cremanium.
Distribution: Colombia, Venezuela.

Miconia cabucu Hoehne, Ostenia 1: 299 pls. 4-6 (1933). Type:—BRAZIL. *Hoehne* 28306 (Holotype: SP; isotypes: A, F, NY, US, S).
Section: Glossocentrum.
Distribution: Brazil Southeast, Brazil South.

Miconia cacatin (Aubl.) S.S.Renner, Mem. New York Bot. Gard. 50: 29 (1989).
Basionym: *Melastoma cacatin* Aubl., Hist. pl. Guiane 1: 437: (1775). *Blakea cacatin* (Aubl.) D.Don ex DC., Prodr. 3: 196 (1828). *Bellucia cacatin* (Aubl.) Sagot, Ann. Sci. Nat., Bot., Ser.6, 15: 332 (1883). Type:—FRENCH GUIANA. *Aublet s.n.* (Holotype: BM).
Section: Tamonea.
Distribution: Suriname, French Guiana.

Miconia cacumina Wurdack, Phytologia 64: 296 (1988). Type:—GUYANA. *Edwards* 107 (Holotype: K; isotypes: K, NY, U, US).
Section: Amblyarrhena.
Distribution: Venezuela, Guyana.

Miconia caelata (Bonpl.) DC., Prodr. 3: 183 (1828).
Basionym: *Melastoma caelatum* Bonpl., Monogr. Melast. 1: 11 (1807). *Acinodendron caelatum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—ECUADOR. *Bonpland s.n.* (Holotype: P; isotypes: BM, P).
Section: Cremanium.
Distribution: Ecuador, Peru.

Miconia caerulea (D.Don) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).
Basionym: *Cremanium caeruleum* D.Don, Mem. Wern. Nat. Hist. Soc. 4: 313 (1823). *Melastoma caeruleum* Pav. ex D.Don, Mem. Wern. Nat. Hist. Soc. 4: 313 (1823), *pro syn.* *Acinodendron caeruleum* (D.Don) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. *Ruiz & Pavón s.n.* (Holotype: OXF?; isotypes: B, destroyed, FI, G, MA?).

Miconia agyna Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 232 (1850).
Miconia caerulea ssp. *vilcabambae* Wurdack, Phytologia 23: 491 (1972).
Section: Cremanium.
Distribution: Peru.

Miconia caesariata Wurdack, Phytologia 38: 294 (1978). Type:—ECUADOR. *Steyermark* 53573 (Holotype: F; isotype: US).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia caesia Cogn. & Gleason, Brittonia 1: 116 (1932). Type:—COLOMBIA. *Lehmann* 7505 (Holotype: K; isotype: BR).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia cajanumana Wurdack, Mem. New York Bot. Gard. 16: 17 (1967). Type:—ECUADOR. *Maguire* 44370 (Holotype: NY; isotype: US).
Section: Amblyarrhena.
Distribution: Ecuador, Peru.

Miconia calignosa Wurdack, Phytologia 38: 296 (1978). Type:—ECUADOR. *Sparre* 18730 (Holotype: S).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia calocoma Almeda, Proc. Calif. Acad. Sci., Ser. 4 46: 144 (1989). Type:—COSTA RICA. *Folsom* 9776 (Holotype: CAS; isotypes: DUKE, MO).
Section: Amblyarrhena.
Distribution: Costa Rica.

Miconia calvescens DC., Prodr. 3: 185 (1828). *Melastoma calvescens* Schrank & Mart. ex DC., Prodr. 3: 185 (1828), *pro syn.* Type:—BRAZIL. *Martius s.n.* (Holotype: M).

Melastoma mandiocanum Raddi, Mem. Soc. Ital. Modena 20: 132: t. 6 (1829).

Melastoma arboreum Vell., Fl. Flum. 4: 182 (1825/1829). *Acinodendron arboreum* (Vell.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Cyanophyllum magnificum Groenl., Rev. Hort., Ser. 4 8: 359 (1859). *Miconia magnifica* (Groenl.) Triana, Trans. Linn. Soc. London 28: 131 (1871).

Miconia arborea Pav. ex Triana, Trans. Linn. Soc. London 28: 108 (1871), *pro syn.*

Miconia velutina L.Linden & Rodigas, Ill. Hort. 41: 331 (1894). *Miconia magnifica* var. *velutina* Rodigas, Ill. Hort. 41: 332 (1894), *pro syn.*

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Panama, Colombia, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast, Brazil South, Paraguay, Argentina; Invasive in Australia, Hawaiian Is., Society Is., Sri Lanka.

Miconia calycina Cogn. in Urban, Symb. Antill. 7: 312 (1912). Type:—HAITI. *Christ* 2183 (Holotype: B, destroyed; isotype: GH).

Miconia cubensis var. *nervulosa* Cogn. in A.DC. & P.DC., Monogr. Phan. 7: 933 (1891).

Miconia cubensis var. *latifolia* Cogn., Jahrb. Königl. Bot. Gart. Berlin 4: 281 (1886).

Miconia cubensis var. *minor* Cogn., Jahrb. Königl. Bot. Gart. Berlin 4: 281 (1886).

Miconia subcorymbosa Britton, Sci. Surv. Porto Rico & Virgin Islands 6: 10 (1925).

Miconia haitiensis Urb. & Ekman, Ark. Bot. 22A (17): 47 (1929).

Section: Chaenopleura.

Distribution: Cuba, Haiti, Dominican Republic, Puerto Rico.

Miconia campanensis Urb. & Ekman, Ark. Bot. 23A (11): 18 (1931). Type:—DOMINICAN REPUBLIC. *Ekman* H11514 (Holotype: S).

Section: Chaenopleura.

Distribution: Dominican Republic.

Miconia campestris (Benth.) Triana, Trans. Linn. Soc. London 28: 113 (1871).

Basionym: *Clidemia campestris* Benth., J. Bot. 2: 308 (1840). *Acinodendron campestre* (Benth.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—GUYANA. *Schomburgk* I 478 (Holotype: K).

Section: Miconia.

Distribution: Venezuela, Guyana, Suriname, Brazil North.

Miconia campii Wurdack, Mem. New York Bot. Gard. 16: 34 (1967). Type:—ECUADOR. *Camp E-5117* (Holotype: US; isotypes: F, GH, MO, NY, U, VEN).

Section: Chaenopleura.

Distribution: Ecuador.

Miconia canaguensis Wurdack, Acta Bot. Venez. 13: 143 (1978). Type:—VENEZUELA. *Wurdack 2744* (Holotype: US; isotypes: NY, VEN).
Section: Octomeris.
Distribution: Venezuela.

Miconia cannabina Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 9: 1145 (1927). Type:—PERU. *Tessmann 5056* (Holotype: B, destroyed; isotype: NY).
Section: Miconia.
Distribution: Colombia, Ecuador, Peru, Brazil North.

Miconia capitellata Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 892 (1891). *Acinodendron capitellatum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—ECUADOR. *Poortmann 162* (Holotype: P; isotype: BR).
Section: Amblyarrhena.
Distribution: Colombia, Ecuador.

Miconia capixaba R.Goldenb., Novon 9: 514 (1999). Type:—BRAZIL. *Thomaz s.n.* (Holotype: MBML; isotypes: CEPEC, K, MO, UEC).
Section: Miconia.
Distribution: Brazil Southeast.

Miconia carassana Cogn. in Martius, Fl. bras. 14 (4): 346 (1887). *Acinodendron carassanum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M; isotype: BR).
Miconia compacta Gleason, Bull. Torrey Bot. Club 58: 230 (1931).
Miconia semota Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 12: 177 (1934).
Section: Miconia.
Distribution: Colombia, Venezuela, Guyana, Ecuador, Peru, Brazil North.

Miconia carneae Cogn., Bot. Jahrb. Syst. 8: 28 (1886). *Acinodendron carneum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COSTA RICA. *Lehmann 38* (Syntype: BR).
Section: Cremanium.
Distribution: Costa Rica, Panama.

Miconia carpishana Wurdack, Phytologia 13: 76 (1966). Type:—PERU. *Maguire 44436* (Holotype: US; isotype: NY).
Section: Amblyarrhena.
Distribution: Peru.

Miconia carvalhoana Baumgratz & D'El Rei Sousa, Bradea 10: 28 (2004). Type:—BRAZIL. *Melo 1692* (Holotype: ALCB; isotypes: K, RB, SPF).
Section uncertain.
Distribution: Brazil Northeast.

Miconia castaneiflora Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 132 (1850). *Acinodendron castaneiflorum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Claussen 334* (Holotype: P).
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia castillensis Wurdack, Mem. New York Bot. Gard. 16: 30 (1967). Type:—ECUADOR. *Camp E-1635* (Holotype: US; isotypes: GH, NY).
Section: Cremanium.
Distribution: Ecuador.

Miconia castrensis Wurdack, Mem. New York Bot. Gard. 16: 15 (1967). Type:—ECUADOR. *Camp E-4728* (Holotype: US; isotypes: B, F, GH, MICH, MO, NY, RB, UC).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia cataractae Triana, Trans. Linn. Soc. London 28: 123 (1871). *Acinodendron cataractae* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Holton s.n.* (Holotype: K; isotypes: BR, NY).
Section: Amblyarrhena.
Distribution: Colombia, Venezuela.

Miconia caucana Gleason, Bull. Torrey Bot. Club 52: 385 (1925). Type:—COLOMBIA. *Holton 909* (Holotype: NY; isotypes: K, PH).
Section: Miconia.
Distribution: Colombia.

Miconia caudata (Bonpl.) DC., Prodr. 3: 187 (1828).
Basionym: *Melastoma caudatum* Bonpl., Monogr. Melast. 1: 13 (1807). *Acinodendron caudatum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Humboldt & Bonpland s.n.* (Holotype: P).
Miconia densiflora Cogn., Bot. Jahrb. Syst. 8: 22 (1886), *nom. illeg.*
Section: Tamonea.
Distribution: Mexico, Guatemala, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Colombia.

Miconia caudigera DC., Prodr. 3: 186 (1828). *Acinodendron caudigerum* (DC.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).
Type:—BRAZIL. *Martius s.n.* (Holotype: M; isotypes: BR, G-DC).
Melastoma caudatum Schrank & Mart. ex DC., Prodr. 3: 186 (1828), *pro syn.*
Section: Miconia.
Distribution: Brazil Northeast, Brazil Southeast.

Miconia cauingia J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (4/1): 394 (1941). Type:—PERU. *Stork 9992* (Holotype: F).
Section: Chaenopleura.
Distribution: Peru.

Miconia cautis Wurdack, Mem. New York Bot. Gard. 10 (5): 169 (1964). Type:—VENEZUELA. *Maguire 37498* (Holotype: US; isotypes: K, NY, VEN).
Section: Miconia.
Distribution: Colombia, Venezuela, Brazil North.

Miconia cazaletii Wurdack, Phytologia 35: 247 (1977). Type:—ECUADOR. *Cazalet 7523* (Holotype: US).
Section: Miconia.
Distribution: Colombia, Ecuador, Peru.

Miconia celaquensis Almeda, Novon 6: 319 (1996). Type:—HONDURAS. *Hawkins 329* (Holotype: CAS; isotypes: EAP, HEH, MO, TEFH).
Section: Octomeris.
Distribution: Honduras.

Miconia centrodesma Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 164 (1850). *Acinodendron centrodesmum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Luschnath 120 in Martius Herb Fl Bras.* 498 (Holotype: P; isotypes: BM, BR, CAS, K, M, MO, NY).

Oxymeris ciliata Mart., Flora 24 (2, Beibl.): 38 (1841). *Clidemia martii* Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 375 (1851).

Miconia buchtienii Cogn., Repert. Spec. Nov. Regni Veg. 8: 2 (1910).

Miconia centrandra Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 361 (1915).

Miconia subtriloba Gleason, Bull. Torrey Bot. Club 58: 429 (1931).

Miconia sylvestris Gleason, Brittonia 2: 321 (1937).

Section: Laceraria.

Distribution: Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Trinidad & Tobago, Guyana, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Northeast, Brazil Southeast.

Miconia centrodesmoides Wurdack, Phytologia 26: 4 (1973). Type:—COLOMBIA. *Robinson* 136 (Holotype: US; isotypes: COL, K).
Section: Laceraria.
Distribution: Colombia, Ecuador, Peru.

Miconia centronioides Gleason, Bull. Torrey Bot. Club 72: 475 (1945). Type:—COLOMBIA. *Cuatrecasas* 14145 (Holotype: NY; isotypes: COL, F, US).
Section: Octomeris.
Distribution: Panama, Colombia, Ecuador.

Miconia centrophora Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 209 (1850). *Acinodendron centrophorum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. *Mathews* 1263 (Holotype: P; isotypes: BR, G, NY, P).
Section: Amblyarrhena.
Distribution: Ecuador, Peru.

Miconia centrosperma Almeda, Brittonia 35: 42 (1983). Type:—PANAMA. *Liesner* 702 (Holotype: CAS; isotypes: BM, MO, PMA, US).
Section: Amblyarrhena.
Distribution: Panama.

Miconia ceramicarpa (DC.) Cogn. in Martius, Fl. bras. 14 (4): 338 (1887).
Basionym: *Clidemia ceramicarpa* DC., Prodr. 3: 160 (1828). *Melastoma ceramicarpum* Schrank & Mart. ex DC., Prodr. 3: 160 (1828), *pro syn.* *Acinodendron ceramicarpum* (DC.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M).

Clidemia coccinea DC., Prodr. 3: 162 (1828).

Clidemia violacea DC., Prodr. 3: 162 (1828). *Melastoma violaceum* Rich. ex DC., Prodr. 3: 162 (1891), *pro syn.* *Miconia ceramicarpa* var. *violacea* (DC.) Cogn. in Martius, Fl. bras. 14 (4): 339 (1887).

Miconia erythropila Steud., Flora 27: 723 (1844).

Miconia iodopila Steud., Flora 27: 723 (1844).

Miconia spondylantha var. *leschenaultiana* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 188 (1850).

Clidemia carassana DC. ex Triana, Trans. Linn. Soc. London 28: 111 (1871), *pro syn.*

Miconia ceramicarpa var. *candolleana* Cogn. in Martius, Fl. bras. 14 (4): 339 (1887).

Miconia ceramicarpa var. *navioensis* Wurdack, Mem. New York Bot. Gard. 10 (4): 40 (1961).

Miconia ceramicarpa var. *crozieriae* Cremers & C.V.Martin, Novon 12: 375 (2002).

Section: Miconia.

Distribution: Venezuela, Guyana, Suriname, French Guiana, Brazil North, Brazil Northeast.

Miconia cerasiflora Urb., Symb. Antill. 9: 112 (1923). Type:—CUBA. *Ekman* 5696 (Holotype: S; isotypes: NY, US).

Miconia cerasiflora var. *setulifera* Urb., Symb. Antill. 9: 113 (1923).

Section: Octomeris.

Distribution: Cuba.

Miconia cercophora Wurdack, Phytologia 13: 74 (1966). Type:—ECUADOR. *Asplund* 19526 (Holotype: S).

Miconia cercophora ssp. *canelosana* Wurdack, Phytologia 26: 397 (1973).

Miconia cercophora ssp. *esetulosa* Wurdack, Phytologia 35: 245 (1977).

Section: Tamonea.

Distribution: Ecuador.

Miconia cernua Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 205 (1850). *Acinodendron cernuum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—VENEZUELA. *Funck* 1553 (Holotype: P; isotypes: BR, G, LE, P, US).

Section: Cremanium.

Distribution: Venezuela.

Miconia chaetodon Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 134 (1850). *Acinodendron chaetodon* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—VENEZUELA. *Linden* 266 (Holotype: P; isotypes: BR, P).
Section: Miconia.
Distribution: Venezuela.

Miconia chamissois Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 179 (1850). *Oxymeris calyprata* Cham. ex Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 179 (1850), *pro syn.* *Acinodendron chamissois* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891).
Type:—BRAZIL. *Sello s.n.* (Lectotype: P; isolectotype: P).

Melastoma calypratum Schrank & Mart ex Triana, Trans. Linn. Soc. London 28: 109 (1871), *pro syn.*

Miconia calyprata Mart. ex Triana, Trans. Linn. Soc. London 28: 109 (1871), *pro syn.*

Miconia langlassaei Standl., Contr. U.S. Natl. Herb. 23: 1068 (1924).

Miconia panamensis Gleason, Phytologia 1: 341 (1939).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Bolivia, Brazil Central-West, Brazil Northeast, Brazil Southeast, Brazil South, Paraguay, Argentina.

Miconia chapensis E.Cotton & W.Meier, Willdenowia 33: 201 (2003). Type:—VENEZUELA. *Meier* 8468 (Holotype: VEN; isotypes: AAU, B, MO, MY, NY, PORT).

Section: Tamonea.

Distribution: Venezuela.

Miconia chartacea Triana, Trans. Linn. Soc. London 28: 119 (1871). *Acinodendron chartaceum* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Burchell* 1396 (Holotype: K; isotype: K).

Miconia chartacea var. *miquelianae* Cogn. in Martius, Fl. bras. 14 (4): 369 (1887).

Miconia chartacea var. *angustifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 839 (1891).

Miconia chartacea var. *brevifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 839 (1891).

Miconia chartacea var. *longifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 838 (1891).

Section: Glossocentrum.

Distribution: Brazil Central-West, Brazil Northeast, Brazil Southeast, Brazil South.

Miconia chinantlana (Naudin) Almeda, Novon 15: 503 (2005).

Basionym: *Staphidium chinantlanum* Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 318 (1851). *Clidemia chinantlana* (Naudin) Triana, Trans. Linn. Soc. London 28: 135 (1871). Type:—MEXICO. *Galeotti* 2945 (Holotype: P; isotype: BR).

Miconia echinoidea Standl. & Steyermark, Publ. Field Mus. Nat. Hist., Bot. Ser. 23 (4): 180 (1944).

Distribution: Mexico, Guatemala.

Section: Octomeris.

Miconia chionophila Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 236 (1850). *Acinodendron chionophilum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *Goudot* s.n. (Holotype: P).

Section: Chaenopleura.

Distribution: Costa Rica, Panama, Colombia, Venezuela, Ecuador, Peru, Bolivia.

Miconia chiriquiensis Almeda, Proc. Calif. Acad. Sci., Ser. 4 42: 303 (1981). Type:—PANAMA. *Almeda* 3535 (Holotype: CAS; isotypes: C, DUKE, F, MO, PMA, US).

Section: Cremanium.

Distribution: Costa Rica, Panama.

Miconia chlorocarpa Cogn., Bull. Acad. Belg., Ser. 3 14: 950 (1887). *Acinodendron chlorocarpum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—COLOMBIA. *André* 3106 (Holotype: K; isotype: BR).

Section: Glossocentrum.

Distribution: Colombia, Ecuador.

Miconia choriophylla Wurdack, Phytologia 23: 499 (1972). Type:—PERU. *Dudley* 10405 (Holotype: US; isotype: NA).

Miconia choriophylla var. *brevifolia* Wurdack, Phytologia 23: 500 (1972).

Section: Cremanium.

Distribution: Peru.

Miconia chrysocoma Gleason, Bull. Torrey Bot. Club 68: 251 (1941). Type:—COLOMBIA. Killip 18035 (Holotype: NY; isotypes: F, US).

Section: Cremanium.

Distribution: Colombia.

Miconia chrysoneura Triana, Trans. Linn. Soc. London 28: 111 (1871). *Acinodendron chrysoneurum* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—MEXICO. Hahn s.n (Holotype: P; isotype: K).

Miconia chrysoneura var. *angustifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 817 (1891).

Section: Miconia.

Distribution: Mexico.

Miconia chrysophylla (Rich.) Urb., Symb. Antill. 4: 459 (1910).

Basionym: *Melastoma chrysophyllum* Rich., Actes Soc. Hist. Nat. Paris 1: 109 (1792). *Acinodendron chrysophyllum* (Rich.) Kuntze, Rev. Gen. Pl. 2: 950 (1891). Type:—FRENCH GUIANA. Leblond s.n. (Holotype: P; isotype: G-DEL).

Melastoma fulvum Rich. in Bonpland, Monogr. Melast. 1: 23 (1807). *Miconia fulva* (Rich.) DC., Prodr. 3: 180 (1828). *Chaenopleura fulva* (Rich.) Griseb., Abh. Königl. Ges. Wiss. Göttingen 9: 55 (1860).

Miconia fulva var. *tinctoria* DC., Prodr. 3: 180 (1828).

Miconia longifolia var. *aubletiana* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 157 (1850). *Miconia fulva* var. *aubletiana* (Naudin) Cogn. in Martius, Fl. bras. 14 (4): 389 (1887).

Miconia fulva var. *angustifolia* Cogn. in Martius, Fl. bras. 14 (4): 389 (1887).

Miconia fulva var. *poeppigii* Cogn. in Martius, Fl. bras. 14 (4): 389 (1887).

Section: Chaenanthera.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Jamaica, Colombia, Venezuela, Trinidad & Tobago, Suriname, Guyana, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West.

Miconia ciliaris Triana, Trans. Linn. Soc. London 28: 105 (1871). *Acinodendron ciliare* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. Ruiz & Pavón 14/65 (Holotype: MA?; isotype: BM).

Section: Tamonea.

Distribution: Peru, Bolivia.

Miconia ciliata (Rich.) DC., Prodr. 3: 179 (1828).

Basionym: *Melastoma ciliatum* Rich., Actes Soc. Hist. Nat. Paris 1: 109 (1792). *Acinodendron ciliatum* (Rich.) Kuntze, Revis. Gen. Pl. 2: 949 (1891). Type:—FRENCH GUIANA. Leblond s.n. in Herb. Richard (Holotype: P).

Melastoma decussatum Vahl, Eclog. Amer. 1: 42 (1797). *Miconia decussata* (Vahl) D. Don ex Loud., Hort. Brit. 1: 174 (1830). *Acinodendron decussatum* (Vahl) Kuntze, Revis. Gen. Pl. 2: 949 (1891).

Miconia barbigera DC., Prodr. 3: 179 (1828). *Hartigia barbigera* (DC.) Miq., Linnaea 18: 623 (1844). *Miconia racemosa* var. *barbigera* (DC.) Cogn. in Martius, Fl. bras. 14 (4): 411 (1888).

Miconia congestiflora DC., Prodr. 3: 180 (1828). *Miconia ciliata* var. *congestiflora* (DC.) Cogn. in Martius, Fl. bras. 14 (4): 409 (1888).

Miconia pileata DC., Prodr. 3: 180 (1828). *Melastoma pileatum* Schrank & Mart. ex DC., Prodr. 3: 180 (1828), pro syn. *Miconia racemosa* var. *pileata* (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 154 (1850). *Acinodendron pileatum* (DC.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Melastoma sessiliflorum Mart. ex DC., Prodr. 3: 180 (1828), pro syn.

Miconia plicata Steud., Nomencl. Bot., ed. 2 (2): 115 (1841).

Clidemia glabriflora K. Presl, Abh. Königl. Böhm. Ges. Wiss., Ser. 5 3: 500 (1845).

Miconia calophylla Crueg., Linnaea 20: 109 (1847). *Miconia cruegeri* Triana, Trans. Linn. Soc. London 28: 131 (1871).

Miconia racemosa var. *latifolia* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 154 (1850).

Miconia racemosa var. *plurilocularis* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 154 (1850). *Miconia ciliata* var. *plurilocularis* (Naudin) Cogn. in Martius, Fl. bras. 14 (4): 410 (1888).

Miconia pileata var. *latifolia* Cogn. in Martius, Fl. bras. 14 (4): 407 (1888).

Miconia pileata var. *longisetosa* Cogn. in Martius, Fl. bras. 14 (4): 407 (1888).

Miconia ciliata var. *attenuata* Cogn. in Martius, Fl. bras. 14 (4): 409 (1888).

Miconia ciliata var. *serrata* Cogn. in Martius, Fl. bras. 14 (4): 410 (1888).

Miconia racemosa var. *urbaniana* Cogn. in Urban, Symb. Antill. 5: 448 (1908).

Miconia virgulata Gleason, Bull. Torrey Bot. Club 52: 386 (1925).

Section: Hartigia.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Jamaica, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Central-West, Brazil Northeast, Brazil Southeast.

Miconia cinchonifolia DC., Prodr. 3: 184 (1828). *Melastoma cinchonifolium* Mart. ex DC., Prodr. 3: 184 (1828), *pro syn.* *Acinodendron cinchonifolium* (DC.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M; isotype: G-DC).

Melastoma leucandrum Schrank ex DC., Prod. 3: 184 (1828), *pro syn.*

Section: Tamonea.

Distribution: Venezuela, Brazil North.

Miconia cinerascens Miq., Linnaea 22: 543 (1849). *Acinodendron cinerascens* (Miq.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Regnell II.109** (Holotype: S).

Miconia proteoides A.St.-Hil. & Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 141 (1850).

Miconia paulensis Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 187 (1850). *Acinodendron paulense* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Melastoma granulosum Willd. ex Triana, Trans. Linn. Soc. London 28: 118 (1871), *pro syn.*

Miconia cinerascens var. *robusta* Wurdack, Sellowia 14: 153 (1962).

Section: Glossocentrum.

Distribution: Brazil Southeast, Brazil South, Paraguay Argentina.

Miconia cinerea Cogn. in Martius, Fl. bras. 14 (4): 290 pl. 60 (1887). *Acinodendron cinereum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—BRAZIL. *Glaziou 11974* (Syntype: BR; isotypes: BR, K, P, R).

Section: Miconia.

Distribution: Brazil Southeast.

Miconia cinnamomifolia (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 68 (1850).

Basionym: *Cremanium cinnamomifolium* DC., Prodr. 3: 194 (1828). Type:—BRAZIL. *Martius s.n.* (Holotype: G-DC).

Miconia martiusiana DC., Prodr. 3: 186 (1828). *Melastoma martianum* Schrank ex DC., Prod. 3: 186 (1828), *pro syn.* *Acinodendron martianum* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Miconia candolleana Triana, Trans. Linn. Soc. London 28: 117 (1871), *nom. illeg.* *Acinodendron candolleanum* (Triana) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Miconia candolleana var. *angustifolia* Cogn. in Martius, Fl. bras. 14 (4): 383 (1887).

Section: Glossocentrum.

Distribution: Brazil Northeast, Brazil Southeast, Brazil South.

Miconia cionotricha L.Uribe, Caldasia 9 (41): 88 (1964). Type:—COLOMBIA. *Hugh-Jones 316* (Holotype: COL; isotypes: K, US).

Section: Miconia.

Distribution: Panama, Colombia, Venezuela.

Miconia cipoensis R.Goldenb., Bol. Bot. Univ. Sao Paulo 18: 29 (1999). Type:—BRAZIL. *Pirani 12699* (Holotype: UEC; isotypes: SPF, US).

Section: Miconia.

Distribution: Brazil Southeast.

Miconia cladonia Gleason, Bull. Torrey Bot. Club 52: 453 (1925). Type:—COLOMBIA. *Pennell 7544* (Holotype: NY; isotypes: GH, PH, US).

Miconia rудis Cogn. & Gleason, Brittonia 1: 123 (1932).

Section: Cremanium.

Distribution: Colombia, Ecuador.

Miconia clathrantha Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 918 (1891). *Acinodendron clathranthum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. Poeppig 1719 (Syntypes: BR, F, NY, US).

Cremanium clathranthum Mart. ex Triana, Trans. Linn. Soc. London 28: 129 (1871), *nom. inval.*

Section: Cremanium.

Distribution: Colombia, Ecuador, Peru.

Miconia cleefii L.Uribe, Caldasia 11 (9): 9 (1975). Type:—COLOMBIA. Cleef 1204 (Holotype: COL; isotypes: U, US).

Section: Cremanium.

Distribution: Colombia.

Miconia clivorum Wurdack, Phytologia 11: 388 (1965). Type:—PERU. Wurdack 1794 (Holotype: US; isotypes: F, GH, K, MICH, UC, USM).

Section: Amblyarrhena.

Distribution: Peru.

Miconia clypeata Wurdack, Phytologia 50: 302 (1982). Type:—COLOMBIA. Luteyn 7325 (Holotype: COL; isotypes: CAS, NY, US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia cocoensis Almeda & Kriebel, Harvard Pap. Bot. 17: 53 (2012). Type:—COSTA RICA. Trusty 270 (Holotype: INB; isotypes: CAS, FTG).

Section: Cremanium.

Distribution: Costa Rica.

Miconia codonostigma Gleason & Wurdack, Phytologia 38: 289 (1978). Type:—COLOMBIA. Cuatrecasas 22468 (Holotype: NY; isotypes: COL, F, MO, U, US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia coelestis (D.Don) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).

Basionym: *Cremanium coeleste* D.Don, Mem. Wern. Nat. Hist. Soc. 4: 314 (1823). *Melastoma coelestis* Pav. ex D.Don, Mem. Wern. Nat. Hist. Soc. 4: 314 (1823), *pro syn.* *Acinodendron coeleste* (D.Don) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. Ruiz & Pavón s.n. (Holotype: OXF?; isotypes: B, destroyed, FI, K).

Miconia chrysopetala Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 233 (1850).

Section: Cremanium.

Distribution: Peru, Bolivia.

Miconia collatata Wurdack, Phytologia 29: 145 (1974). Type:—BRAZIL. Irwin 17991 (Holotype: US; isotypes: F, GH, MO, NY, S, SP, TEX).

Section: Glossocentrum.

Distribution: Bolivia, Brazil North, Brazil Central-West, Brazil Southeast, Brazil South, Paraguay, Argentina.

Miconia collayensis Wurdack, Mem. New York Bot. Gard. 16: 13 (1967). Type:—ECUADOR. Camp E-4890 (Holotype: US; isotype: NY).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia colliculosa Almeda, Proc. Calif. Acad. Sci., Ser. 4 52: 33 (2000). Type:—COSTA RICA. Herrera 3129 (Holotype: CAS; isotypes: CAS, CR, INB, MEXU, MO).

Section: Amblyarrhena.

Distribution: Costa Rica, Panama.

Miconia coloradensis Almeda, Proc. Calif. Acad. Sci., Ser. 4 42: 305 (1981). Type:—PANAMA. *Folsom* 6143 (Holotype: CAS; isotypes: MO, PMA).

Section: Amblyarrhena.

Distribution: Panama.

Miconia commutata Almeda, Novon 15: 504 (2005). Type:—COSTA RICA. *Hammel* 11237 (Holotype: CAS; isotype: DUKE).

Section: Miconia.

Distribution: Costa Rica, Panama.

Miconia comosa Cogn. in Martius, Fl. bras. 14 (4): 408 (1888). *Acinodendron comosum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Spruce* 4438 (Syntypes: BR, K; isosyntypes: BM, NY, TCD).

Section: Cremanium.

Distribution: Peru.

Miconia compressa Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 128 (1850). *Acinodendron compressum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Herb. Martius* s.n. (Holotype: P).

Section: Miconia.

Distribution: Brazil Northeast.

Miconia compressicaulis Wurdack, Brittonia 9: 104 (1957). Type:—COLOMBIA. *Core* 1284 (Holotype: US; isotype: NY).

Section: Cremanium.

Distribution: Colombia.

Miconia comptifolia Wurdack, Phytologia 55: 138 (1984). Type:—BRAZIL. *Boyan* 291 (Holotype: INPA).

Section: Jucunda.

Distribution: Brazil North.

Miconia concinna Almeda, Proc. Calif. Acad. Sci., Ser. 4 42: 307 (1981). *Topobea micrantha* Pittier, J. Wash. Acad. Sci. 14 (19): 451 (1924). Type:—PANAMA. *Pittier* 3276 (Holotype: US).

Section: Chaenopleura.

Distribution: Panama.

Miconia condylata Wurdack, Phytologia 7: 236 (1960). Type:—PERU. *Ferreyra* 1105 (Holotype: US; isotype: NY).

Section: Amblyarrhena.

Distribution: Peru.

Miconia confertiflora Almeda, Proc. Calif. Acad. Sci., Ser. 4 42: 309 (1981). Type:—COSTA RICA. *Almeda* 2908 (Holotype: CAS; isotypes: CR, F, MO, US).

Section: Chaenopleura.

Distribution: Costa Rica.

Miconia conformis Wurdack, Phytologia 52: 64 (1982). Type:—ECUADOR. *Jaramillo* 3527 (Holotype: AAU; isotype: QCA).

Section: Miconia.

Distribution: Ecuador.

Miconia coniophora Urb. & Ekman, Ark. Bot. 22A (17): 42 (1929). Type:—HAITI. *Ekman* H8067 (Holotype: S; isotypes: A, C, NY, S, US).

Section: Chaenopleura.

Distribution: Haiti, Dominican Republic.

Miconia contrerasii Wurdack, Wrightia 5: 226 (1976). Type:—GUATEMALA. Lundell 19406 (Holotype: US; isotypes: F, LL, S).
Section: Octomeris.
Distribution: Guatemala.

Miconia cookii Gleason, Phytologia 1: 43 (1933). Type:—PERU. Cook 1742 (Holotype: US).
Section: Amblyarrhena.
Distribution: Peru, Bolivia.

Miconia corallina Spring, Flora 20 (2, Beibl.): 77 (1837). *Acinodendron corallinum* (Spring) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. Martius s.n. (Holotype: M).
Miconia soltmanniana Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 132 (1850). *Soltmannia splendens* Klotzsch ex Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 132 (1850), *pro syn.*
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia corazonica Wurdack, Mem. New York Bot. Gard. 16: 35 (1967). Type:—ECUADOR. Camp E-1664 (Holotype: NY; isotype: US)
Section: Chaenopleura.
Distribution: Ecuador.

Miconia cordata Triana, Trans. Linn. Soc. London 28: 114 (1871). *Acinodendron cordatum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU or BOLIVIA. Pearce s.n. (Holotype: K).
Section: Miconia.
Distribution: Peru, Bolivia.

Miconia cordifolia Wurdack, Phytologia 31: 498 (1975). Type:—COLOMBIA. Uribe-Uribe 3873 (Holotype: US; isotype: COL).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia coriacea (Sw.) DC., Prodr. 3: 189 (1828).
Basionym: *Melastoma coriaceum* Sw., Prodr. 1: 70 (1788). *Acinodendron coriaceum* (Sw.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—GUADELOUPE. Swartz s.n. (Holotype: BM; isotype: S).
Melastoma cymbifolium Rich., Bonpl. Melast. 57 (1809).
Section: Cremanium.
Distribution: Costa Rica, Guadeloupe, Dominica.

Miconia cornifolia (Desr.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 126 (1850).
Basionym: *Melastoma cornifolium* Desr. in Lamarck, Encycl. Méth. Bot. 4: 51 (1797). *Conostegia cornifolia* (Desr.) Ser. ex DC., Prodr. 3: 175 (1828). Type:—MARTINIQUE. S.col., s.n. (Holotype: P).
Melastoma cinnamomifolium Jacq., Collectanea 2: 111 (1788). *Miconia cinnamomifolia* (Jacq.) Triana, Trans. Linn. Soc. London 28: 101 (1871), *nom. illeg.* *Acinodendron cinnamomifolium* (Jacq.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).
Melastoma pictum Vahl, Eclog. Amer. 3: 15 (1807). *Miconia picta* (Vahl) Wurdack, Phytologia 7: 235 (1960).
Melastoma lutescens Vahl, Eclog. Amer. 3: 17 (1807). *Conostegia lutescens* (Vahl.) Ser. ex DC., Prodr. 3: 175 (1828).
Miconia candolleana Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).
Melastoma semicrenatum Rich. in Bonpland, Monogr. Melast. 1: 69 (1811). *Conostegia semicrenata* (Rich.) Ser. ex DC., Prodr. 3: 175 (1828). *Miconia semicrenata* (Rich.) D.Don in Loudon, Hort. Brit. 1: 174 (1830). *Tetrazygia semicrenata* (Rich.) Griseb., Fl. Brit. W. Indian Is. 255 (1860).
Miconia striata var. *vincentina* Domin, Preslia 9: 4 (1930).
Miconia elongata Stehle & Quentin, Fl. Guadeloupe Depend. 2 (3): 91 (1949).
Section: Laceraria.
Distribution: Guadeloupe, Dominica, Martinique, Santa Lucia, Barbados, Grenada.

Miconia coronata (Bonpl.) DC., Prodr. 3: 187 (1828).

Basionym: *Melastoma coronatum* Bonpl., Monogr. Melast. 1: 130 (1816). *Acinodendron coronatum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Linden* 817 (Holotype: G).

Miconia strangulata Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 117 (1850).

Section: Tamonea.

Distribution: Colombia.

Miconia coronifera Wurdack, Brittonia 9: 104 (1957). Type:—COLOMBIA. *Little* 8518 (Holotype: NY; isotypes: B, COL, US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia correae Almeda, Proc. Calif. Acad. Sci., Ser. 4 52: 37 (2000). Type:—PANAMA. *Dressler* 5806 (Holotype: CAS; isotype: PMA).

Section: Jucunda.

Distribution: Costa Rica, Panama.

Miconia corymbiformis Cogn., Bull. Acad. Belg., Ser. 3 14: 964 (1887). *Acinodendron corymbiforme* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *André* 3061 (Syntypes: BR, K, NY).

Section: Chaenopleura.

Distribution: Colombia, Ecuador.

Miconia cosangensis Wurdack, Phytologia 39: 323 (1978). Type:—ECUADOR. *Asplund* 9564 (Holotype: S).

Section: Cremanium.

Distribution: Ecuador.

Miconia costaricensis Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 887 (1891). *Acinodendron costaricense* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COSTA RICA. *Wendland* 936 (Holotype: GOET; isotype: BR).

Miconia costaricensis var. *pittieri* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 888 (1891).

Section: Amblyarrhena.

Distribution: Mexico, Costa Rica, Panama, Colombia.

Miconia cowanii Wurdack, Mem. New York Bot. Gard. 10 (4): 37 (1961). Type:—BRAZIL. *Cowan* 38399 (Holotype: NY; isotypes: RB, US).

Section: Miconia.

Distribution: Brazil North.

Miconia crassifolia Triana, Trans. Linn. Soc. London 28: 125 (1871). *Acinodendron crassifolium* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Mathews* 1261 (Syntypes: BR, K, NY).

Miconia crassifolia var. *parvifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 882 (1891).

Miconia crassifolia var. *rotundifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 882 (1891).

Section: Amblyarrhena.

Distribution: Peru.

Miconia crassinervia Cogn. in Martius, Fl. bras. 14 (4): 391 (1887). *Acinodendron crassinerve* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Riedel* 1396 (Holotype: LE; isotypes: BR, K, P).

Section: Miconia.

Distribution: Venezuela, Brazil North.

Miconia crassipes Triana, Trans. Linn. Soc. London 28: 116 (1871). *Acinodendron crassipes* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Mathews* 1300 (Holotype: K; isotypes: BR, K).

Section: Miconia.

Distribution: Peru.

Miconia crassistigma Cogn., Bot. Jahrb. Syst. 42: 143 (1909). Type:—PERU. *Weberbauer* 3407 (Holotype: B, destroyed; isotypes: BR, MOL).
Section: Cremanium.
Distribution: Peru.

Miconia creribullata Wurdack, Phytologia 64: 298 (1988). Type:—ECUADOR. *Dobson* 1080 (Holotype: US; isotypes: CONN, S).
Section: Cremanium.
Distribution: Ecuador.

Miconia cremadena Gleason, Bull. Torrey Bot. Club 66: 417 (1939). Type:—COSTA RICA. *Skutch* 3394 (Holotype: NY; isotypes: K, MO, S, US).
Section: Cremanium.
Distribution: Costa Rica.

Miconia cremophylla Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 228 (1850). *Acinodendron cremophyllum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Mathews* 1297 (Holotype: P; isotypes: G, K, NY, P).
Section: Cremanium.
Distribution: Peru, Bolivia.

Miconia cretacea Gleason, Amer. J. Bot. 19: 744 (1932). Type:—BOLIVIA. *Buchtien* 1025 (Holotype: US).
Section: Miconia.
Distribution: Peru, Bolivia, Brazil North.

Miconia crinita Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 194 (1850). *Acinodendron crinitum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Linden* 1067 (Holotype: P; isotypes: BR, P).

Miconia cardiophylla Cogn., Bull. Acad. Belg., Ser. 3 14: 960 (1887). *Acinodendron cardiophyllum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Miconia crinita ssp. *australis* Wurdack, Phytologia 31: 500 (1975).
Section: Amblyarrhena.
Distribution: Colombia, Ecuador.

Miconia crocata Almeda, Proc. Calif. Acad. Sci., Ser. 4 52: 39 (2000). Type:—PANAMA. *Almeda* 5930 (Holotype: CAS; isotypes: DUKE, MEXU, MO, NY, PMA, TEX, US).
Section: Cremanium.
Distribution: Panama.

Miconia crocea (Desr.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).
Basionym: *Melastoma croceum* Desr. in Lamarck, Encycl. Méth. Bot. 4: 55 (1797). *Cremanium croceum* (Desr.) DC., Prodr. 3: 194 (1828). *Acinodendron croceum* (Desr.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU or ECUADOR. *de Jussieu* s.n. (Holotype: P-JU).

Melastoma setinode Bonpl., Monogr. Melast. 1: 3 (1807). *Miconia setinodis* (Bonpl.) DC., Prodr. 3: 179 (1828).
Acinodendron setinode (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Miconia theresiae Cogn. in Bayern, Beih. Bot. Centralbl. 13: 80 (1902).

Miconia crocea var. *setosa* Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 12: 181 (1934).

Section: Cremanium.

Distribution: Colombia, Ecuador.

Miconia cruenta Triana, Trans. Linn. Soc. London 28: 122 (1871). *Acinodendron cruentum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Triana* s.n. (Holotype: BM; isotypes: COL, K, US).

Miconia neurocarpa Triana, Trans. Linn. Soc. London 28: 108 (1871). *Acinodendron neurocarpum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia cuatrecasae Markgr. in Cuatrecasas, Trab. Mus. Nac. Ci. Nat., Ser. Bot. 26: 27 (1933). Type:—COLOMBIA. Cuatrecasas 2509 (Syntype: MA; isosyntype: US).
Section: Cremanium.
Distribution: Colombia.

Miconia cubatanensis Hoehne, Anexos Mem. Inst. Butantan, Secc. Bot. 1 (5): 139 (1922). Type:—BRAZIL. Hoehne 3341 (Holotype: SP; isotypes: BM, US).

Miconia revoluta Miq., Linnaea 22: 544 (1849), *nom. illeg.* *Acinodendron revolutum* (Miq.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Miconia revoluta var. *nitida* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 1190 (1891).
Section: Glossocentrum.
Distribution: Brazil Central-West, Brazil Northeast, Brazil Southeast, Brazil South.

Miconia cubensis (C.Wright ex Griseb.) C.Wright in Sauvalle, Anales Acad. Ci. Med. Habana 5: 465 (1869).

Basionym: *Pleurochaenia cubensis* C.Wright ex Griseb., Cat. Pl. Cub. 101 (1866). *Acinodendron cubense* (C.Wright ex Griseb.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). *Tamonea cubensis* (C.Wright ex Griseb.) Krasser in Engler & Prantl, Nat. Pflanzenfam. 3 (7): 188 (1893). Type:—CUBA. Wright 2514 (Holotype: GOET; isotypes: BM, GH, GOET, MO, NY).
Section: Chaenopleura.
Distribution: Cuba.

Miconia cundinamarcensis Wurdack, Caldasia 7 (35): 332 (1957). Type:—COLOMBIA. Uribe-Urabe 2677 (Holotype: NY; isotypes: COL, F, U).

Section: Cremanium.
Distribution: Colombia.

Miconia cuneata Triana, Trans. Linn. Soc. London 28: 113 (1871). *Staphidium cuneatum* Berg ex Triana, Trans. Linn. Soc. London 28: 113 (1871), *pro syn.* *Acinodendron cuneatum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. Poeppig 2047 (Syntypes: B, destroyed, BR, NY).

Section: Laceraria.
Distribution: Ecuador, Peru, Bolivia.

Miconia cuprea Wurdack, Mem. New York Bot. Gard. 16: 27 (1967). Type:—ECUADOR. Camp E-4736 (Holotype: US; isotypes: F, GH, NY).

Section: Cremanium.
Distribution: Ecuador.

Miconia curta (Gleason) Wurdack, Mem. New York Bot. Gard. 10 (5): 170 (1964).

Basionym: *Graffenrieda curta* Gleason in Steyermark, Fieldiana, Bot. 28: 431 (1952). Type:—VENEZUELA. Steyermark 58026a (Holotype: F; isotype: VEN).

Miconia curta ssp. *ptariensis* Wurdack, Mem. New York Bot. Gard. 10 (5): 170 (1964).

Section: Miconia.
Distribution: Venezuela.

Miconia cuspidata Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 163 163 (1850). Type:—BRAZIL. Poeppig 3037 (Holotype: P; isotypes: BR, P).

Section: Miconia.
Distribution: Peru, Brazil North, Brazil Central-West, Brazil Northeast.

Miconia cuspidatissima Pittier, J. Wash. Acad. Sci. 14 (19): 449 (1924). Type:—PANAMA. Pittier 3224 (Holotype: US; isotype: F).

Section: Cremanium.
Distribution: Costa Rica, Panama.

Miconia cutucuensis Wurdack, Mem. New York Bot. Gard. 16: 9 (1967). Type:—ECUADOR. *Camp E-1297* (Holotype: NY; isotype: US).
Section: Miconia.
Distribution: Ecuador.

Miconia cyanocarpa Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 221 (1850). *Acinodendron cyanocarpum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BOLIVIA. *Weddell 4205* (Holotype: P).
Miconia cyanocarpa var. *parvifolia* Cogn., Bull. Torrey Bot. Club 23: 17 (1896).
Miconia cyanocarpa var. *hirsuta* Cogn., Bull. Torrey Bot. Club 23: 277 (1896). *Acinodendron cyanocarpum* var. *hirsutum* (Cogn.) Kuntze, Revis. Gen. Pl. 3: 93 (1898).
Section: Cremanium.
Distribution: Peru, Bolivia.

Miconia cyathanthera Triana, Trans. Linn. Soc. London 28: 127 (1871). *Cyathanthera lanceolata* Pohl, Pl. Bras. Icon. Descr. 2: 131 (1831). *Acinodendron cyathantherum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Pohl s.n.* (Holotype: W; isotypes: BM, M).
Section: Cremanium.
Distribution: Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia danielii Almeda, Novon 15: 506 (2005). Type:—COSTA RICA. *Almeda 2876* (Holotype: CAS; isotype: CR).
Section: Cremanium.
Distribution: Costa Rica.

Miconia dapsiliflora Wurdack, Phytologia 26: 7 (1973). Type:—ECUADOR. *Asplund 17463* (Holotype: S).
Section: Amblyarrhena.
Distribution: Colombia, Ecuador.

Miconia dasyclada Wurdack, Phytologia 23: 492 (1972). Type:—PERU. *Dudley 11829* (Holotype: US; isotype: NA).
Section: Cremanium.
Distribution: Peru.

Miconia decipiens Cogn., Bull. Acad. Belg., Ser. 3 14: 948 (1887). *Acinodendron decipiens* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *André 3455* (Holotype: K; isotypes: BR, K, NY).
Miconia nigricalyx Gleason, Bull. Torrey Bot. Club 57: 67 (1930).
Section: Miconia.
Distribution: Colombia.

Miconia decurrens Cogn., Bol. Mus. Paraense Hist. Nat. Ethnogr. 5: 253 (1909). Type:—PERU. *Ducke 7603* (Holotype: BR; isotypes: MG, RB, US).
Section: Jucunda.
Distribution: Mexico, Nicaragua, Colombia, Venezuela, Ecuador, Peru, Bolivia, Brazil North.

Miconia delicatula A.Rich. in Sagra, Hist. Fis. Cuba, Bot. 10: 268 (1845). *Acinodendron delicatulum* (A.Rich.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). *Tamonea delicatula* (A.Rich.) Jenn., Ann. Carnegie Mus. 11 (1-2): 210 (1917). *Tetrazygia delicatula* (A.Rich.) Borhidi, Acta Bot. Hung. 23: 37 (1977). Type:—CUBA. *La Sagra s.n.* (Holotype: P?).

Miconia chlorophylla C.Wright in Sauvalle, Anales Acad. Ci. Med. Habana 5: 464 (1869).
Section: Octomeris.
Distribution: Cuba.

Miconia demissifolia Wurdack, Phytologia 23: 489 (1972). Type:—PERU. *Madison 10168* (Holotype: US; isotypes: MO, NA).
Section: Cremanium.
Distribution: Peru.

Miconia densifolia Cogn., Bot. Jahrb. Syst. 42: 145 (1909). Type:—PERU. *Weberbauer 3404* (Holotype: B, destroyed; isotype: BR).
Section: Cremanium.
Distribution: Peru.

Miconia denticulata Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 214 (1850). *Acinodendron denticulatum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Bonpland 3451* (Holotype: P; isotype: P).

Miconia centrophora var. *subintegerrima* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 883 (1891).

Miconia chrysanthera Cogn., Bot. Jahrb. Syst. 42: 141 (1909).

Section: Amblyarrhena.

Distribution: Colombia, Ecuador, Peru.

Miconia desmantha Benth., Pl. Hartw. 181 (1845). *Acinodendron desmanthum* (Benth.) Kuntze, Revis. Gen. Pl. 1: 243 (1891). Type:—COLOMBIA. *Hartweg 1010* (Holotype: K; isotypes: BM, F, LD, US).

Miconia bourgeana Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 772 (1891). *Acinodendron bourgeanum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Miconia carioana Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 773 (1891). *Acinodendron carioanum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia.

Miconia despentesii Urb., Symb. Antill. 8: 496 (1921). Type:—DOMINICAN REPUBLIC. *Fuertes 1491* (Lectotype: NY).

Section: Cremanium.

Distribution: Dominican Republic.

Miconia diaphanea Gleason, Recueil Trav. Bot. Neerl. 32: 213 (1935). Type:—SURINAME. *Gonggrijp 186* (Holotype: U).

Section: Laceraria.

Distribution: Colombia, Guyana, Suriname, French Guiana, Peru, Brazil North.

Miconia dichroa Cogn. in de Candolle & de Candolle, Monogr. Phan. 7 1192 (1891). *Acinodendron dichroum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Glaziou s.n.* (Holotype: BR).

Section: Cremanium.

Distribution: Brazil Southeast.

Miconia dichrophylla J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 189 (1929). Type:—PERU. *Macbride 5495* (Holotype: F; isotypes: G, K, US).

Miconia elaeagnoides Cogn. in Martius, Fl. bras. 14 (4): 390 (1887), nom. illeg. *Acinodendron elaeagnoides* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Section: Miconia.

Distribution: Peru, Bolivia, Brazil North, Brazil West-Central.

Miconia diegogomezii Kriebel & Almeda, Harvard Pap. Bot. 17: 55 (2012). Type:—COSTA RICA. *Lépiz 347* (Holotype: INB; isotypes: CAS, CR, MO).

Section: Cremanium.

Distribution: Costa Rica.

Miconia dielsiana Urb., Ark. Bot. 23A (11): 22 (1931). Type:—DOMINICAN REPUBLIC. *Ekman H13229* (Holotype: S; isotype: NY).

Section: Chaenopleura.

Distribution: Dominican Republic.

Miconia dielsii Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 12: 180 (1934). Type:—ECUADOR. *Diels* 952 (Holotype: B, destroyed).
Section: Cremanium.
Distribution: Ecuador, Peru.

Miconia difficilis Triana, Trans. Linn. Soc. London 28: 122 (1871). *Acinodendron difficile* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—ECUADOR. *Spruce* 4972 (Syntypes: BM, CAS, K, NY, TCD).
Section: Amblyarrhena.
Distribution: Colombia, Ecuador.

Miconia dioica Wurdack, Mem. New York Bot. Gard. 10 (5): 173 (1964). Type:—VENEZUELA. *Steyermark* 1226 (Holotype: US; isotypes: CAS, F, GH, NY, U, VEN).
Section: Cremanium.
Distribution: Venezuela.

Miconia dipsacea Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 138 (1850). *Melastoma amplexicaule* Pav. ex Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 138 (1850), *pro syn.* *Acinodendron dipsaceum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Poeppig s.n.* (Holotype: P).
Section: Miconia.
Distribution: Peru.

Miconia discolor DC., Prodr. 3: 184 (1828). *Acinodendron discolor* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Langsdorff s.n.* (Holotype: B, destroyed; isotype: P).
Miconia discolor var. *subconcolor* Cogn. in Martius, Fl. bras. 14 (4): 284 (1887).
Miconia kleinii Brade, Sellowia 12: 144 (1960).
Section: Glossocentrum.
Distribution: Brazil Southeast, Brazil South, Paraguay, Argentina.

Miconia dispar Benth., Hooker's J. Bot. Kew Gard. Misc. 2: 241 (1850). *Acinodendron dispar* (Benth.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Spruce s.n.* (Holotype: K).
Section: Miconia.
Distribution: Venezuela, French Guiana, Peru, Bolivia, Brazil North, Brazil Central-West.

Miconia dissimulans Wurdack, Mem. New York Bot. Gard. 16: 34 (1967). Type:—ECUADOR. *Espinosa* 1545 (Holotype: US; isotype: NY).
Section: Cremanium.
Distribution: Ecuador.

Miconia dissita Almeda, Novon 15: 508 (2005). Type:—PANAMA. *Almeda* 6246 (Holotype: CAS; isotypes: MO, PMA).
Section: Cremanium.
Distribution: Costa Rica, Panama.

Miconia dissitiflora Almeda, Proc. Calif. Acad. Sci., Ser. 4 46: 146 (1989). Type:—COSTA RICA. *Almeda* 3093 (Holotype: CAS; isotypes: BM, BR, CR, DUKE, F, MO, NY, US).
Section: Amblyarrhena.
Distribution: Costa Rica.

Miconia dissitinervia Kriebel, Almeda & Estrada, Proc. Calif. Acad. Sci., Ser. 4 56 (37): 678 (2005). Type:—COSTA RICA. *Estrada* 3101 (Holotype: CR; isotypes: CAS, INB, MO).
Section: Amblyarrhena.
Distribution: Costa Rica, Panama.

Miconia divaricata Gardner, London J. Bot. 2: 345 (1843). *Acinodendron divaricatum* (Gardner) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Gardner* 391 (Holotype: BM; isotypes: CAS, F, G, K, NY, P, S, US).
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia divergens Triana, Trans. Linn. Soc. London 28: 127 (1871). *Acinodendron divergens* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Triana* 3999 (Holotype: BM; isotype: COL).
Section: Cremanium.
Distribution: Colombia, Ecuador.

Miconia divisoriana Wurdack, Phytologia 16: 175 (1968). Type:—PERU. *Ferreyra* 1669 (Holotype: US; isotype: USM).
Section: Laceraria.
Distribution: Peru.

Miconia dodecandra (Desr.) Cogn. in Martius, Fl. bras. 14 (4): 243 (1887).
Basionym: *Melastoma dodecandrum* Desr. in Lamarck, Encycl. Méth. Bot. 4: 46 (1797). *Acinodendron dodecandrum* (Desr.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—NOT LOCATED.

Melastoma swartzianum Rich. in Bonpland, Monogr. Melast. 1: 74 (1811). *Diplochita swartziana* (Rich.) DC., Prodr. 3: 177 (1828).

Miconia fothergilla var. *gracilipes* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 120 (1850). *Miconia dodecandra* var. *gracilipes* (Naudin) Cogn. in Martius, Fl. bras. 14 (4): 244 (1887).

Miconia dodecandra var. *longifolia* Cogn. in Martius, Fl. bras. 14 (4): 244 (1887).

Section: Tamonea.

Distribution: Mexico, Belize, Guatemala, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Cuba, Jamaica, Haiti, Dominican Republic, Dominica, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast, Brazil South.

Miconia dodsonii Wurdack, Mem. New York Bot. Gard. 16: 23 (1967). Type:—ECUADOR. *Dodson* 681 (Holotype: US; isotype: MO).

Section: Cremanium.

Distribution: Ecuador.

Miconia dolichopoda Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 217 (1850). *Acinodendron dolichopodium* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—VENEZUELA. *Funck* 93 (Holotype: P; isotypes: BM, BR, G, P).
Section: Cremanium.

Distribution: Costa Rica, Colombia, Venezuela.

Miconia dolichorrhyncha Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 166 (1850). *Acinodendron dolichorrhynchum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Mathews* 1291 (Holotype: P; isotypes: BM, K, NY, P).
Eurychaenia fulva Griseb., Fl. Brit. W. Indian Is. 259 (1860). *Acinodendron fulvum* (Griseb.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Miconia solmsii Cogn. in Martius, Fl. bras. 14 (4): 393 (1887).

Miconia stellipilis Cogn. in Rusby, Bull. New York Bot. Gard. 4: 360 (1907).

Miconia pilgeriana Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 363 (1915).

Miconia buntingii Wurdack, Phytologia 21: 357 (1971).

Section: Chaenanthera.

Distribution: Costa Rica, Panama, Colombia, Venezuela, Trinidad & Tobago, Guyana, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West.

Miconia domingensis Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 931 (1891). *Acinodendron domingense* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—HAITI. de Menonville s.n. (Holotype: MPU - "Herb. Thouin"?; isotype: P?).

Section: Chaenopleura.

Distribution: Haiti, Dominican Republic.

Miconia donaeana Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850). *Acinodendron donaeanum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Pavón s.n.* (Holotype: BM).

Chitonita caudata D.Don, Mem. Wern. Nat. Hist. Soc. 4: 319 (1823). *Diplochita caudata* (D.Don) DC., Prodr. 3: 177 (1828).

Miconia longicaudata Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 738 (1891). *Acinodendron longicaudatum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Miconia leucantha Gleason, Bull. Torrey Bot. Club 63: 534 (1936).

Section: Tamonea.

Distribution: Guatemala, Honduras, Costa Rica, Nicaragua, Panama, Colombia, Venezuela, Peru.

Miconia dorsaliporosa R.Goldenb. & Reginato, J. Torrey Bot. Soc. 136: 294 (2009). Type:—BRAZIL. *Paixão* 851 (Holotype: CEPEC; isotypes: RB, UPCB).

Section: Glossocentrum.

Distribution: Brazil Northeast, Brazil Southeast.

Miconia dorsiloba Gleason, Brittonia 2: 320 (1937). Type:—COSTA RICA. *Skutch* 2816 (Holotype: NY; isotypes: GH, K, MICH, MO, S, US).

Section: Laceraria.

Distribution: Nicaragua, Costa Rica, Panama, Colombia, Ecuador, Peru.

Miconia duckei Cogn., Bol. Mus. Paraense Hist. Nat. Ethnogr. 5: 254 (1909). Type:—PERU. *Ducke* 7586 (Holotype: BR; isotypes: MG, RB, US).

Section: Laceraria.

Distribution: Peru, Bolivia, Brazil North.

Miconia dudleyi Wurdack, Phytologia 21: 356 (1971). Type:—PERU. *Dudley* 11489 (Holotype: US; isotype: NA).

Section: Miconia.

Distribution: Ecuador, Peru.

Miconia dumetosa Cogn., Bot. Jahrb. Syst. 42: 144 (1909). Type:—PERU. *Weberbauer* 4389 (Holotype: B, destroyed; isotypes: BR, MOL).

Section: Cremanium.

Distribution: Peru.

Miconia dunstervillei Wurdack, Acta Bot. Venez. 4: 60 (1970). Type:—VENEZUELA. *Steyermark* 100904 (Holotype: US; isotype: VEN).

Section: Amblyarrhena.

Distribution: Colombia, Venezuela.

Miconia dura Triana, Trans. Linn. Soc. London 28: 131 (1871). *Acinodendron durum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Sellow s.n.* (Holotype: B, destroyed).

Section: Uncertain

Distribution: Brazil Southeast (probably).

Miconia egensis Cogn. in Martius, Fl. bras. 14 (4): 374 (1887). *Acinodendron egense* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Poeppig* 2583 (Holotype: BR).

Miconia panicularis Gleason, Bull. Torrey Bot. Club 52: 384 (1925).

Section: Glossocentrum.

Distribution: Venezuela, Guyana, Suriname, French Guiana, Ecuador, Peru, Brazil North, Brazil Central-West.

Miconia egregia Wurdack, Phytologia 39: 326 (1978). Type:—PERU. *Sandeman* 4086 (Holotype: K).

Section: Cremanium.

Distribution: Peru.

Miconia eichleri Cogn. in Martius, Fl. bras. 14 (4): 362 (1887). *Acinodendron eichleri* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. Glaziou 8679 (Syntype: BR; isosyntypes: C, K, P, US).
Miconia eichleri var. *australis* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 1190 (1891).
Section: Glossocentrum.
Distribution: Brazil Southeast, Brazil South.

Miconia elaeodendron (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850).
Basionym: *Cremanium elaeodendrum* DC., Prodr. 3: 191 (1828). *Acinodendron elaeodendron* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. Herb. Deless. s.n. (Holotype: G-DC; isotypes: BR, G, L).
Miconia verticillata Triana, Trans. Linn. Soc. London 28: 117 (1871). *Acinodendron verticillatum* (Triana) Kuntze, Revis. Gen. Pl. 2: 953 (1891).
Section: Chaenanthera.
Distribution: Brazil Southeast.

Miconia elaeoides Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 223 (1850). *Acinodendron elaeoides* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. Goudot s.n. (Holotype: P).
Miconia pallida Gleason, Bull. Torrey Bot. Club 57: 72 (1930).
Miconia ligustriflora Cogn. ex Cuatrec., Trab. Mus. Cienc. Nat., Ser. Bot. 33: 91 (1936).
Section: Cremanium.
Distribution: Colombia, Venezuela.

Miconia elata (Sw.) DC., Prodr. 3: 182 (1828).
Basionym: *Melastoma elatum* Sw., Prodr. 70 (1788). *Acinodendron elatum* (Sw.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—JAMAICA. Swartz s.n. (Lectotype: BM).
Miconia eurychaenoides Griseb., Pl. Wright. 1: 185 (1860).
Section: Miconia.
Distribution: Mexico, Belize, Honduras, Nicaragua, Costa Rica, Panama, Cuba, Jamaica, Colombia, Venezuela, French Guiana, Ecuador, Peru, Bolivia, Brazil North.

Miconia elegans Cogn. in Martius, Fl. bras. 14 (4): 312 (1887). *Acinodendron elegans* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. Glaziou s.n. (Syntype: BR).
Miconia elegans var. *latialata* Cogn. in Martius, Fl. bras. 14 (4): 313 (1887).
Miconia elegans var. *pauciflora* Cogn. in Martius, Fl. bras. 14 (4): 313 (1887).
Section: Miconia.
Distribution: Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia elongata Cogn. in Britton, Bull. Torrey Bot. Club 17: 91 (1890). *Acinodendron elongatum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BOLIVIA. Rusby 2283 (Holotype: NY; isotypes: BR, CAS, MICH, NY, PH, US).
Section: Cremanium.
Distribution: Peru, Bolivia.

Miconia elvirae Wurdack, Phytologia 69: 321 (1990). Type:—VENEZUELA. Luteyn 9635 (Holotype: VEN; isotypes: CAS, NY, PORT, US, VEN).
Section: Cremanium.
Distribution: Colombia, Venezuela.

Miconia emendata Wurdack, Phytologia 35: 249 (1977). Type:—PERU. Tessmann 5238 (Holotype: S; isotype: NY).
Section: Miconia.
Distribution: Panama, Colombia, Peru, Brazil North.

Miconia eremita L.Uribe, Caldasia 8: 538 (1962). Type:—COLOMBIA. Uribe-Uribe 3625 (Holotype: COL; isotypes: COL, US).
Section: Tamonea.
Distribution: Colombia.

Miconia eriocalyx Cogn. in Martius, Fl. bras. 14 (4): 342 (1887). *Acinodendron eriocalyx* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. Poeppig 2245 (Syntype: BR; isosyntype: NY).
Section: Miconia.
Distribution: Peru, Bolivia, Brazil North.

Miconia erioclada Triana, Trans. Linn. Soc. London 28: 111 (1871). *Acinodendron eriocladum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. Spruce s.n. (Holotype: K).

Miconia acreana Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 360 (1915).
Section: Miconia.
Distribution: Colombia, Ecuador, Peru, Bolivia, Brazil North.

Miconia eriodonta DC., Prodr. 3: 185 (1828). *Acinodendron eriodontum* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).
Type:—FRENCH GUIANA. Poiteau s.n. (Holotype: G-DEL).

Miconia eriodonta var. *oblongifolia* DC., Prodr. 3: 185 (1828).
Miconia bracteolaris Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 150 (1850).
Miconia bracteolaris var. *cardiophora* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 151 (1850).
Section: Miconia.
Distribution: French Guiana, Brazil North.

Miconia ernstii Wurdack, Phytologia 11: 385 (1965). Type:—DOMINICA. Ernst 1723 (Holotype: US; isotypes: B, CAS).

Section: Glossocentrum.
Distribution: Dominica.

Miconia erosa Gleason, Bull. Torrey Bot. Club 57: 70 (1930). Type:—COLOMBIA. Toro 1139 (Holotype: NY).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia espinosae Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 15: 378 (1941). Type:—ECUADOR. Espinosa 758 (Holotype: B, destroyed).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia eugeniooides Triana, Trans. Linn. Soc. London 28: 109 (1871). *Acinodendron eugeniooides* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—VENEZUELA. Spruce 3531 (Holotype, K; isotypes: BM, BR, GH, K, NY, TCD).
Oxymeris cuspidata Triana, Trans. Linn. Soc. London 28: 93 (1871). *Leandra cuspidata* (Triana) Cogn. in Martius, Fl. bras. 14 (4): 175 (1886). *Acinodendron cuspidatum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Section: Miconia.
Distribution: Colombia, Venezuela, Brazil North.

Miconia expansa Gleason, Bull. Torrey Bot. Club 63: 536 (1936). Type:—PERU. Klug 3201 (Holotype: NY, isotypes: BC, F, GH, K, MO, S, US, WIS).
Section: Amblyarrhena.
Distribution: Peru.

Miconia explicita Wurdack, Phytologia 26: 3 (1973). Type:—ECUADOR. Harling 4475 (Holotype: S; isotypes: LL, NY).
Section: Octomeris.
Distribution: Ecuador.

Miconia falcata Cogn., Bot. Jahrb. Syst. 42: 139 (1909). Type:—PERU. Weberbauer 4998 (Holotype: B, destroyed; isotype: BR).
Section: Miconia.
Distribution: Peru.

Miconia fallax DC., Prodr. 3: 181 (1828). *Acinodendron fallax* (DC.) Kuntze, Revis. Gen. Pl. 1: 244 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M).

Section: Miconia.

Distribution: Venezuela, Guyana, Suriname, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast, Paraguay.

Miconia fanshawei Wurdack, Mem. New York Bot. Gard. 10 (4): 36 (1961). Type:—GUYANA. *Fanshawe F2872* (Holotype: NY; isotypes: K).

Section: Laceraria.

Distribution: Guyana.

Miconia fasciculata Gardner, London J. Bot. 1: 533 (1842). *Acinodendron fasciculatum* (Gardner) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Gardner 183* (Holotype: BM; isotypes: G, K, NY, P, US).

Miconia amygdalina Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 141 (1850).

Miconia persicifolia Naudin & A.St.-Hil. ex Triana, Trans. Linn. Soc. London 28: 119 (1871), *pro syn.*

Miconia fasciculata var. *robusta* Cogn. in Martius, Fl. bras. 14 (4): 355 (1887).

Miconia fasciculata ssp. *catharinensis* Wurdack, Sellowia 14: 154 (1962).

Section: Glossocentrum.

Distribution: Brazil Northeast, Brazil Southeast, Brazil South.

Miconia favosa (Desr.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 190 (1850).

Basionym: *Melastoma favosum* Desr. in Lamarck, Encycl. Méth. Bot. 4: 47 (1797). *Acinodendron favosum* (Desr.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—HAITI. *Jaeger 43170* (Holotype: LE; isotypes: A, B, destroyed, S).

Section: Chaenopleura.

Distribution: Haiti.

Miconia ferreyrae Wurdack, Phytologia 7: 235 (1960). Type:—PERU. *Ferreyra 4297* (Holotype: US).

Section: Amblyarrhena.

Distribution: Peru.

Miconia ferruginata DC., Prodr. 3: 181 (1828). *Melastoma ferrugineum* Schrank & Mart. ex DC., Prod. 3: 181 (1828), *pro syn.* *Acinodendron ferruginatum* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M).

Miconia ferruginata var. *latifolia* DC., Prodr. 3: 181 (1828).

Melastoma pluknetii Schrank & Mart. ex DC., Prodr. 3: 181 (1828), *pro syn.*

Miconia triseriata Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 145 (1850).

Section: Miconia.

Distribution: Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia ferruginea (Desr.) DC., Prodr. 3: 182 (1828).

Basionym: *Melastoma ferrugineum* Desr. in Lamarck, Encycl. Méth. Bot. 4: 42 (1797). *Acinodendron ferrugineum* (Desr.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—HAITI. *Martin s.n.* (Holotype: P?).

Section: Chaenopleura.

Distribution: Haiti, Dominican Republic.

Miconia fictilis (J.F.Macbr.) Michelangeli & R.Goldenb., Phytotaxa 53: 30 (2012).

Basionym: *Icaria fictilis* J.F.Macbr., Publ. Field Mus. Nat. Hist. Bot. Series 4: 191 (1929). Type:—PERU. *Macbride 4382* (Holotype: F; isotypes: K, NY, US).

Section: Chaenopleura.

Distribution: Peru.

Miconia filamentosa Gleason, Bull. Torrey Bot. Club 65: 579 (1938). Type:—PERU. *Klug 938* (Holotype: NY; isotype: US).

Miconia filamentosa ssp. *caquetana* Wurdack, Phytologia 52: 63 (1982).

Section: Miconia.

Distribution: Colombia, Ecuador, Peru.

Miconia firma J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (4/1): 411 (1941). Type:—PERU. *Weberbauer* 6389 (Holotype: F).
Section: Chaenopleura.
Distribution: Peru.

Miconia fissa Gleason, Bull. Torrey Bot. Club 59: 363 (1932). Type:—BRAZIL. *Holt* 528 (Holotype: NY; isotype: US).
Section: Jucunda.
Distribution: Colombia, Venezuela, Brazil North.

Miconia flaccida Gleason, Bull. Torrey Bot. Club 58: 226 (1931). Type:—PERU. *Killip* 25625 (Holotype: NY; isotypes: F, K, US).
Section: Tamonea.
Distribution: Peru.

Miconia flammea Casar., Nov. Stirp. Bras. Dec. 85 (1845). *Acinodendron flammeum* (Casar.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Riedel* s.n. (Holotype: TO).
Miconia flammea var. *major* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 839. (1891).
Section: Glossocentrum.
Distribution: Brazil Southeast, Brazil South.

Miconia flavescens Cogn. in Britton, Bull. Torrey Bot. Club 17: 93 (1890). *Acinodendron flavescens* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BOLIVIA. *Rusby* 2288 (Holotype: NY; isotypes: BR, F, MICH, NY, US).
Section: Chaenopleura.
Distribution: Bolivia.

Miconia floccosa Cogn., Bot. Jahrb. Syst. 42: 140 (1909). Type:—PERU. *Weberbauer* 2200 (Holotype: B, destroyed; isotypes: BR, MOL).
Section: Cremanium.
Distribution: Peru.

Miconia floribunda (Bonpl.) DC., Prodr. 3: 188 (1828).
Basionym: *Melastoma floribundum* Bonpl., Monogr. Melast. 1: 123 (1816). *Acinodendron floribundum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Bonpland* s.n. (Holotype: P).
Section: Amblyarrhena.
Distribution: Colombia, Venezuela, Ecuador, Peru.

Miconia fluminensis Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 362 (1915). Type:—BRAZIL. *Ule* 4229 (Holotype: B, destroyed; isotype: HBG).
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia formosa Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 842 (1891). *Acinodendron formosum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Glaziou* 16911 (Holotype: BR; isotypes: C, F, G, K, NY, P, R).
Miconia formosa var. *angustifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 842 (1891).
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia fosbergii Wurdack, Phytologia 11: 390 (1965). Type:—ECUADOR. *Fosberg* 22971 (Holotype: US; isotype: US).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia fosteri Wurdack in Harling & Sparre, Fl. Ecuador 13: 393 (1980). Type:—COLOMBIA. *Klug* 1864 (Holotype: NY; isotype: US).

Section: Miconia.

Distribution: Colombia, Ecuador, Peru, Bolivia, Brazil North.

Miconia foveolata Cogn., Jahrb. Königl. Bot. Gart. Berlin 4: 281 (1886). *Acinodendron foveolatum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PUERTO RICO. *Sintenis* 1335 (Holotype: B, destroyed; isotypes: BM, BR, CORD, GH, GOET, JE, K, L, LD, LIL, M, MICH, MO, MSC, NY, S, US).

Section: Chaenopleura.

Distribution: Puerto Rico.

Miconia fragilis Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 171 (1850). *Acinodendron naudinii* Kuntze, Revis. Gen. Pl. 2: 949 (1891). Type:—FRENCH GUIANA. *Martin* s.n. (Holotype: P; isotypes: BR, US).

Section: Miconia.

Distribution: Venezuela, Guyana, French Guiana.

Miconia fragrans Cogn. in Urban, Symb. Antill. 7: 526 (1913). Type:—TOBAGO. *Broadway* 4045 (Holotype: BR; isotypes: BM, GH, MO, NY, S, U, US).

Section: Miconia.

Distribution: Venezuela, Trinidad & Tobago, Guyana, Brazil North.

Miconia francavillana Cogn. in Martius, Fl. bras. 14 (4): 237 pl. 48 (1887). *Acinodendron francavillanum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Blanchet* 3448 (Lectotype: G-BOISS; isolectotype: BR).

Section: Jucunda.

Distribution: Guyana, Suriname, French Guiana, Brazil Northeast, Brazil Southeast.

Miconia friedmanorum Almeda & Umaña, Novon 3: 5 (1993). Type:—COSTA RICA. *Herrera* 1900 (Holotype: CR; isotypes: BR, CAS, F, MA, MEXU, MO, P, USJ).

Section: Miconia.

Distribution: Costa Rica, Panama.

Miconia frontinoana Cogn. & Gleason, Brittonia 1: 117 (1932). Type:—COLOMBIA. *Lehmann* 7500 (Holotype: K; isotypes: BR, F, NY, US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia fruticulosa Cogn., Bot. Jahrb. Syst. 42: 146 (1909). Type:—PERU. *Weberbauer* 3383 (Holotype: B, destroyed; isotype: BR).

Section: Chaenopleura.

Distribution: Peru.

Miconia fuertesii Cogn. in Urban, Symb. Antill. 7: 527 (1913). Type:—DOMINICAN REPUBLIC. *Fuertes* 1905 (Lectotype: NY).

Section: Chaenopleura.

Distribution: Haiti, Dominican Republic.

Miconia fuliginosa Wurdack, Mem. New York Bot. Gard. 16: 26 (1967). Type:—ECUADOR. *Prieto* P-314 (Holotype: US; isotypes: F, GH, NY, VEN).

Section: Cremanium.

Distribution: Ecuador.

Miconia fulvostellata L.O.Williams, Fieldiana, Bot. 29 (10): 571 (1963). Type:—GUATEMALA. *Steyermark* 49143a (Holotype: F).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Nicaragua.

Miconia funckii Wurdack, Phytologia 22: 400 (1972). *Octomeris schlimii* Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 379 (1851). *Heterotrichum schlimii* (Naudin) Triana, Trans. Linn. Soc. London 28: 134 (1871). Type:—VENEZUELA. *Funck 313* (Holotype: P).

Section: Octomeris.

Distribution: Venezuela.

Miconia furfuracea (Vahl) Griseb., Fl. Brit. W. Indian Is. 257 (1860).
Basionym: *Melastoma furfuraceum* Vahl, Eclog. Amer. 3: 13 (1807). *Acinodendron furfuraceum* (Vahl) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—"INDIA OCCIDENTALI". *Ziegler s.n.* (Lectotype: C).
Melastoma purpurascens Vahl ex DC., Prod. 3: 180 (1828). *Miconia purpuracea* (Vahl) Stehle & Quentin, Fl. Guad. Depend 2: 87 (1949).
Section: Tamonea.
Distribution: Jamaica, Guadeloupe, Dominica, Martinique, Santa Lucia.

Miconia galactantha Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850). *Cremanium serrulatum* Pav. ex D.Don, Mem. Wern. Nat. Hist. Soc. 4: 314 (1823). *Melastoma serrulatum* Pav. ex DC., Prodr. 3: 192 (1828), pro syn. *Miconia serrulata* (Pav. ex D.Don) Triana, Trans. Linn. Soc. London 28: 129 (1871), nom. illeg. *Acinodendron serrulatum* (Pav. ex D.Don) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Pavón s.n.* (Holotype: BM; isotypes: BR, P).
Section: Cremanium.
Distribution: Peru, Bolivia.

Miconia gentryi Wurdack, Phytologia 48: 241 (1981). Type:—PANAMA/COLOMBIA FRONTIER. *Gentry 28624* (Holotype: US; isotypes: COL, MO, PMA).
Section: Octomeris.
Distribution: Panama, Colombia.

Miconia gibba Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 12: 179 (1934). Type:—ECUADOR. *Diels 934* (Holotype: B, destroyed).
Section: Amblyarrhena.
Distribution: Colombia, Ecuador.

Miconia gigantea Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 937 (1891). *Acinodendron giganteum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Glaziou 17548* (Holotype: BR; isotypes: K, P).
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia gilva Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 836 (1891). *Acinodendron gilvum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Glaziou 11974* (Holotype: BR; isotypes: C, K, P, R).
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia glaberrima (Schltdl.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 243 (1850).
Basionym: *Melastoma glaberrimum* Schltdl., Linnaea 13 (5): 421 (1839). *Acinodendron glaberrimum* (Schltdl.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—MEXICO. *Schiiede s.n.* (Holotype: HAL; isotype: CAS).
Miconia pinetorum Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 229 (1850). *Acinodendron pinetorum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Miconia brachystyla Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 230 (1850).

Miconia myriocarpa Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 231 (1850).

Miconia glaberrima var. *australis* J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 184 (1929).

Section: Cremanium.

Distribution: Mexico, Belize, Guatemala, Honduras, El Salvador, Colombia, Peru.

Miconia glabrata Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 875 (1891). *Acinodendron glabratum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—MEXICO. *Galeotti 2963J* (Syntype: BR; isosyntypes: BR, LE).

Section: Amblyarrhena.
Distribution: Mexico.

Miconia glandulifera Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 737 (1891). *Acinodendron glanduliferum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Mathews* 1729 (Syntypes: G-BOISS, LE; isosyntypes: BM, BR, NY).

Section: Tamonea.
Distribution: Peru.

Miconia glandulistylo Wurdack, Phytologia 38: 287 (1978). Type:—ECUADOR. *Steyermark* 54268 (Holotype: F; isotype: NY).

Section: Amblyarrhena.
Distribution: Ecuador.

Miconia glaucescens Triana, Trans. Linn. Soc. London 28: 115 (1871). *Miconia tunicata* Griseb. ex Triana, Trans. Linn. Soc. London 28: 115 (1871), *pro syn.* *Acinodendron glaucescens* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Lechler* 2402 (Holotype, B, destroyed; isotypes: BR, F, K, NY, S).

Section: Miconia.
Distribution: Ecuador, Peru.

Miconia glazioviana Cogn. in Martius, Fl. bras. 14 (4): 301 (1887). *Acinodendron glaziovianum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Glaziou* 13850 (Holotype: BR; isotypes: BR, K, MO, P, R).

Section: Miconia.
Distribution: Brazil Southeast.

Miconia gleasoniana Wurdack, Brittonia 9: 105 (1957). Type:—COLOMBIA. *Pennell* 9366 (Holotype: NY; isotype: US).

Section: Cremanium.
Distribution: Colombia.

Miconia globulifera Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 139 (1850). *Acinodendron globuliferum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—MEXICO. *Schiede* 559 (Holotype: HAL; isotypes: CAS, F).

Section: Octomeris.
Distribution: Mexico, Guatemala.

Miconia globuliflora (Rich.) Cogn. in Martius, Fl. bras. 14 (4): 418 (1888).

Basionym: *Melastoma globuliflorum* Rich., Actes Soc. Hist. Nat. Paris 1: 109 (1792). *Acinodendron globuliflorum* (Rich.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—FRENCH GUIANA (PROBABLY MARTINIQUE). *Leblond s.n.* (Holotype: G-DEL; isotype: P).

Melastoma myricoides Desr. in Lamarck, Encycl. Meth. Bot. 4 (51) (1797). *Miconia myricoides* (Desr.) Triana, Trans. Linn. Soc. London 28: 129 (1871).

Melastoma thyrsoideum Vahl, Eclog. Am. 3: 13 (1807).

Miconia sieberi DC., Prodr. 3: 189 (1828). *Cremanium sieberi* (DC.) Griseb., Fl. Brit. W. Indian Is. 262 (1860).

Miconia vulcanica Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 213 (1850). *Acinodendron vulcanicum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). *Miconia globuliflora* var. *vulcanica* (Naudin) R.A.Howard & E.A.Kellogg, J. Arnold Arbor. 67: 243. (1986).

Miconia abortiva Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 232 (1850).

Miconia globuliflora Cham. ex Triana, Trans. Linn. Soc. London 28: 129 (1871), *pro syn.*

Miconia globuliflora var. *dominicae* R.A.Howard & E.A.Kellogg, J. Arnold Arbor. 67: 242 (1986).

Section: Cremanium.

Distribution: Costa Rica, Panama, Guadeloupe, Dominica, Martinique, Venezuela.

Miconia glomerata Triana, Trans. Linn. Soc. London 28: 110 (1871). *Acinodendron glomeratum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Ruiz & Pavón* 14/58 (Holotype MA?; isotypes: BM, F).

Section: Miconia.

Distribution: Peru.

Miconia glomerulifera Cogn., Bull. Torrey Bot. Club 23: 15 (1896). Type:—BOLIVIA. *Bang* 2856 (Holotype: BR; isotypes: CORD, GH, K, M, MIN, MO, NY, PH, US).

Section: Cremanium.

Distribution: Bolivia.

Miconia glutinosa Cogn., Bot. Jahrb. Syst. 42: 141 (1909). Type:—PERU. *Weberbauer* 1325 (Holotype: B, destroyed; isotype: BR).

Section: Amblyarrhena.

Distribution: Peru.

Miconia glyptophylla Wurdack, Phytologia 39: 322 (1978). Type:—ECUADOR. *Camp E-4566* (Holotype: US; isotypes: NY, S).

Section: Cremanium.

Distribution: Ecuador.

Miconia gonioclada Triana, Trans. Linn. Soc. London 28: 128 (1871). *Acinodendron goniocladium* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—ECUADOR. *Seemann* 766 (Holotype: K; isotypes: BR, TCD).

Section: Cremanium.

Distribution: Ecuador.

Miconia goniostigma Triana, Trans. Linn. Soc. London 28: 124 (1871). *Acinodendron goniostigma* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Triana* 3968 (Holotype: BM; isotypes: BR, K, NY, P).

Miconia megastigma Gleason, Bull. Torrey Bot. Club 68: 250 (1941).

Section: Amblyarrhena.

Distribution: Panama, Colombia, Ecuador.

Miconia gossypina Triana, Trans. Linn. Soc. London 28: 123 (1871). *Acinodendron gossypinum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Triana s.n.* (Holotype: BM).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia goudotii Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 155 (1850). *Acinodendron goudotii* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Goudot s.n.* (Holotype: P; isotype: BR).

Section: Miconia.

Distribution: Colombia, Venezuela.

Miconia gracilis Triana, Trans. Linn. Soc. London 28: 107 (1871). *Acinodendron gracile* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—NICARAGUA. *Seemann* 38 (Syntype: K).

Miconia ochroleuca Standl., Publ. Field Mus. Nat Hist., Bot. Ser. 11 (4): 138 (1932).

Section: Miconia.

Distribution: Mexico, Belize, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador.

Miconia grandidentata Almeda, Proc. Calif. Acad. Sci., Ser. 4 42: 311 (1981). Type:—COSTA RICA. *Almeda* 3627 (Holotype: CAS).

Section: Chaenopleura.

Distribution: Costa Rica, Panama.

Miconia grandiflora Cogn., Bot. Jahrb. Syst. 8: 25 (1886). *Acinodendron grandiflorum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Lehmann* 3742 (Holotype: K?; isotypes: BR, P?, US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia grandifoliata R.Goldenb. & Michelangeli, Phytotaxa 52: 29 (2012). *Miconia grandifolia* Ule, Notizbl. Bot. Gart. Mus. Berlin-Dahlem 6: 359 (1915), nom. illeg. Type:—PERU. Ule 6207 (Lectotype: K; isolectotype: HBG). Section: Miconia. Distribution: Ecuador, Peru, Brazil North.

Miconia gratissima Benth. ex Triana, Trans. Linn. Soc. London 28: 101 (1871). *Acinodendron gratissimum* (Benth. ex Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. Spruce 1189 (Holotype: K; isotypes: F, GH, K, M, NY, TCD).

Section: Jucunda. Distribution: Nicaragua, Panama, Colombia, Venezuela, Guyana, Suriname, French Guiana, Bolivia, Brazil North, Brazil Central-West.

Miconia grayana Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 896 (1891). *Acinodendron grayanum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—ECUADOR. *Grisar s.n.* (Syntypes: GH, LE, P; isosyntype: BR). Section: Amblyarrhena. Distribution: Ecuador.

Miconia grayumii Almeda, Proc. Calif. Acad. Sci., Ser. 4 46: 209 (1989). Type:—COSTA RICA. *McDowell* 576 (Holotype: CAS; isotypes: DUKE, MO). Section: Miconia. Distribution: Nicaragua, Costa Rica.

Miconia griffisii J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 188 (1929). Type:—PERU. *Macbride* 4382 (Holotype: F; isotypes: NY, US). Section: Chaenopleura. Distribution: Peru.

Miconia grisea Cogn., Bot. Jahrb. Syst. 42: 147 (1909). Type:—PERU. *Weberbauer* 3346 (Holotype: B, destroyed; isotypes: BR, F, MOL). Section: Chaenopleura. Distribution: Peru.

Miconia grossidentata Wurdack, Phytologia 50: 298 (1982). Type:—VENEZUELA. *Guanchez* 913 (Holotype: US; isotypes: MO, PORT, VEN). Section: Miconia. Distribution: Venezuela.

Miconia guaiquinimae Wurdack, Mem. New York Bot. Gard. 10 (4): 41 (1961). Type:—VENEZUELA. *Maguire* 33024 (Holotype: NY; isotypes: F, K, RB, S, US, VEN). *Miconia guaiquinimae* ssp. *angustifolia* Wurdack, Mem. New York Bot. Gard. 10 (4): 42 (1961). Section: Glossocentrum. Distribution: Venezuela.

Miconia guatemalensis Cogn. in Smith, Bot. Gaz. 16: 5 (1891). *Acinodendron guatemalense* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—GUATEMALA. von Türckheim 1317 (Syntype: BR; isosyntypes: K, NY, PH, US). Section: Octomeris. Distribution: Mexico, Guatemala, Honduras, El Salvador, Nicaragua.

Miconia guayaquilensis (Bonpl.) D.Don, Mem. Wern. Nat. Hist. Soc. 4: 316 (1823). Basionym: *Melastoma guayaquilense* Bonpl., Monogr. Melast. 1: 114 (1816). *Acinodendron guayaquilense* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—ECUADOR. *Bonpland s.n.* (Holotype: P). Section: Miconia. Distribution: Ecuador.

Miconia hadrophylla Wurdack, Phytologia 38: 290 (1978). Type:—COLOMBIA. *Garcia-Barriga 12946* (Holotype: US; isotype: COL).
Section: Amblyarrhena.
Distribution: Colombia, Ecuador.

Miconia hamata Cogn., Bot. Jahrb. Syst. 42: 141 (1909). Type:—PERU. *Weberbauer 4447* (Holotype: B, destroyed; isotypes: BR, MOL).
Section: Amblyarrhena.
Distribution: Peru.

Miconia harlingii Wurdack, Phytologia 26: 10 (1973). Type:—ECUADOR. *Harling 8175* (Holotype: GB; isotypes: MO, NY, S, US).
Section: Chaenopleura.
Distribution: Colombia, Ecuador.

Miconia haughtii (Gleason) Wurdack, Phytologia 7: 237 (1960).
Basionym: *Pachydesmia haughtii* Gleason, Phytologia 2: 430 (1948). Type:—COLOMBIA. *Haught 5201* (Holotype: NY; isotypes: COL, US).
Section: Chaenopleura.
Distribution: Colombia.

Miconia heliotropoides Triana, Trans. Linn. Soc. London 28: 116 (1871). *Acinodendron heliotropoides* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Gardner 3167* (Syntypes: BM, K, NY).
Section: Miconia.
Distribution: Bolivia, Brazil North, Brazil Central-West.

Miconia hematostemon Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 193 (1850). *Acinodendron hematostemon* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Linden 1062* (Holotype: P; isotypes: BR, NY).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia hemenostigma Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 230 (1850). *Acinodendron hemenostigma* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—MEXICO. *Galeotti 2908* (Holotype: P; isotype: BR).
Section: Cremanium.
Distribution: Mexico, Guatemala.

Miconia herpetica DC., Prodr. 3: 181 (1828). *Melastoma herpeticum* Schrank & Mart. ex DC., Prodr. 3: 181 (1828), *pro syn.* *Acinodendron herpeticum* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M; isotypes: BR, G-DC).
Miconia detergilis Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 149 (1850).
Miconia herpetica var. *acutifolia* Cogn. in Martius, Fl. bras. 14 (4): 287 (1887).
Section: Miconia.
Distribution: Brazil Northeast, Brazil Southeast.

Miconia herrerae Gleason, Amer. J. Bot. 19: 745 (1932). Type:—PERU. *Herrera 1575* (Holotype: US).
Section: Glossocentrum.
Distribution: Peru.

Miconia herzogii Cogn., Meded. Rijks-Herb. 19: 80 (1913). Type:—BOLIVIA. *Herzog 2285* (Holotype: BR; isotypes: S, M).
Section: Cremanium.
Distribution: Bolivia.

Miconia heterochaeta Wurdack, Phytologia 55: 140 (1984). Type:—ECUADOR. *Lugo 5431* (Holotype: GB; isotypes: F, NY, QCA, US).
Section: Miconia.
Distribution: Ecuador, Peru.

Miconia heteromera Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 124 (1850). *Acinodendron heteromerum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Poepig 1911* (Holotype: P; isotype: BR).
Clidemia simulans J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (4/1): 494 (1941).
Section: Miconia.
Distribution: Peru.

Miconia heterothrix Gleason & Wurdack, Phytologia 16: 173 (1968). Type:—MEXICO. *Hinton 9936* (Holotype: US; isotypes: DS, F, GH, K, LIL, RSA, U).
Section: Amblyarrhena.
Distribution: Mexico.

Miconia heterotricha Wurdack, Phytologia 5: 57 (1954). Type:—COLOMBIA. *Sandeman 6042* (Holotype: K; isotypes: COL, US).
Section: Miconia.
Distribution: Colombia.

Miconia hexamera Wurdack, Mem. New York Bot. Gard. 16: 21 (1967). Type:—ECUADOR. *Camp E-5185* (Holotype: US; isotypes: CAS, F, GH, MO, NY, SP, U, VEN).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia hexapetala Wurdack, Phytologia 48: 240 (1981). Type:—PERU. *Schunke 11432* (Holotype: US; isotypes: BR, F, MISSA, MO, NY, RB, RSA, TEX).
Section: Octomeris.
Distribution: Peru.

Miconia hildeana Kriebel & Almeda, Harvard Pap. Bot. 17: 60 (2012). Type:—PANAMA. *McPherson 11090* (Holotype: CAS; isotypes: MO, PMA).
Section: Cremanium.
Distribution: Costa Rica, Panama.

Miconia hirsutivena Gleason, Phytologia 2: 305 (1947). Type:—ECUADOR. *Steyermark 54142* (Holotype: NY; isotype: F).
Section: Cremanium.
Distribution: Ecuador.

Miconia hirta Cogn. in Rusby, Mem. Torrey Bot. Club 3: 31 (1893). Type:—BOLIVIA. *Bang 491* (Holotype: BR; isotypes: GH, K, M, MICH, NY, PH).
Section: Amblyarrhena.
Distribution: Bolivia.

Miconia hirtella Cogn. in Martius, Fl. bras. 14 (4): 423 (1888). *Acinodendron hirtellum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Blanchet 40 or 3792* (Syntype: BR; isosyntypes: BM, P).

Miconia hirtella var. *ovata* Cogn. in Martius, Fl. bras. 14 (4): 424 (1888).

Section: Cremanium.
Distribution: Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia histothrix Wurdack, Brittonia 40: 12 (1988). Type:—PERU. *Vigo 7573* (Holotype: US; isotype: MO).
Section: Miconia.
Distribution: Peru.

Miconia holosericea (L.) DC., Prodr. 3: 181 (1828).

Basionym: *Melastoma holosericeum* L., Sp. Pl. 1: 390 (1753). *Pleroma holosericea* (L.) D.Don, Mem. Wern. Nat. Hist. Soc. 4: 295 (1823). *Acinodendron holosericeum* (L.) Kuntze, Revis. Gen. Pl. 1: 244 (1891). Type:—BRAZIL OR SURINAME. *Hort. Cliff. s.n.* (Lectotype: BM).

Melastoma mucronatum Desr. in Lamarck, Encycl. Méth. Bot. 4: 46 (1797). *Diplochita mucronata* (Desr.) DC., Prodr. 3: 177 (1828). *Miconia mucronata* (Desr.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 120 (1850). *Miconia holosericea* var. *mucronata* (Desr.) Cogn. in Martius, Fl. bras. 14 (4): 237 (1887). *Acinodendron mucronatum* (Desr.) Kuntze, Revis. Gen. Pl. 1: 244 (1891).

Diplochita bracteata DC., Prodr. 3: 176 (1828). *Melastoma bracteatum* Mart. & Schrank ex DC., Prodr. 3: 176 (1828), pro syn. *Miconia holosericea* var. *bracteata* (DC.) Cogn. in Martius, Fl. bras. 14 (4): 237 (1887).

Miconia holosericea var. *subquintuplinervia* Cogn. in Martius, Fl. bras. 14 (4): 237 (1887).

Section: Jucunda.

Distribution: Mexico, Belize, Guatemala, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Trinidad & Tobago, Guyana, Surinam, French Guiana, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia hondurensis Donn.Sm., Bot. Gaz. 40: 3 (1905). Type:—HONDURAS. *Wilson* 575 (Holotype: US; isotype: NY).

Miconia gatunensis Pittier, J. Wash. Acad. Sci. 14 (19): 447 (1924).

Section: Tamonea.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama.

Miconia hookeriana Triana, Trans. Linn. Soc. London 28: 125 (1871). *Miconia pulverulenta* Hook., Bot. Mag. 89: 5411 (1863), nom. illeg. *Acinodendron hookerianum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Veitch* s.n. (Holotype: K).

Section: Miconia.

Distribution: Colombia, Ecuador, Peru.

Miconia hospitalis Wurdack, Phytologia 23: 475 (1972). Type:—PERU. *Schunke A-120* (Holotype: US).

Section: Tamonea.

Distribution: Peru.

Miconia howardiana Judd, Salzman & Skean, Brittonia 47: 416 (1995). Type:—DOMINICAN REPUBLIC. *Judd* 5178 (Holotype: FLAS; isotypes: DUKE, F, FLAS, JBSD, MICH, MO, NY, S, US).

Section: Chaenopleura.

Distribution: Dominican Republic.

Miconia huanucensis Wurdack, Phytologia 24: 201 (1972). Type:—PERU. *Asplund* 13173 (Holotype: S; isotype: S).

Section: Amblyarrhena.

Distribution: Peru.

Miconia huberi Wurdack, Mem. New York Bot. Gard. 50: 103 (1989). Type:—VENEZUELA. *Huber* 9519 (Holotype: US; isotypes: K, NY, U, VEN).

Section: Tamonea.

Distribution: Venezuela.

Miconia huigrensis Wurdack, Mem. New York Bot. Gard. 16: 14 (1967). Type:—ECUADOR. *Camp E-3375* (Holotype: US; isotypes: B, COL, F, GH, MO, NY).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia hutchisonii Wurdack, Phytologia 14: 273 (1967). Type:—PERU. *Hutchison* 6638 (Holotype: US; isotypes: F, K, M, MICH, MO, NY, UC, US, USM).

Section: Cremanium.

Distribution: Peru.

Miconia hyemalis A.St.-Hil. & Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 142 (1850). *Acinodendron hyemale* (A.St.-Hil. & Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Gaudichaud* 1362 (Holotype: P).
Section: Cremanium.
Distribution: Brazil Southeast, Brazil South, Uruguay.

Miconia hygrophila Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 220 (1850). *Acinodendron hygrophilum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BOLIVIA. *Weddell* 4705 (Holotype: P; isotype: NY).
Section: Cremanium.
Distribution: Peru, Bolivia.

Miconia hylophila Wurdack, Phytologia 55: 139 (1984). Type:—ECUADOR. *Brandbyge* 33885 (Holotype: US; isotype: AAU).
Section: Tamonea.
Distribution: Ecuador.

Miconia hymenanthera Triana, Trans. Linn. Soc. London 28: 130 (1871). *Acinodendron hymenantherum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—ECUADOR. *Jameson* 706 (Holotype: K; isotypes: BM, US).
Section: Chaenopleura.
Distribution: Colombia, Ecuador.

Miconia hyperprasina Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 186 (1850). *Acinodendron hyperprasimum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—MEXICO. *Linden* 640 (Holotype: P; isotypes: BR, G, K).
Section: Miconia.
Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica.

Miconia hypoides Urb. & Ekman, Ark. Bot. 22A (17): 38 (1929). Type:—HAITI. *Ekman* H7552 (Holotype: S; isotypes: NY, US).
Section: Chaenopleura.
Distribution: Haiti.

Miconia hypoleuca (Benth.) Triana, Trans. Linn. Soc. London 28: 119 (1871).
Basionym: *Chaenopleura hypoleuca* Benth., J. Bot. 2: 315 (1840). Type:—GUYANA. *Schomburgk* 392 (Holotype: K; isotypes: BM, E).
Chaenopleura ferruginea Crueg., Linnaea 20: 112 (1847). *Miconia cruegeriana* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).
Miconia blanchetiana Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 202 (1850).
Acinodendron hypoleucum (Benth.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).
Section: Chaeanthera.
Distribution: Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Brazil North, Brazil Northeast, Brazil Southeast.

Miconia ibaguensis (Bonpl.) Triana, Trans. Linn. Soc. London 28: 110 (1871).
Basionym: *Melastoma ibaguense* Bonpl., Monogr. Melast. 1: 105 (1815). *Tschudya ibaguensis* (Bonpl.) Griseb., Abh. Königl. Ges. Wiss. Gottingen 9: 47 (1860). *Acinodendron ibaguense* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Bonpland* 371 (Holotype: P).
Miconia auriculata DC., Prodr. 3: 185 (1828). *Melastoma auriculatum* Mart. ex DC., Prodr. 3: 185 (1828), *pro syn.*
Miconia auriculata var. *theiformis* DC., Prodr. 3: 185 (1828). *Melastoma theiforme* Schrank & Mart. ex DC., Prodr. 3: 185 (1828), *pro syn.* *Acinodendron theiforme* (DC.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Miconia maximiliana DC., Prodr. 3: 186 (1828). *Melastoma maximilianum* Mart. ex DC., Prod. 3: 186 (1828), *pro syn.*
Leandra ciliata Cham., Linnaea 10: 36 (1835). *Clidemia chamissois* Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 375 (1851).
Miconia lineata Schltld., Linnaea 13: 423 (1839). *Melastoma lineatum* Schltld., Linnaea 13: 423 (1839), *pro syn.*
Clidemia maculata Benth., J. Bot. 2: 310 (1840).
Clidemia miconioides Benth., J. Bot. 2: 310 (1840).
Miconia eriophora Miq., Comm. Phytogr. 2: 86 (1840).

Clidemia glabrata Steud., Flora 27 (42): 721 (1844). *Staphidium glabratum* (Steud.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 324 (1851). *Miconia ibaguensis* var. *glabrata* (Steud.) Cogn. in Martius, Fl. bras. 14 (4): 332 (1887).

Miconia eriophora f. *minor* Miq., Linnaea 18: 282 (1844).

Miconia auriculata var. *quintuplinervia* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 176 (1850).

Melastoma pilosum Sessé & Moç., Fl. mexic., ed. 2: 107 (1894).

Miconia sanctiramoni Kraenzl., Vierteljahrsschr. Naturf. Ges. Zurich 76: 157 (1931).

Clidemia virgata Pittier, Bol. Soc. Venez. Ci. Nat. 11: 24 (1947).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Cuba, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast, Brazil South, Paraguay.

Miconia ibarrae Almeda, Proc. Calif. Acad. Sci., Ser. 4 46: 212 (1989). Type:—MEXICO. *Sinaca* 738 (Holotype: CAS; isotypes: CAS, MEXU).

Section: Miconia.

Distribution: Mexico.

Miconia icosandra Gleason, Bull. Torrey Bot. Club 58: 239 (1931). Type:—PERU. *Killip* 25781 (Holotype: NY; isotypes: F, K, US).

Section: Glossocentrum.

Distribution: Peru.

Miconia idiogena Wurdack, Phytologia 31: 496 (1975). Type:—ECUADOR. *Harling* 12403 (Holotype: GB; isotypes: S, US).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia idroboi Wurdack, Phytologia 18: 154 (1969). Type:—COLOMBIA. *Idrobo* 837 (Holotype: US; isotypes: COL, U, US).

Section: Miconia.

Distribution: Colombia, Peru.

Miconia igniaria Bonpl. ex Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 197 (1850). *Acinodendron igniarium* (Bonpl. ex Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Bonpland* 3513 (Holotype: P).

Section: Amblyarrhena.

Distribution: Peru.

Miconia iluensis Wurdack, Mem. New York Bot. Gard. 10 (5): 165 (1964). Type:—VENEZUELA. *Maguire* 33436 (Holotype: US; isotypes: NY, VEN).

Section: Tamonea.

Distribution: Venezuela.

Miconia imbricata Gleason, Bull. Torrey Bot. Club 52: 382 (1925). Type:—COLOMBIA. *Pennell* 10503 (Holotype: NY; isotypes: GH, US).

Section: Octomeris.

Distribution: Colombia.

Miconia imitans Wurdack, Phytologia 31: 497 (1975). Type:—ECUADOR. *Lugo* 2421 (Holotype: GB; isotypes: F, MO, S, US).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia impetiolaris (Sw.) D.Don ex DC., Prodr. 3: 183 (1828).

Basionym: *Melastoma impetiolare* Sw., Prodr. 70 (1788). *Acinodendron impetiolare* (Sw.) Kuntze, Revis. Gen. Pl. 1: 244 (1891). *Tamonea impetiolaris* (Sw.) O.F.Cook & G.N.Collins, Contr. U.S. Natl. Herb. 8: 249 (1903). Type:— "IND. OCC." Swartz s.n. (Lectotype: S).

Melastoma macrophyllum Desr. in Lamarck, Encycl. Méth. Bot. 4: 44 (1797).

Miconia wydleriana DC., Mem. Melast. 77 (1828).

Miconia macrophylla Macfad., Fl. Jamaica 2: 84 (1850).

Miconia impetiolaris var. *pandurifolia* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 137 (1850).

Miconia impetiolaris var. *spruceana* Cogn. in Martius, Fl. bras. 14 (4): 272 (1887).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Cuba, Jamaica, Haiti, Dominican Republic, Puerto Rico, Guadeloupe, Dominica, Colombia, Venezuela, Guyana, Ecuador, Peru, Brazil North.

Miconia inaequalifolia Triana, Trans. Linn. Soc. London 28: 110 (1871). *Acinodendron inaequalifolium* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—VENEZUELA. *Spruce* s.n. (Holotype: K; isotype: BR)

Section: Miconia.

Distribution: Colombia, Venezuela, Brazil North.

Miconia inaequidens (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).

Basionym: *Cremanium inaequidens* DC., Prodr. 3: 194 (1828). *Acinodendron inaequidens* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Martius* s.n. (Holotype: M; isotype: G-DC).

Section: Chaenanthera.

Distribution: Brazil Southeast.

Miconia inamoena Pilg., Verh. Bot. Vereins Prov. Brandenburg 47: 171 (1905). Type:—PERU. *Ule* 6735 (Holotype: B, destroyed; isotypes: HBG, K, MG).

Section: Amblyarrhena.

Distribution: Peru.

Miconia inanis Cogn. & Gleason, Brittonia 1: 118 (1932). Type:—ECUADOR. *Lehmann* 5658 (Holotype: K; isotype: BR).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia incachacana Wurdack, Phytologia 23: 483 (1972). Type:—BOLIVIA. *Steinbach* 9459 (Holotype: US; isotypes: F, MO, U).

Section: Amblyarrhena.

Distribution: Bolivia.

Miconia inconspicua Miq., Linnaea 22: 542 (1849). *Acinodendron inconspicuum* (Miq.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Regnell* II.110 (Holotype: S).

Miconia excelsa A.St.-Hil. ex Triana, Trans. Linn. Soc. London 28: 126 (1871).

Miconia inconspicua var. *lanceolata* Cogn. in Martius, Fl. bras. 14 (4): 415 (1888).

Miconia conferta Cogn. in Martius, Fl. bras. 14 (4): 416 (1888). *Acinodendron confertum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Section: Cremanium.

Distribution: Brazil Southeast, Brazil South.

Miconia incurva Gleason, Bull. Torrey Bot. Club 65: 580 (1938). Type:—COSTA RICA. *Skutch* 3273 (Holotype: NY; isotypes: K, MO, S, US).

Miconia austin-smithii Standl. & L.O.Williams, Brittonia 15: 25 (1963).

Section: Amblyarrhena.

Distribution: Costa Rica.

Miconia ingens Wurdack, Phytologia 14: 268 (1967). Type:—COLOMBIA. von Sneidern 751 (Holotype: S; isotypes: NY, S, US).

Section: Octomeris.

Distribution: Colombia.

Miconia innata Gleason, Phytologia 2: 304 (1947). Type:—ECUADOR. Steyermak 54309 (Holotype: F).

Section: Amblyarrhena.

Distribution: Ecuador, Peru.

Miconia insueta Wurdack, Brittonia 28: 139 (1976). Type:—COLOMBIA. Barclay 6790 (Holotype: US; isotype: COL).

Section: Cremanium.

Distribution: Colombia.

Miconia insularis Gleason in Johnston, Sargentia 8: 234 (1949). Type:—PANAMA. Johnston 448 (Holotype: NY; isotypes: BM, GH, MO, US).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Costa Rica, Panama, Ecuador.

Miconia intricata Triana, Trans. Linn. Soc. London 28: 105 (1871). *Acinodendron intricatum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. Triana 4003 (Holotype: BM; isotypes: BR, CAS, E, G, K, NY, P).

Section: Tamonea.

Distribution: Costa Rica, Panama, Colombia, Ecuador.

Miconia ioneura Griseb., Abh. Konigl. Ges. Wiss. Gottingen 24: 129 (1879). *Acinodendron ioneurum* (Griseb.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—ARGENTINA. Hieronymus 449 (Holotype: B, destroyed; isotypes: BR, CORD, K, S, US).

Section: Cremanium.

Distribution: Bolivia, Argentina.

Miconia irwinii Wurdack, Phytologia 29: 144 (1974). Type:—BRAZIL. Irwin 9385 (Holotype: US; isotypes: F, NY, S, SP, TEX).

Section: Miconia.

Distribution: Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia iteophylla Almeda, Proc. Calif. Acad. Sci., Ser. 4 46: 214 (1989). Type:—PANAMA. Hammel 3393 (Holotype: CAS; isotype: MO).

Section: Miconia.

Distribution: Panama.

Miconia jahni Pittier, J. Wash. Acad. Sci. 14 (19): 450 (1924). Type:—VENEZUELA. Jahn 876 (Holotype: VEN; isotype: US).

Miconia cuneifolia Gleason, Bull. Torrey Bot. Club 52: 451 (1925).

Section: Cremanium.

Distribution: Colombia, Venezuela, Ecuador.

Miconia javorkaeana Borhidi, Acta Bot. Hung. 29: 187 (1983). *Graffenrieda cordifolia* Alain, Contr. Ocas. Mus. Hist. Nat. Colegio "De La Salle" 14: 1 (1955). *Miconia cordifolia* (Alain) Borhidi, Abstracta Botanica (Budapest) 5: 32 (1977), nom. illeg.. Type:—CUBA. Clemente 4407 (Holotype: HAC; isotype: NY).

Section: Octomeris.

Distribution: Cuba.

Miconia jefensis Almeda, Proc. Calif. Acad. Sci., Ser. 4 52: 43 (2000). Type:—PANAMA. Almeda 5826 (Holotype: CAS; isotypes: MO, NY, PMA).

Section: Miconia.

Distribution: Panama.

Miconia jentaculorum Wurdack, Phytologia 11: 393 (1965). Type:—COLOMBIA. Cuatrecasas 13530 (Holotype: US; isotypes: COL, K, NY).

Section: Cremanium.

Distribution: Colombia.

Miconia jimenezii Judd & R.S.Beaman, Brittonia 40: 386 (1988). Type:—DOMINICAN REPUBLIC. Judd 5192 (Holotype: FLAS; isotypes: A, DUKE, F, FLAS, JBSD, MO, NY, S, US).

Section: Chaenopleura.

Distribution: Dominican Republic.

Miconia jitotolana Wurdack, Phytologia 14: 270 (1967). Type:—MEXICO. Breedlove 11951 (Holotype: US; isotypes: DS, F, LL, WIS).

Section: Octomeris.

Distribution: Mexico.

Miconia johnwurdackiana Baumgratz & D'El Rei Sousa, Bradea 10: 26 (2004). Type:—BRAZIL. Hatschbach 47884 (Holotype: RB; isotypes: CAS, FLOR, K, MBM).

Section Uncertain

Distribution: Brazil Northeast.

Miconia jorgensenii Wurdack, Mem. New York Bot. Gard. 16: 28 (1967). Type:—ECUADOR. Jorgensen JP-58 (Holotype: US; isotype: NY).

Section: Cremanium.

Distribution: Ecuador.

Miconia jucunda (DC.) Triana, Trans. Linn. Soc. London 28: 101 (1871).

Basionym: *Osbeckia jucunda* DC., Prodr. 3: 139 (1828). *Rhexia jucunda* Schrank & Mart. ex DC., Prodr. 3: 139 (1828), *pro syn. Graffenrieda jucunda* (DC.) Mart., Nov. Gen. Sp. 3: 145 (1832). *Acinodendron jucunda* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). *Tamonea jucunda* (DC.) Krasser in Engler & Prantl, Nat. Pflanzenfam. 3 (7): 188 (1893). Type:—BRAZIL. Martius s.n. (Holotype: M; isotypes: G-DC, M).

Miconia staminea var. *oblongata* DC., Prodr. 3: 187 (1828).

Jucunda martiana Cham., Linnaea 9: 456 (1834). *Miconia jucunda* var. *martiana* (Cham.) Cogn. in Martius, Fl. bras. 14 (4): 229 (1887).

Jucunda sellowiana Cham., Linnaea 9: 456 (1834). *Miconia jucunda* var. *selloana* (Cham.) Cogn. in Martius, Fl. bras. 14 (4): 229 (1887).

Jucunda olfersiana Cham., Linnaea 9: 457 (1834). *Miconia olfersiana* (Cham.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 123 (1850). *Miconia jucunda* var. *olfersiana* (Cham.) Cogn. in Martius, Fl. bras. 14 (4): 229 (1887).

Miconia jucunda var. *parvifolia* Cogn. in Martius, Fl. bras. 14 (4): 230 (1887).

Graffenrieda brevicalcarata Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 10: 53 (1927).

Section: Jucunda.

Distribution: Brazil Northeast, Brazil Southeast, Brazil South.

Miconia juruensis Pilg., Verh. Bot. Vereins Prov. Brandenburg 47: 172 (1905). Type:—BRAZIL. Ule 5827 (Lectotype: K; isolectotypes: HBG, US).

Miconia japuraensis Cogn., Bol. Mus. Paraense Hist. Nat. Ethnogr. 5: 253 (1909).

Section: Tamonea.

Distribution: Colombia, Peru, Bolivia, Brazil North.

Miconia kappellei Almeda & Kriebel, Harvard Pap. Bot. 17: 61 (2012). Type:—COSTA RICA. Herrera 5911 (Holotype: INB; isotypes: CAS, CR, MO).

Section: Cremanium.

Distribution: Costa Rica.

Miconia kappleri Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 162 (1850). *Acinodendron kappleri* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—FRENCH GUIANA. *Kappler* 1675 (Holotype: P; isotypes: M, NY).
Section: Miconia.
Distribution: Venezuela, Trinidad & Tobago, Suriname, French Guiana, Brazil Northeast.

Miconia kavanayensis Wurdack, Phytologia 18: 152 (1969). Type:—VENEZUELA. *Lasser* 1879 (Holotype: VEN; isotype: NY).
Section: Miconia.
Distribution: Venezuela, Brazil North.

Miconia klotzschii Triana, Trans. Linn. Soc. London 28: 118 (1871). *Acinodendron klotzschii* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Sello* 1126 (Holotype: B, destroyed; isotype: K).
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia klugii Gleason, Bull. Torrey Bot. Club 58: 240 (1931). Type:—PERU. *Klug* 1141 (Holotype: NY; isotypes: F, US).
Section: Glossocentrum.
Distribution: Peru, Bolivia.

Miconia koepckeana Wurdack, Phytologia 23: 485 (1972). Type:—PERU. *Dudley* 13143 (Holotype: US).
Section: Amblyarrhena.
Distribution: Peru.

Miconia kollmannii R.Goldenb. & Reginato, Brittonia 59: 337 (2007). Type:—BRAZIL. *Kollmann* 5064 (Holotype: MBML; isotypes: RB, UPCB).
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia kraenzlinii Cogn., Bot. Jahrb. Syst. 8: 22 (1886). *Acinodendron kraenzlinii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Lehmann* 3025 (Holotype: B, destroyed; isotypes: BR, US).
Miconia kraenzlinii ssp. *aequatorialis* Wurdack, Phytologia 35: 244 (1977).
Section: Tamonea.
Distribution: Colombia, Ecuador.

Miconia kriegeriana Baumgratz & Chiaveg., Acta Bot. Brasil. 20: 484 (2006). Type:—BRAZIL. *Chiavegatto* 87 (Holotype: RB; isotypes: CESJ, SPF).
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia krugii Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 932 (1891). *Acinodendron krugii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—HAITI. *de Menonville* s.n. (Holotype: MPU - "Herb. Thouin", NOT LOCATED)

Miconia ocoana Urb. & Ekman, Ark. Bot. 23A (11): 20 (1931).
Section: Chaenopleura.
Distribution: Haiti, Dominican Republic.

Miconia kuntzei Cogn. in Kuntze, Revis. Gen. Pl. 3 (2): 93 (1898). *Acinodendron kuntzei* (Cogn.) Kuntze, Revis. Gen. Pl. 3: 93 (1898). Type:—BOLIVIA. *Kuntze* s.n. (Holotype: B, destroyed; isotypes: BR, NY).
Section: Amblyarrhena.
Distribution: Bolivia.

Miconia labiakiana R.Goldenb. & C.V.Martin, Harvard Pap. Bot. 13: 225 (2008). *Miconia plumosa* Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 9: 54 (1927), nom. illeg. Type:—BRAZIL. *Luetzelburg* 12336 (Holotype: M; isotype: US).

Section: Octomeris.

Distribution: Brazil Southeast.

Miconia lacera (Bonpl.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 152 (1850).

Basionym: *Melastoma lacerum* Bonpl., Monogr. Melast. 1: 9 (1807). *Clidemia lacera* (Bonpl.) DC., Prodr. 3: 164 (1828). *Acinodendron lacerum* (Bonpl.) Kuntze, Revis. Gen. Pl. 1: 244 (1891). Type:—COLOMBIA. *Humboldt & Bonpland s.n.* (Holotype: P).

Miconia villosissima Crueg., Linnaea 20: 109 (1847).

Clidemia rhodolasia Beurl., Kongl. Vetensk. Acad. Handl. 40: 126 (1854).

Section: Hartigia.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Trinidad & Tobago, Ecuador, Peru.

Miconia lachnoclada Wurdack, Phytologia 23: 501 (1972). Type:—PERU. *Madison 10378* (Holotype: US; isotype: NA).

Section: Chaenopleura.

Distribution: Peru.

Miconia laciniata Wurdack, Phytologia 69: 320 (1990). Type:—PERU. *Sagástegui 12994* (Holotype: US; isotypes: F, HUT).

Section: Cremanium.

Distribution: Peru.

Miconia laeta Cogn. in Martius, Fl. bras. 14 (4): 614 (1888). *Acinodendron laetum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Glaziou n. B* (Holotype: B, destroyed; isotype: BR).

Miconia laeta var. *fastigiata* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 854 (1891).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia laetivirens L.Uribe, Caldasia 9: 457 (1966). Type:—COLOMBIA. *Uribe-Uribe 5564* (Holotype: COL; isotypes: NY, US).

Section: Cremanium.

Distribution: Colombia.

Miconia laevigata (L.) DC., Prodr. 3: 188 (1828).

Basionym: *Melastoma laevigatum* L., Syst. Nat., ed. 10 2: 1022 (1759). *Acinodendron laevigatum* (L.) Kuntze, Revis. Gen. Pl. 1: 244 (1891). Type:—JAMAICA. *S.coll., s.n.* (Lectotype: LINN 559.10).

Melastoma nicotianifolium Desr. in Lamarck, Encycl. Méth. Bot. 4: 53 (1797). *Miconia nicotianifolia* (Desr.) DC., Prodr. 3: 188 (1828).

Melastoma piluliferum Vahl, Eclog. Amer. 3: 15 (1807).

Melastoma expansum Vahl, Eclog. Amer. 3: 21 (1807).

Melastoma brachiatum Vahl, Eclog. Amer. 3: 24 (1807).

Melastoma virescens Vahl, Eclog. Amer. 3: 18 (1807). *Miconia virescens* (Vahl) Triana, Trans. Linn. Soc. London 28: 107 (1871).

Melastoma ambiguum Bonpl., Monogr. Melast. 1: 55 (1809). *Miconia ambigua* (Bonpl.) DC., Prodr. 3: 189 (1828). *Acinodendron ambiguum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Melastoma pendulifolium Bonpl., Monogr. Melast. 1: 79 (1811). *Miconia pendulifolia* (Bonpl.) DC., Prodr. 3: 187 (1828).

Melastoma portoricense Spreng., Neue Entdeck. 3: 61 (1822).

Miconia havanensis DC., Prodr. 3: 188 (1828).

Miconia ceanothina DC., Prodr. 3: 189 (1828).

Melastoma acuminatum Sieber ex K.Presl, Isis (Oken) 21: 272 (1828).

Melastoma asperum K.Presl, Isis (Oken) 21: 272 (1828).

Melastoma cuspidatum Willd. ex Cham., Linnaea 9: 458 (1834).

Melastoma striatum Willd. ex Cham., Linnaea 9: 458 (1834).

Jucunda bredemeyriana Cham., Linnaea 9: 458 (1834).

Jucunda rudolphiana Cham., Linnaea 9: 458 (1834).

Miconia parviflora Macfad., Fl. Jamaica 2: 90 (1850).

Miconia microcarpa Macfad. ex Griseb., Fl. Brit. W. Indian Is. 257 (1860), *pro syn.*

Miconia affinis Macfad. ex Griseb., Fl. Brit. W. Indian Is. 257 (1860), *pro syn.* *Acinodendron affine* (Macfad.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, El Salvador, Honduras, Nicaragua, Costa Rica, Cuba, Jamaica, Haiti, Dominican Republic, Puerto Rico, Virgin Islands, Saint Kitts & Nevis, Antigua, Montserrat, Guadeloupe, Dominica, Martinique, Santa Lucia, Saint Vincent, Barbados, Grenada, Saba, Colombia, Venezuela, Trinidad & Tobago, Ecuador, Peru.

Miconia laevipilis Wurdack, Phytologia 22: 402 (1972). *Heterotrichum racemosum* Wurdack, Phytologia 8: 171 (1962).

Type:—VENEZUELA. Pittier 15469 (Holotype: US; isotype: VEN).

Section: Octomeris.

Distribution: Venezuela.

Miconia lagunensis Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 358 (1915). Type:—BRAZIL. Ule 1667 (Holotype: B, destroyed; isotypes: HBG, S).

Section: Glossocentrum.

Distribution: Brazil South.

Miconia lambayequensis Wurdack, Phytologia 64: 297 (1988). Type:—PERU. Quiroz 1976 (Holotype: F; isotype: US).

Section: Amblyarrhena.

Distribution: Peru.

Miconia lamprarrhena Triana, Trans. Linn. Soc. London 28: 122 (1871). *Acinodendron lamprarhenum* (Triana) Kuntze, Revis. Gen. Pl. 1: 244 (1891). *Acinodendron lamprarrhenum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951. (1891).

Type:—COLOMBIA. Triana 4041 (Holotype: BM; isotype: US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia lamprophylla Triana, Trans. Linn. Soc. London 28: 108 (1871). *Acinodendron lamprophyllum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. Triana 1033 (Holotype: BM; isotypes: COL, K, P, US).

Section: Miconia.

Distribution: Nicaragua, Costa Rica, Colombia, Venezuela, Ecuador, Peru, Bolivia, Brazil North.

Miconia lanata (DC.) Triana, Trans. Linn. Soc. London 28: 113 (1871).

Basionym: *Clidemia lanata* DC., Prodr. 3: 162 (1828). *Melastoma lanatum* Lamb. ex DC., Prod. 3: 162 (1828), *pro syn.* *Staphidium lanatum* (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 324 (1851). *Tschudya lanata* (DC.) Griseb., Fl. Brit. W. Indian Is. 250 (1860). *Acinodendron lanatum* (DC.) Kuntze, Rev. Gen. Pl. 2: 951 (1891). Type:—TRINIDAD. Sieber 66 (Syntypes: G-DC, MO).

Section: Miconia.

Distribution: Cuba, Puerto Rico, Colombia, Trinidad & Tobago, Peru, Bolivia, Brazil North, Brazil Central-West.

Miconia lanceolata (Desr.) DC., Prodr. 3: 190 (1828).

Basionym: *Melastoma lanceolatum* Desr. in Lamarck, Encycl. Méth. Bot. 4: 43 (1797). *Acinodendron lanceolatum* (Desr.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—DOMINICAN REPUBLIC. Martin s.n. (Holotype: P-JU).

Miconia christii Cogn. in Urban, Symb. Antill. 6: 27 (1909).

Section: Chaenopleura.

Distribution: Haiti, Dominican Republic.

Miconia lappacea (DC.) Triana, Trans. Linn. Soc. London 28: 111 (1871).

Basionym: *Clidemia lappacea* DC., Prodr. 3: 162 (1828). *Melastoma lappaceum* Rich. ex DC., Prod. 3: 162 (1828), *pro syn.* *Staphidium lappaceum* (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 324 (1851). *Acinodendron lappaceum* (DC.) Kuntze, Rev. Gen. Pl. 2: 951 (1891). Type:—FRENCH GUIANA. *Richard s.n.* (Holotype: P).

Miconia lappacea var. *angustifolia* Cogn. in Martius, Fl. bras. 14 (4): 333 (1887).

Section: Miconia.

Distribution: Guyana, Suriname, French Guiana, Brazil North.

Miconia larenensis Gleason in Steyermark, Fieldiana, Bot. 28 (2): 436 (1952). Type:—VENEZUELA. *Steyermark 55275* (Holotype: F; isotypes: A, NY, VEN, US).

Section: Amblyarrhena.

Distribution: Venezuela.

Miconia lasiocalyx Cogn., Bull. Torrey Bot. Club 23: 278 (1896). Type:—BOLIVIA. *Bang 2344* (Holotype: BR; isotypes: CAS, CORD, F, GH, K, MIN, M, MICH, MO, NY, PH, S, US).

Miconia scabriuscula Cogn., Meded. Rijks-Herb. 19: 80 (1913).

Section: Amblyarrhena.

Distribution: Colombia, Ecuador, Peru, Bolivia.

Miconia lasiostyla Gleason, Bull. Torrey Bot. Club 58: 243 (1931). Type:—PERU. *Sawada 77* (Holotype: F).

Section: Amblyarrhena.

Distribution: Peru.

Miconia lasseri Gleason in Lasser & Maguire, Bol. Soc. Venez. Ci. Nat. 15: 103 (1954). Type:—VENEZUELA. *Lasser 1685* (Holotype: NY; isotypes: F, VEN).

Section: Jucunda.

Distribution: Venezuela, Guyana, Suriname, French Guiana, Brazil North.

Miconia latecrenata (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 239 (1850).

Basionym: *Cremanium latecrenatum* DC., Prodr. 3: 194 (1828). Type:—BRAZIL. *Martius s.n.* (Holotype: M; isotype: G-DC).

Melastoma pyramidale Mart. & Schrank ex DC., Prodr. 3: 194 (1828), *pro syn.*

Melastoma taquari Vell., Flora Flum. 4: 182 (1825/1829). *Acinodendron taquari* (Vell.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Cremanium chaenopleuroides Gardner, London J. Bot. 2: 349 (1843). *Miconia chaenopleuroides* (Gardner) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).

Section: Chaenanthera.

Distribution: Brazil Northeast, Brazil Southeast, Brazil South.

Miconia lateriflora Cogn., Bol. Mus. Paraense Hist. Nat. Ethnogr. 5: 255 (1909). Type:—BRAZIL. *Baker 116* (Holotype: BR; isotype MG).

Octopleura ciliata Triana, Trans. Linn. Soc. London 28: 146 (1871). *Ossaea ciliata* (Triana) Cogn. in de Candolle & de Candolle, Monogr. Phan 7: 1067 (1891).

Miconia tessmannii Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 9: 266 (1925).

Ossaea disparilis Standl., Contrib. Arnold Arbor. 5: 120 (1933). *Miconia disparilis* (Standl.) R.O.Williams, Fl. Trinidad & Tobago 1: 388 (1934).

Ossaea disparilis var. *adenophora* Standl., Contr. Arnold Arbor. 5: 121 (1933). *Miconia lateriflora* var. *adenophora* (Standl.) C.Nelson, Ceiba 42: 39 (2001).

Miconia lateriflora ssp. *monticellensis* Wurdack, Phytologia 16: 171 (1968).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Northeast.

Miconia latifolia (D.Don) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850).

Basionym: *Cremanium latifolium* D.Don, Mem. Wern. Nat. Hist. Soc. 4: 312 (1823). *Acinodendron latifolium* (D.Don) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. Ruiz & Pavón s.n. (Holotype: OXF; isotypes: B, destroyed, FI, MA).

Chiloporus andinus Naudin, Ann. Sci. Nat., Bot., Ser. 3 4: 57 (1845). *Miconia andina* (Naudin) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 236 (1850). *Acinodendron andinum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Cremanium nitidum Griseb. ex Triana, Trans. Linn. Soc. London 28: 130 (1871), *pro syn.*

Miconia epiphytica Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 934 (1891). *Acinodendron epiphyticum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Section: Chaenopleura.

Distribution: Colombia, Venezuela, Ecuador, Peru, Bolivia.

Miconia latistigma Cogn. in Rusby, Bull. New York Bot. Gard. 4: 360 (1907). Type:—BOLIVIA. Rusby s.n. (Holotype: NY; isotype: BR).

Section: Cremanium.

Distribution: Bolivia.

Miconia laurina (D.Don) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).

Basionym: *Cremanium laurinum* D.Don, Mem. Wern. Nat. Hist. Soc. 4: 313 (1823). *Acinodendron laurinum* (D.Don) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. Ruiz & Pavón s.n. (Holotype: OXF; isotypes: MA?, US).

Section: Cremanium.

Distribution: Peru.

Miconia laxa Wurdack, Phytologia 41: 2 (1978). Type:—ECUADOR. Sparre 18735 (Holotype: S; isotypes: NY, TEX, US).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia leandroides Cogn. & Gleason, Brittonia 1: 121 (1932). Type:—ECUADOR. Lehmann 6690 (Holotype: K; isotype: BR).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia lechleri Triana, Trans. Linn. Soc. London 28: 124 (1871). *Acinodendron lechleri* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. Lechler 2565 (Holotype: K; isotypes: BR, GOET).

Section: Miconia.

Distribution: Peru.

Miconia ledifolia (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 224 (1850).

Basionym: *Cremanium ledifolium* DC., Prodr. 3: 193 (1828). *Acinodendron ledifolium* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—ECUADOR. Bonpland s.n. (Holotype: P; isotypes: G-DC, S, SI).

Section: Cremanium.

Distribution: Ecuador.

Miconia lehmannii Cogn., Bot. Jahrb. Syst. 8: 29 (1886). *Acinodendron lehmannii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. Lehmann 28 (Holotype: BR; isotype: US).

Section: Cremanium.

Distribution: Colombia, Venezuela.

Miconia leiotricha Wurdack, Phytologia 22: 402 (1972). *Heterotrichum lucidum* Triana, Trans. Linn. Soc. London 28: 134 (1871). *Staphidium lucidum* Berg ex Triana, Trans. Linn. Soc. London 28: 134 (1871), *pro syn.* Type:—VENEZUELA. Fendler 1845 (Holotype: K; isotypes: BR, F, GH, NY, PH).

Section: Octomeris.

Distribution: Venezuela.

Miconia lennartanderssonii Michelangeli & R.Goldenb., Phytotaxa 52: 30 (2012). *Miconia anderssonii* Wurdack, Phytologia 52: 66 (1982), nom. illeg. Type:—ECUADOR. Harling 19235 (Holotype: US; isotype: GB).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia lenticellata Alain, Contr. Ocas. Mus. Hist. Nat. Colegio "De La Salle" 14: 3 (1955). Type:—CUBA. Alain 3746 (Holotype: HAC; isotype: NY).
Section: Octomeris.
Distribution: Cuba.

Miconia lepidota DC., Prodr. 3: 180 (1828). *Melastoma lepidotum* Schrank & Mart. ex DC., Prodr. 3: 180 (1828), pro syn. *Acinodendron lepidotum* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M). *Miconia schomburgkii* Benth., J. Bot. 2: 312 (1840).

Miconia bifrons Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 178 (1850).

Miconia naudiniana Miq., Stirp. Surinam. Select. 50 (1850).

Section: Miconia.

Distribution: Colombia, Venezuela, Guyana, Suriname, French Guiana, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia leptantha Urb. & Ekman, Ark. Bot. 22A (17): 35 (1929). Type:—HAITI. Ekman H10153 (Holotype: S).

Section: Glossocentrum.

Distribution: Haiti.

Miconia leucocarpa DC., Prodr. 3: 182 (1828). *Melastoma leucocarpum* Mart. & Schrank ex DC., Prod. 3: 182 (1828), pro syn. *Acinodendron leucocarpum* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M; isotypes: BR, G-DC).

Miconia pohliana Cogn. in Martius, Fl. bras. 14 (4): 349 (1887). *Acinodendron pohlianum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Section: Miconia.

Distribution: Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia licropora Wurdack, Phytologia 50: 302 (1982). Type:—COLOMBIA. Fosberg 19266 (Holotype: US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia liebmannii Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 821 (1891). *Acinodendron liebmannii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—MEXICO. Liebmann 62 (Holotype: BR; isotypes: C, G-DC, P).
Section: Miconia.

Distribution: Mexico.

Miconia liesneri Wurdack, Phytologia 48: 239 (1981). Type:—VENEZUELA. Steyermark 119642 (Holotype: US; isotypes: MO, VEN).

Section: Tamonea.

Distribution: Colombia, Venezuela.

Miconia ligulata Almeda, Proc. Calif. Acad. Sci., Ser. 4 46: 216 (1989). *Leandra consimilis* Gleason, Ann. Missouri Bot. Gard. 45: 268 (1958). Type:—PANAMA. Allen 2702 (Holotype: NY).

Section: Miconia.

Distribution: Nicaragua, Costa Rica, Panama, Colombia, Venezuela.

Miconia ligustrina (Sm.) Triana, Trans. Linn. Soc. London 28: 128 (1871).

Basionym: *Melastoma ligustrinum* Sm. in Rees, Cycl. 23: 27 (1812). *Auliphas ligustrina* (Sm.) Raf., Sylva Tellur. 95 (1838). *Acinodendron ligustrinum* (Sm.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Mutis s.n.* (Holotype: LINN 559.15).

Miconia rhamnoidea Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 226 (1850).

Miconia ligustrina var. *setulinodis* Wurdack, Phytologia 39: 325 (1978).

Section: Cremanium.

Distribution: Colombia, Ecuador.

***Miconia ligustroides* (DC.) Naudin**, Ann. Sci. Nat., Bot., Ser. 3 16: 167 (1850).

Basionym: *Cremanium ligustroides* DC., Prodr. 3: 194 (1828). *Melastoma ligustrinum* Mart. & Schrank ex DC., Prod. 3: 194 (1828), *pro syn.* *Acinodendron ligustroides* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). *Tamonea ligustroides* (DC.) Krasser in Engler & Prantl, Nat. Pflanzenfam. 3 (7): 188 (1893). Type:—BRAZIL. *Martius s.n.* (Holotype: M).

Melastoma densifrons Schrank & Mart. ex DC., Prodr. 3: 194 (1828), *pro syn.*

Melastoma nitens Schrank & Mart. ex DC., Prodr. 3: 194 (1828), *pro syn.*

Melastoma repandulum Schrank & Mart. ex DC., Prodr. 3: 194 (1828), *pro syn.*

Melastoma trineurum Vell., Fl. Flum. 4: 112 (1825/1829).

Miconia ligustroides var. *cordifolia* Cogn. in Martius, Fl. bras. 14 (4): 384 (1887).

Section: Glossocentrum.

Distribution: Brazil Central-West, Brazil Northeast, Brazil Southeast, Brazil South.

***Miconia lilacina* Triana**, Trans. Linn. Soc. London 28: 129 (1871). *Acinodendron lilacinum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Pearce s.n.* (Holotype: K; isotypes: BM, BR).

Section: Cremanium.

Distribution: Peru.

***Miconia limitaris* Wurdack**, Acta Bot. Venez. 13: 155 (1978). Type:—COLOMBIA. *Cuatrecasas* 25228 (Holotype: US; isotype: COL).

Section: Cremanium.

Distribution: Colombia, Venezuela.

***Miconia lithophila* L.Uribe**, Caldasia 11: 12 (1975). Type:—COLOMBIA. *Klug* 6178 (Holotype: COL; isotypes: NY, US).

Section: Amblyarrhena.

Distribution: Colombia.

***Miconia littlei* Wurdack**, Phytologia 13: 73 (1966). Type:—ECUADOR. *Little* 6177 (Holotype: US; isotype: K).

Section: Tamonea.

Distribution: Ecuador.

***Miconia livida* Triana**, Trans. Linn. Soc. London 28: 121 (1871). *Acinodendron lividum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. *Mathews* 1180 (Holotype: K; isotypes: BM, E, K).

Miconia donnellsmithii Cogn. in Smith, Bot. Gaz. 20: 287 (1895).

Section: Amblyarrhena.

Distribution: Mexico, Guatemala, Honduras, Costa Rica, Panama, Ecuador, Peru, Bolivia.

***Miconia lonchophylla* Naudin**, Ann. Sci. Nat., Bot., Ser. 3 16: 176 (1850). *Acinodendron lonchophyllum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—VENEZUELA. *Funck* 94 (Holotype: P; isotypes: BR, G, P).

Miconia sylvicola Pittier, Bol. Soc. Venez. Ci. Nat. 9: 123 (1944).

Section: Miconia.

Distribution: Mexico, Honduras, Costa Rica, Panama, Colombia, Venezuela.

***Miconia longibracteata* Almeda**, Proc. Calif. Acad. Sci., Ser. 4 42: 312 (1981). Type:—COSTA RICA. *Wilbur* 9845 (Holotype: DUKE; isotypes: CAS, US).

Section: Chaenopleura.

Distribution: Costa Rica.

Miconia longicuspidata S.S.Renner & R.Goldenb., Phytokeys 3: 36 (2011). *Miconia longicuspis* Herzog, Feddes Repert. Nov. Sp. 7: 64 (1909), nom. illeg. Type:—BOLIVIA. Herzog 326 (Holotype: Z).
Section: Miconia.
Distribution: Bolivia.

Miconia longicuspis Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 851 (1891). *Acinodendron longicuspe* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. Glaziou 16918 (Lectotype: BR; isolectotypes: K, MO, NY, P, R, Z).
Section: Glossocentrum.
Distribution: Brazil Southeast, Brazil South.

Miconia longifolia (Aubl.) DC., Prodr. 3: 184 (1828).
Basionym: *Melastoma longifolium* Aubl., Hist. pl. Guiane 1: 432 (1775). *Chaenopleura longifolia* (Aubl.) Griseb., Fl. Brit. W. Indian Is. 260 (1860). *Acinodendron longifolium* (Aubl.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—FRENCH GUIANA. Aublet s.n. (Holotype: BM).

Miconia lambertiana DC., Prodr. 3: 185 (1828).

Section: Glossocentrum.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Northeast.

Miconia longiracemosa Gleason, Bull. Torrey Bot. Club 58: 232 (1931). Type:—PERU. Williams 4249 (Holotype: NY; isotype: US).
Section: Miconia.
Distribution: Peru.

Miconia longisepala Gleason, Bull. Torrey Bot. Club 63: 536 (1936). Type:—PERU. Klug 3233 (Holotype: NY; isotypes: GH, K, MO, S, US, WIS).
Section: Amblyarrhena.
Distribution: Peru.

Miconia longisetosa Wurdack, Phytologia 41: 6 (1978). Type:—ECUADOR. Sodiro 490 (Holotype: BR).
Section: Chaenopleura.
Distribution: Ecuador.

Miconia longispicata Triana, Trans. Linn. Soc. London 28: 117 (1871). *Acinodendron longispicatum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. Spruce 2280 (Syntypes: BM, CAS, K, M, NY, TCD).

Miconia longipedunculata Cogn. in Martius, Fl. bras. 14 (4): 266: pl. 54 (1887). *Acinodendron longipedunculatum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Miconia longispicata var. *minor* Cogn. in Martius, Fl. bras. 14 (4): 267 (1887).

Section: Miconia.

Distribution: Colombia, Venezuela, Guyana, French Guiana, Bolivia, Brazil North.

Miconia loretensis Pilg., Verh. Bot. Vereins Prov. Brandenburg 47: 172 (1905). Type:—PERU. Ule 6734 (Holotype: B, destroyed; isotypes: HBG, MG).
Section: Miconia.
Distribution: Peru, Bolivia.

Miconia loreyoides Triana, Trans. Linn. Soc. London 28: 121 (1871). *Acinodendron loreyoides* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. Triana 4028 (Syntype: BM; isosyntypes: K, P).

Miconia killipii Gleason, Bull. Torrey Bot. Club 52: 387 (1925).

Section: Amblyarrhena.

Distribution: Costa Rica, Panama, Colombia, Guyana, Ecuador, Peru.

Miconia lourteigiana Wurdack, Phytologia 20: 380 (1970). Type:—PERU. Wurdack 1919 (Holotype: US; isotype: F).
Section: Miconia.

Distribution: Colombia, Venezuela, Ecuador, Peru, Bolivia, Brazil North.

Miconia loxensis (Bonpl.) DC., Prodr. 3: 186 (1828).

Basionym: *Melastoma loxense* Bonpl., Monogr. Melast. 1: 100 (1815). *Acinodendron loxensis* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—ECUADOR. *Bonpland* 3513 (Holotype: P; isotypes: BM, G-DEL).

Section: Cremanium.

Distribution: Ecuador, Peru.

Miconia luciana Gleason, J. Arnold Arbor. 28: 334 (1947). Type:—ST. LUCIA. *Beard* 477 (Holotype: NY; isotypes: A, K).

Section: Octomeris.

Distribution: Santa Lucia.

Miconia lucida Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 196 (1850). *Acinodendron lucidum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—VENEZUELA. *Linden* 307 (Lectotype: P; isolectotypes: BR, GH, P).

Miconia lucida ssp. *pariensis* Wurdack, Phytologia 16: 182 (1968).

Section: Amblyarrhena.

Distribution: Venezuela.

Miconia lugonis Wurdack, Phytologia 26: 397 (1973). Type:—ECUADOR. *Lugo* 1729 (Holotype: US; isotypes: F, GB, S).

Section: Miconia.

Distribution: Colombia, Ecuador, Peru, Brazil North.

Miconia lugubris Cogn., Bot. Jahrb. Syst. 42: 142 (1909). Type:—PERU. *Weberbauer* 3394 (Holotype: B, destroyed; isotypes: BR, MOL).

Section: Cremanium.

Distribution: Peru.

Miconia lundellii (Wurdack) Almeda, Novon 15: 510 (2005).

Basionym: *Clidemia lundellii* Wurdack, Wrightia 5: 227 (1976). Type:—GUATEMALA. *Lundell* 19626 (Holotype: US; isotypes: F, LL, S, US).

Section: Octomeris.

Distribution: Guatemala.

Miconia lurida Cogn. in Martius, Fl. bras. 14 (4): 356 (1887). *Acinodendron luridum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Riedel* s.n. (Syntype: BR; isosyntypes: K, LE).

Section: Glossocentrum.

Distribution: Brazil Northeast, Brazil Southeast.

Miconia luteola Cogn. in Urban, Symb. Antill. 6: 26 (1909). Type:—HAITI. *Ekman* H1407 (Neotype: S; isoneotypes: GH, IJ).

Miconia viejana Urb. & Ekman, Ark. Bot. 23A (11): 20 (1931).

Section: Chaenopleura.

Distribution: Haiti, Dominican Republic.

Miconia lutescens (Bonpl.) DC., Prodr. 3: 189 (1828).

Basionym: *Melastoma lutescens* Bonpl., Monogr. Melast. 1: 39 (1807). *Acinodendron lutescens* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—ECUADOR. *Bonpland* s.n. (Holotype: P; isotype: BM).

Miconia lutescens ssp. *piurensis* Wurdack, Phytologia 5: 59 (1954).

Section: Amblyarrhena.

Distribution: Ecuador, Peru.

Miconia luteynii Wurdack, Phytologia 50: 301 (1982). Type:—COLOMBIA. *Luteyn 7512* (Holotype: COL; isotypes: CAS, COL, MO, NY, US).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia lymanii Wurdack, Sellowia 14: 164 (1962). Type:—BRAZIL. *Smith 7878* (Holotype: US; isotypes: GH, HBR, NY, R).
Section: Cremanium.
Distribution: Brazil Southeast, Brazil South.

Miconia macayana Judd & Skean, Bull. Florida State Mus., Biol. Sci. 32: 145 (1987). Type:—HAITI. *Skean 1428* (Holotype: FLAS; isotypes: A, JBSD, NY).
Section: Chaenopleura.
Distribution: Haiti.

Miconia macbrydeana Wurdack, Phytologia 41: 3 (1978). Type:—ECUADOR. *MacBryde 983* (Holotype: US).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia machinazana C.Ulloa & D.A.Neill, Phytokeys 12: 35-46 (2012). Type:—ECUADOR. *Neill 16909* (Holotype: MO; isotypes: AAU, CAS, LOJA, M, NY, QCNE).
Section: Cremanium.
Distribution: Ecuador.

Miconia macrantha Triana, Trans. Linn. Soc. London 28: 123 (1871). *Acinodendron macranthum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Triana 8421* (Holotype: BM).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia macrodon (Naudin) Wurdack, Phytologia 22: 400 (1972).
Basionym: *Octomeris macrodon* Naudin, Ann. Sci. Nat., Bot., Ser. 3 4: 53 (1845). *Heterotrichum macrodon* (Naudin) Planch. ex Hook., Bot. Mag. 75: 4421 (1849). Type:—VENEZUELA. *Linden 265* (Lectotype: P; isolectotype: P).
Section: Octomeris.
Distribution: Venezuela.

Miconia macrothyrsa Benth., J. Bot. 2: 312 (1840). *Acinodendron macrothyrsum* (Benth.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—GUYANA. *Schomburgk I 398* (Holotype: K; isotypes: BM, NY, TCD).
Miconia pseudorubiginosa Cogn. in Urban, Symb. Antill. 7: 311 (1912).

Section: Miconia.
Distribution: Brazil Southeast, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast.

Miconia macrotis Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 749 (1891). *Acinodendron macrote* (Cogn.) Kuntze, Revis. Gen. Pl. 1: 244 (1891). Type:—TRINIDAD. *Crueger s.n.* (Syntype: K; isosyntype: BR).
Miconia macrotis var. *canescens* Gleason, Bull. Torrey Bot. Club 52: 381 (1925).
Section: Tamonea.
Distribution: Colombia, Venezuela, Trinidad & Tobago, Ecuador, Peru, Brazil North, Brazil Central-West.

Miconia madisonii Wurdack, Phytologia 23: 482 (1972). Type:—PERU. *Madison 10295* (Holotype: US; isotypes: MO, NA).
Section: Amblyarrhena.
Distribution: Peru.

Miconia madrensis Standl., Contr. U.S. Natl. Herb. 23: 1063 (1924). Type:—MEXICO. Rose 2226 (Holotype: US).
Section: Amblyarrhena.
Distribution: Mexico.

Miconia magdalena Triana, Trans. Linn. Soc. London 28: 115 (1871). Type:—COLOMBIA. Holton 910 (Holotype: K; isotypes: BR, NY, PH).
Acinodendron magdalena (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891).
Section: Miconia.
Distribution: Colombia, Venezuela.

Miconia maguirei Gleason, Bull. Torrey Bot. Club 75: 552 (1948). Type:—GUYANA. Maguire 23401 (Holotype: NY; isotype: K).
Section: Miconia.
Distribution: Guyana.

Miconia majalis Cogn., Bull. Acad. Belg., Ser. 3 14: 953 (1887). *Acinodendron majale* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. André 2768 (Holotype: K; isotypes: BR, F, NY).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia malatestae J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 181 (1929). Type:—PERU. Macbride 5176 (Holotype: F; isotypes: NY, US).
Section: Miconia.
Distribution: Peru.

Miconia manauara R.Goldenb., Caddah & Michelangeli, Brittonia 63: 246 246 (2011). Type:—BRAZIL. Ferreira 6723 (Holotype: INPA; isotypes: NY, US).
Section: Miconia.
Distribution: Brazil North.

Miconia mandonii Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 911 (1891). *Acinodendron mandonii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BOLIVIA. Mandon 646 (Syntype: BR; isosyntypes: CAS, K, MO, MPU, NY, P, S).
Section: Cremanium.
Distribution: Bolivia.

Miconia manicata Cogn. & Gleason, Brittonia 1: 122 (1932). Type:—COLOMBIA. Lehmann 6688 (Holotype: K; isotypes: BR, F, NY, US).
Miconia manicata ssp. *ecuadorensis* Wurdack, Phytologia 39: 328 (1978).
Section: Cremanium.
Distribution: Colombia, Venezuela, Ecuador.

Miconia mansfeldiana Urb. & Ekman, Ark. Bot. 23A (11): 21 (1931). Type:—DOMINICAN REPUBLIC. Ekman H11737 (Holotype: S; isotypes: EHH, GH, NY, S, US).
Section: Chaenopleura.
Distribution: Dominican Republic.

Miconia mapirensis Gleason, Bull. Torrey Bot. Club 68: 249 (1941). Type:—BOLIVIA. Krukoff 11215 (Holotype: NY; isotypes: A, K, LP, MO, S, U, UC, US, WIS).
Section: Amblyarrhena.
Distribution: Peru, Bolivia.

Miconia marginata Triana, Trans. Linn. Soc. London 28: 110 (1871). *Acinodendron marginatum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. Spruce 2469 (Holotype: K; isotypes: BM, BR, GH, K, NY, TCD).

Section: Miconia.

Distribution: Colombia, Venezuela, Guyana, Brazil North.

Miconia mariae Wurdack, Mem. New York Bot. Gard. 10 (5): 168 (1964). Type:—VENEZUELA. Steyermark 1369 (Holotype: US; isotypes: NY, VEN).

Section: Laceraria.

Distribution: Venezuela, Guyana.

Miconia markgraffii Urb. & Ekman, Ark. Bot. 22A (17): 41 (1929). Type:—HAITI. Ekman H7673 (Holotype: S).

Section: Chaenopleura.

Distribution: Haiti.

Miconia maroana Wurdack, Phytologia 16: 176 (1968). Type:—VENEZUELA. Williams 14431 (Holotype: US; isotypes: F, NY, VEN).

Section: Miconia.

Distribution: Colombia, Venezuela, Brazil North.

Miconia martiniana Gleason, Bull. Torrey Bot. Club 58: 233 (1931). Type:—PERU. Williams 7212 (Holotype: NY).

Section: Miconia.

Distribution: Peru.

Miconia martinicensis Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 913 (1891). *Acinodendron martinicense* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—MARTINIQUE. Hahn 1174 (Lectotype: P).

Section: Cremanium.

Distribution: Guadeloupe, Martinique.

Miconia matthaei Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 176 (1850). *Acinodendron matthaei* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—PERU. Mathews 1299 (Holotype: P; isotypes: BR, E, K).

Miconia matthaei var. *undulata* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 813 (1891).

Miconia wilsonii Cogn. in Urban, Symb. Antill. 7: 311 (1912).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Cuba, Jamaica, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West.

Miconia mattogrossensis Hoehne, Anexos Mem. Inst. Butantan, Secc. Bot. 1 (5): 147 (1922). Type:—BRAZIL. Hoehne 5447 (Syntype: SP).

Section: Glossocentrum.

Distribution: Brazil North, Brazil Central-West.

Miconia maximowicziana Cogn. in Martius, Fl. bras. 14 (4): 355 (1887). *Acinodendron maximowiczianum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. Riedel 1205 (Holotype: BR; isotypes: K, P).

Miconia maximowicziana var. *major* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 833 (1891).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia mazanana J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (4/1): 428 (1941). Type:—PERU. Schunke 374 (Holotype: F; isotypes: A, NY, S, US).

Section: Tamonea.

Distribution: Colombia, Ecuador, Peru, Brazil North.

Miconia mazatecana J.R.Santiago, Novon 20: 78 (2010). Type:—MEXICO. Santiago 2076 (Holotype: MEXU; isotypes: CAS, ENCB, FCME, IEB, MO, OAX, US, XAL).

Section: Amblyarrhena.

Distribution: Mexico.

Miconia mcvaughii Wurdack, Phytologia 16: 179 (1968). Type:—MEXICO. *McVaugh 13978* (Holotype: MICH).

Section: Amblyarrhena.

Distribution: Mexico.

Miconia media (D.Don) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850).

Basionym: *Cremanium medium* D.Don, Mem. Wern. Nat. Hist. Soc. 4: 313 (1823). *Acinodendron medium* (D.Don) Kuntze, Revis. Gen. Pl. 2: 951 (1891). *Tamonea media* (D.Don) Krasser in Engler & Prantl, Nat. Pflanzenfam. 3 (7): 142 (1893). Type:—PERU. *Ruiz & Pavón s.n.* (Holotype: OXF?; isotypes: B, destroyed, MA?).

Cremanium tinctorium DC., Prodr. 3: 193 (1828). *Miconia tinctoria* (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 229 (1850).

Miconia media ssp. *borealis* Wurdack, Phytologia 9: 421 (1964).

Miconia media ssp. *cajamaricensis* Wurdack, Phytologia 14: 274 (1967).

Miconia media ssp. *ecuadorensis* Wurdack, Mem. New York Bot. Gard. 16: 29 (1967).

Section: Cremanium.

Distribution: Ecuador, Peru.

Miconia mediocris Wurdack, Phytologia 39: 324 (1978). Type:—ECUADOR. *Camp E-705* (Holotype: NY; isotype: US).

Section: Cremanium.

Distribution: Ecuador.

Miconia medusa Gleason, Phytologia 2: 429 (1948). Type:—ECUADOR. *Espinosa 1559* (Holotype: NY; isotype: US).

Section: Cremanium.

Distribution: Ecuador.

Miconia megalantha Gleason, Phytologia 1: 33 (1933). Type:—COLOMBIA. *Lawrance 431* (Holotype: NY; isotypes: A, GH, MO, S, US, WIS).

Section: Octomeris.

Distribution: Colombia.

Miconia melanotricha (Triana) Gleason, Brittonia 3: 101 (1939).

Basionym: *Clidemia melanotricha* Triana, Trans. Linn. Soc. London 28: 136 (1871). Type:—COSTA RICA. *Hoffmann s.n.* (Holotype: B, destroyed; isotype: F).

Miconia atrosanguinea Cogn., Bot. Jahrb. Syst. 8: 23 (1886). *Acinodendron atrosanguineum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Miconia melanotricha var. *panamensis* Gleason, Ann. Missouri Bot. Gard. 45: 302 (1958).

Section: Chaenopleura.

Distribution: Costa Rica, Panama.

Miconia melinonis Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 125 (1850). *Acinodendron melinonis* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—FRENCH GUIANA. *Melinon 178* (Holotype: P; isotypes: BR, NY, U, US).

Section: Jucunda.

Distribution: French Guiana, Brazil North.

Miconia mellina DC., Prodr. 3: 182 (1828). *Melastoma mellinum* Schrank & Mart. ex DC., Prod. 3: 182 (1828), *pro syn.* *Acinodendron mellinum* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M; isotypes: BR, G-DC).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia membranacea Triana, Trans. Linn. Soc. London 28: 115 (1871). *Acinodendron membranaceum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Ruiz & Pavón s.n.* (Holotype MA?; isotypes: BM, BR).

Section: Miconia.

Distribution: Peru.

Miconia mendoncae Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 772 (1891). *Acinodendron mendoncae* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Mendonca* 1216 (Holotype: B, destroyed; isotype: BR).

Section: Miconia.

Distribution: Brazil Southeast.

Miconia meridensis Triana, Trans. Linn. Soc. London 28: 102 (1871). *Acinodendron meridense* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—VENEZUELA. *Moritz* 946 (Syntypes: BR, K, NY, PH).

Section: Octomeris.

Distribution: Venezuela.

Miconia mesmeana Gleason, Phytologia 1: 44 (1933). Type:—COLOMBIA. *Killip* 20012 (Holotype: NY; isotypes: A, GH, US, VEN).

Miconia mesmeana ssp. *longipetiolata* Wurdack, Phytologia 22: 414 (1972).

Miconia mesmeana ssp. *jabonensis* Wurdack, Phytologia 22: 415 (1972).

Section: Cremanium.

Distribution: Colombia, Venezuela.

Miconia metallica (Naudin) Triana, Trans. Linn. Soc. London 28: 106 (1871).

Basionym: *Cyanophyllum metallicum* Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 325 (1851). *Acinodendron metallicum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—VENEZUELA. *Funk* 1078 (Syntypes: BM, BR, G, MPU)

Section: Tamonea.

Distribution: Venezuela.

Miconia mexicana (Bonpl.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850).

Basionym: *Melastoma mexicanum* Bonpl., Monogr. Melast. 1: 128 (1816). *Conostegia mexicana* (Bonpl.) Ser. ex DC., Prodr. 3: 175 (1828). *Acinodendron mexicanum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—MEXICO? Not located (Holotype: P?).

Melastoma tunicatum Bonpl., Monogr. Melast. 1: 141 (1816). *Conostegia tunicata* (Bonpl.) Ser. ex DC., Prodr. 3: 175 (1828). *Miconia tunicata* (Bonpl.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 243 (1850). *Acinodendron tunicatum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 953 (1891).

Miconia conostegioides Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 127 (1850). *Miconia mexicana* var. *conostegioides* (Naudin) Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 763 (1891).

Miconia tococoides Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 127 (1850).

Miconia lauriformis Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 189 (1850). *Acinodendron lauriforme* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Miconia humilis Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 764 (1891). *Acinodendron humile* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). *Tamonea humilis* (Cogn.) Krasser in Engler & Prantl, Nat. Pflanzenfam. 3 (7): 188 (1893).

Miconia purpusii Brandegee, Univ. Calif. Publ. Bot. 10: 185 (1922).

Section: Laceraria.

Distribution: Mexico, Guatemala, Honduras, El Salvador, Nicaragua, Costa Rica.

Miconia micayana Wurdack, Phytologia 7: 237 (1960). *Amphitoma flavescens* Gleason, Bull. Torrey Bot. Club 52: 378 (1925). Type:—COLOMBIA. *Killip* 11884 (Holotype: NY).

Section: Cremanium.

Distribution: Colombia.

Miconia michelangeliana R.Goldenb. & L.Kollmann, Blumea 55: 139 (2010). Type:—BRAZIL. *Kollmann* 10743 (Holotype: MBML; isotypes: CEPEC, NY, RB, UPCB).

Section: Tamonea.

Distribution: Brazil Southeast.

Miconia micrantha Cogn., Bull. Torrey Bot. Club 23: 16 (1896). Type:—BOLIVIA. Bang 2858 (Holotype: BR; isotypes: K, NY).
Section: Cremanium.
Distribution: Venezuela, Bolivia.

Miconia micropetala Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 882 (1891). *Acinodendron micropetalum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—ECUADOR. Jameson s.n. (Holotype: US; isotypes: BR, US).

Miconia setinervia Cogn., Bot. Jahrb. Syst. 42: 143 (1909).

Miconia torta Gleason, Phytologia 3: 349 (1950).

Section: Cremanium.

Distribution: Ecuador, Peru, Bolivia.

Miconia miles-morganii J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 184 (1929). Type:—PERU. Macbride 4318 (Holotype: F; isotypes: NY, US).

Section: Cremanium.

Distribution: Peru.

Miconia militis Wurdack, Phytologia 16: 180 (1968). Type:—MEXICO. Rzedowski 160 (Holotype: MICH; isotype: CAS).

Section: Amblyarrhena.

Distribution: Mexico.

Miconia mimica Gleason, Bull. Torrey Bot. Club 52: 451 (1925). Type:—COLOMBIA. Killip 7936 (Holotype: NY; isotypes: GH, US).

Section: Cremanium.

Distribution: Colombia.

Miconia minuta Gleason, Bull. Torrey Bot. Club 52: 450 (1925). Type:—COLOMBIA. Rusby 791 (Holotype: NY).

Section: Cremanium.

Distribution: Colombia, Ecuador.

Miconia minutiflora (Bonpl.) DC., Prodr. 3: 189 (1828).

Basionym: *Melastoma minutiflorum* Bonpl., Monogr. Melast. 1: 50 (1809). *Acinodendron minutiflorum* (Bonpl.) Kuntze, Revis. Gen. Pl. 1: 245 (1891). Type:—FRENCH GUIANA. Bonpland s.n. (Holotype: P).

Cremanium trinitatis Crueg., Linnaea 20: 111 (1847). *Miconia trinitatis* (Crueg.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 246 (1850).

Glossocentrum collinum Crueg., Linnaea 20: 111 (1847). *Miconia glossocentra* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 243 (1850).

Miconia melanodendron Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 162 (1850).

Melastoma pendulum Salzm. ex Griseb., Fl. Brit. W. Indian Is. 257 (1860).

Cremanium paniculatum Mart. ex Triana, Trans. Linn. Soc. London 28: 118 (1871), *pro syn.*

Miconia minutiflora var. *latifolia* Cogn. in Martius, Fl. bras. 14 (4): 385 (1887).

Miconia borealis Gleason, Bull. Torrey Bot. Club 55: 118 (1928).

Section: Glossocentrum.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Cuba, Colombia, Venezuela, Trinidad & Tobago, Suriname, Guyana, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia miocarpa Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 237 (1850). *Acinodendron miocarpum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. Goudot s.n. (Holotype: P; isotype: BR).

Section: Cremanium.

Distribution: Colombia.

Miconia mirabilis (Aubl.) L.O.Williams, Fieldiana, Bot. 29 (10): 574 (1963).

Basionym: *Fothergilla mirabilis* Aubl., Hist. pl. Guiane 1: 441 (1775). *Fothergilla guianensis* Aubl., Hist. pl. Guiane 1: 441 (1775). *Tamonea guianensis* Aubl., Hist. pl. Guiane 1: 441 (1775). *Melastoma tamonea* Sw., Prodr. 70 (1788). *Lieutautia mirabilis* (Aubl.) Buch'oz, Pl. Nouv. Dec. 10 (1779). *Melastoma fothergilla* Desr. in Lamarck, Encycl. Méth. Bot. 4: 45 (1797). *Chitonia fothergilla* (Desr.) D.Don, Mem. Wern. Nat. Hist. Soc. 4: 318 (1823). *Diplochita fothergilla* (Desr.) DC., Prodr. 3: 176 (1828). *Abrophaes mirabilis* (Aubl.) Raf., Sylva Tellur. 99 (1838). *Miconia fothergilla* (Desr.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 119 (1850). *Miconia guianensis* (Aubl.) Cogn., Jahrb. Königl. Bot. Gart. Berlin 4: 280 (1886). *Acinodendron guianense* (Aubl.) Kuntze, Revis. Gen. Pl. 1: 244 (1891). *Tamonea fothergilla* (Desr.) O.F.Cook & G.N.Collins, Contr. U.S. Natl. Herb. 8: 249 (1903). Type:—FRENCH GUIANA. Aublet s.n. (Holotype: BM).

Melastoma compressum Vahl, Eclog. Amer. 3: 23 (1807).

Diplochita florida DC., Prodr. 3: 176 (1828). *Miconia florida* (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850). *Acinodendron floridum* (DC.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Diplochita rosea Macfad., Fl. Jamaica 2: 74 (1850).

Miconia guianensis var. *ovalis* Cogn. in Martius, Fl. bras. 14 (4): 246 (1887).

Miconia guianensis var. *vulgaris* Cogn. in Martius, Fl. bras. 14 (4): 246 (1887).

Section: Tamonea.

Distribution: Mexico, Belize, Nicaragua, Costa Rica, Cuba, Jamaica, Haiti, Dominican Republic, Puerto Rico, Saint Kitts & Nevis, Antigua, Montserrat, Guadeloupe, Dominica, Martinique, Santa Lucia, Grenada, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Peru, Brazil North, Brazil Northeast, Brazil Southeast.

Miconia mituana Wurdack, Phytologia 13: 75 (1966). Type:—COLOMBIA. Maguire 44089 (Holotype: NY; isotypes: COL, NY).

Section: Tamonea.

Distribution: Colombia, Venezuela.

Miconia mocquerysii Wurdack, Phytologia 22: 404 (1972). Type:—VENEZUELA. Mocquerys 986 (Holotype: P; isotype: P).

Section: Miconia.

Distribution: Colombia, Venezuela.

Miconia modica J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 182 (1929). Type:—PERU. Weberbauer 7552 (Holotype: F; isotypes: NY, S).

Section: Amblyarrhena.

Distribution: Peru.

Miconia moensis (Britton) Alain, Contr. Ocas. Mus. Hist. Nat. Colegio "De La Salle" 14: 4 (1955).

Basionym: *Tamonea moensis* Britton, Mem. Torrey Bot. Club 16: 91 (1920). Type:—CUBA. Shafer 8073 (Holotype: A).

Section: Chaenopleura.

Distribution: Cuba.

Miconia molesta Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 854 (1891). *Acinodendron molestum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. Glaziou 17542 (Syntype: BR; isosyntypes: BR, C, K, P, R).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia molinopampana Wurdack, Phytologia 13: 78 (1966). Type:—PERU. Wurdack 1389A (Holotype: US; isotypes: F, GH, MO, NY, UC, USM).

Section: Cremanium.

Distribution: Peru.

Miconia mollicula Triana, Trans. Linn. Soc. London 28: 113 (1871). *Acinodendron molliculum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. Triana s.n. (Holotype: BM)

Section: Miconia.

Distribution: Colombia, Venezuela.

Miconia mollis Triana, Trans. Linn. Soc. London 28: 113 (1871). *Acinodendron molle* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. Spruce 4177 (Holotype: K; isotypes: BM, BR, K, NY, TCD).

Section: Miconia.

Distribution: Peru.

Miconia molybdea Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 185 (1850). *Acinodendron molybdeum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BOLIVIA. *De Castelnau* 40 (Lectotype: P).

Miconia titanea Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 203 (1850). *Miconia molybdea* var. *titanea* (Naudin) Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 827 (1891).

Section: Miconia.

Distribution: Bolivia, Argentina.

Miconia monciona Urb. & Ekman, Ark. Bot. 23A (11): 17 (1931). Type:—DOMINICAN REPUBLIC. *Ekman H12807* (Holotype: S; isotypes: GH, NY, S, US).

Section: Cremanium.

Distribution: Dominican Republic.

Miconia monzonensis Cogn., Bot. Jahrb. Syst. 42: 140 (1909). Type:—PERU. *Weberbauer* 3420 (Holotype: B, destroyed; isotypes: BR, MOL).

Miconia pterogona Gleason, Amer. J. Bot. 19: 746 (1932).

Miconia monzonensis ssp. *cuzcoensis* Wurdack, Phytologia 23: 484 (1972).

Section: Amblyarrhena.

Distribution: Peru.

Miconia moorei Wurdack, Phytologia 48: 242 (1981). Type:—MEXICO. *Moore 2003* (Holotype: BH; isotype: US).

Section: Octomeris.

Distribution: Mexico.

Miconia morii Almeda, Proc. Calif. Acad. Sci., Ser. 4 52: 46 (2000). Type:—PANAMA. *Mori 3773* (Holotype: CAS; isotypes: MO, PMA, WIS).

Section: Tamonea.

Distribution: Panama.

Miconia mornicola A.C.Nicolson, Phytologia 68: 120 (1990). Type:—DOMINICA. *Lack 1* (Holotype: BM; isotypes: CAS, US).

Section: Cremanium.

Distribution: Dominica.

Miconia mourae Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 830 (1891). *Acinodendron mourae* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Moura 132* (Holotype: BR).

Section: Miconia.

Distribution: Brazil Northeast, Brazil Southeast.

Miconia mulleola Wurdack, Acta Bot. Venez. 4: 61 (1970). Type:—COLOMBIA. *Cuatrecasas 12935* (Holotype: US; isotype: COL).

Section: Amblyarrhena.

Distribution: Colombia, Venezuela.

Miconia multiflora Cogn. in Britton, Bull. Torrey Bot. Club 17: 60 (1890). *Acinodendron multiflorum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BOLIVIA. *Rusby 2254* (Holotype: NY; isotypes: BR, CAS, GH, MICH, MIN, PH, US, WIS).

Section: Miconia.

Distribution: Venezuela, Bolivia.

Miconia multiglandulosa Cogn. in Urban, Symb. Antill. 5: 448 (1908). Type:—HAITI. *Buch 691* (Holotype: B, destroyed; isotype: BR); *Ekman H4347* (Neotype: S; isoneotypes: EHH, GH, NY, US).

Section: Chaenopleura.

Distribution: Haiti.

Miconia multinervia Cogn. in Martius, Fl. bras. 14 (4): 259 (1887). *Acinodendron multinerve* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Langsdorff 2150a* (Holotype: LE).

Miconia multinervia var. *minor* Cogn. in Martius, Fl. bras. 14 (4): 260 (1887).

Section: Octomeris.

Distribution: Brazil Southeast.

Miconia multiplinervia Cogn., Bot. Jahrb. Syst. 8: 26 (1886). *Acinodendron multiplinerve* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Lehmann 34* (Holotype: BR; isotype: US).

Miconia alfredii Kraenzl., Vierteljahrsschr. Naturf. Ges. Zürich 76: 155 (1931).

Section: Amblyarrhena.

Distribution: Costa Rica, Panama, Colombia, Ecuador.

Miconia multispicata Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 131 (1850). Type:—JAMAICA. *Purdie s.n.* (Holotype: P; isotype: K).

Cremanium integrifolium Macfad., Fl. Jamaica 2: 93 (1850). *Acinodendron integrifolium* (Macfad.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Miconia perplexans Sprague, Trans. & Proc. Bot. Soc. Edinb. 22: 432 (1905).

Miconia valenciana Kraenzl., Vierteljahrsschr. Naturf. Ges. Zürich 76: 154 (1931).

Section: Miconia.

Distribution: Nicaragua, Costa Rica, Panama, Jamaica, Colombia, Venezuela, Ecuador, Peru, Bolivia, Brazil North.

Miconia muricata (D.Don) Triana, Trans. Linn. Soc. London 28: 102 (1871).

Basionym: *Axinaea muricata* D.Don, Mem. Wern. Nat. Hist. Soc. 4: 321 (1823). *Acinodendron muricatum* (D.Don) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Ruiz & Pavón s.n.* (Holotype: OXF?; isotypes: B, destroyed, BR, FI, G, MA?)

Section: Tamonea.

Distribution: Peru, Bolivia.

Miconia mutabilis (DC.) Triana, Trans. Linn. Soc. London 28: 112 (1871).

Basionym: *Clidemia mutabilis* DC., Prodr. 3: 162 (1828). *Melastoma mutabile* Rich. ex DC., Prodr. 3: 162 (1828), *pro syn.* *Staphidium mutabile* (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 324 (1851). *Acinodendron mutabile* (DC.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—FRENCH GUIANA. *Richard s.n.* (Holotype: P).

Section: Miconia.

Distribution: French Guiana.

Miconia mutisiana Markgr. in Cuatrecasas, Trab. Mus. Nac. Ci. Nat., Ser. Bot. 26: 26 (1933). Type:—COLOMBIA. *Cuatrecasas 2506* (Holotype: MA).

Section: Cremanium.

Distribution: Colombia.

Miconia myriantha Benth., J. Bot. 2: 314 (1840). Type:—GUYANA. *Schomburgk I 507* (Holotype: K; isotypes: BR, E, F, NY, TCD, US).

Section: Glossocentrum.

Distribution: Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Peru, Bolivia, Brazil North.

Miconia myrtiformis Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850). *Cremanium calophyllum* D.Don, Mem. Wern. Nat. Hist. Soc. 4: 313 (1823). *Miconia calophylla* (D.Don) Triana, Trans. Linn. Soc. London 28: 129 (1871), *nom. illeg.*

Acinodendron calophyllum (D.Don) Kuntze, Revis. Gen. Pl. 2: 950 (1891). Type:—PERU. Ruiz & Pavón s.n. (Holotype: OXF?; isotypes: B, destroyed, BR, FI, MA?).

Melastoma obovatum Pav. ex DC., Prodr. 3: 192 (1828), *pro syn.*

Section: Cremanium.

Distribution: Peru.

Miconia myrtillifolia Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 198 (1850). *Acinodendron myrtillifolium* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—VENEZUELA. Linden 459 (Lectotype: P; isolectotypes: BR, GH, NY, P).

Section: Amblyarrhena.

Distribution: Colombia, Venezuela.

Miconia namandensis Wurdack, Phytologia 38: 297 (1978). Type:—ECUADOR. Espinosa E-1077 (Holotype: NY).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia nambyquarae Hoehne, Anexos Mem. Inst. Butantan, Secc. Bot. 1 (5): 131 (1922). Type:—BRAZIL. Kuhlmann 6800a (Holotype: SP; isotype: R).

Section: Miconia.

Distribution: Brazil North, Brazil Central-West.

Miconia napoana Wurdack, Phytologia 31: 494 (1975). Type:—ECUADOR. Lugo 2529 (Holotype: US; isotypes: GB, MO, NY, S).

Section: Tamonea.

Distribution: Colombia, Ecuador, Peru.

Miconia nasella Wurdack, Mem. New York Bot. Gard. 16: 18 (1967). Type:—ECUADOR. Camp E-1332 (Holotype: US; isotypes: COL, NY).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia navioensis Wurdack, Mem. New York Bot. Gard. 10 (4): 38 (1961). Type:—BRAZIL. Cowan, 38118 (Holotype: NY; isotypes: F, RB, S, US).

Section: Miconia.

Distribution: Brazil North.

Miconia neblinensis Wurdack, Mem. New York Bot. Gard. 18 (2): 288 (1969). Type:—BRAZIL. Maguire 60484 (Holotype: NY; isotypes: CAS, NY, US).

Section: Cremanium.

Distribution: Brazil North.

Miconia nematophora Urb. & Ekman, Ark. Bot. 22A (17): 34 (1929). Type:—HAITI. Ekman H6218 (Holotype: S; isotypes: NY, S, US).

Section: Miconia.

Distribution: Haiti, Dominican Republic.

Miconia nerifolia Triana, Trans. Linn. Soc. London 28: 127 (1871). *Acinodendron nerifolium* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. Mathews 24 (Holotype: K; isotypes: BR, K).

Miconia nerifolia var. *brevifolia* Cogn., Bot. Jahrb. Syst. 42: 143 (1909).

Section: Cremanium.

Distribution: Peru.

Miconia nervosa (Sm.) Triana, Trans. Linn. Soc. London 28: 111 (1871).

Basionym: *Melastoma nervosum* Sm. in Rees, Cycl. 23: 31 (1812). *Acinodendron nervosum* (Sm.) Kuntze, Revis. Gen. Pl. 1: 245 (1891). Type:—JAMAICA? (erroneous but exact locality is unknown). *Shakespear s.n.* (Holotype: LINN-HS782-22).

Clidemia spicata Pav. ex D.Don, Mem. Wern. Nat. Hist. Soc. 4: 308 (1823).

Clidemia spicaeformis DC., Prodr. 3: 160 (1828).

Melastoma spicatum Pav. ex DC., Prod. 3: 160 (1828), *pro syn.*

Miconia spondylantha DC., Prodr. 3: 184 (1828). *Melastoma spondylanthum* Mart. ex DC., Prodr. 3: 184 (1828), *pro syn.* *Clidemia spondylantha* Mart. ex Triana, Trans. Linn. Soc. London 28: 111 (1871), *pro syn.* *Tschudya spondylantha* (DC.) Griseb., Fl. Brit. W. Indian Is. 251 (1860).

Miconia fockeana Miq., Comm. Phytogr. 2: 88 (1840).

Miconia pseudonervosa Cogn. in Martius, Fl. bras. 14 (4): 337 pl. 67 (1887). *Acinodendron pseudonervosum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Miconia corallocarpa S.Moore, Trans. Linn. Soc. London, Bot. Ser. 2 4: 363 (1895).

Miconia camposnovaesii Hoehne, Anexos Mem. Inst. Butantan, Secc. Bot. 1 (5): 136 (1922).

Miconia septuplinervis Pittier, J. Wash. Acad. Sci. 14 (19): 447 (1924).

Miconia nervosa var. *mediana* J.F.Macbr., Publ. Field Mus. Nat Hist., Bot. Ser. 4: 181 (1929).

Section: Miconia.

Distribution: Belize, Guatemala, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia neurotricha Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 885 (1891). *Acinodendron neurotrichum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Triana s.n.* (Syntype: BM?; isotypes: BR, G, US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia nigricans Cogn., Bot. Jahrb. Syst. 42: 146 (1909). Type:—PERU. *Weberbauer* 4392 (Holotype: B, destroyed; isotype: BR).

Section: Chaenopleura.

Distribution: Peru.

Miconia nigripes Cogn. & Gleason, Brittonia 1: 119 (1932). Type:—COLOMBIA. *Lehmann* 5474 (Holotype: K; isotypes: BR, CAS, F, K, NY, S, US).

Section: Octomeris.

Distribution: Colombia.

Miconia nitida (D.Don) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850).

Basionym: *Cremanium nitidum* D.Don, Mem. Wern. Nat. Hist. Soc. 4: 312 (1823). *Acinodendron nitidum* (D.Don) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Ruiz & Pavón s.n.* (Holotype: OXF?; isotypes: B, destroyed, BR, FI, G, MA, US).

Miconia macrostoma Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 235 (1850).

Section: Chaenopleura.

Distribution: Peru.

Miconia nitidissima Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 748 (1891). *Acinodendron nitidissimum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—VENEZUELA. *Linden* 454 (Holotype: G-DEL; isotypes: BR, MPU).

Section: Amblyarrhena.

Distribution: Venezuela.

Miconia nobilis Gleason, Bull. Torrey Bot. Club 58: 235 (1931). Type:—PERU. *Killip* 26006 (Holotype: NY; isotype: US).

Section: Miconia.

Distribution: Peru.

Miconia nodosa Cogn., Bull. Acad. Belg., Ser. 3 14: 962 (1887). *Acinodendron nodosum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Jameson* 447 (Syntype: K; isotypes: BR, K, TCD).
Section: Cremanium.
Distribution: Colombia, Ecuador.

Miconia notabilis Triana, Trans. Linn. Soc. London 28: 103 (1871). *Acinodendron notabile* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—COLOMBIA. *Triana* 4021 (Holotype: BM; isotypes: COL, US).
Section: Octomeris.
Distribution: Colombia, Ecuador.

Miconia novemnervia Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 216 (1850). *Acinodendron novemnerve* (Naudin) Kuntze, Revis. Gen. Pl. 1: 245 (1891). Type:—VENEZUELA. *Linden* 290 (Holotype: P; isotypes: BR, G).
Section: Octomeris.
Distribution: Venezuela.

Miconia nubicola Proctor, Bull. Inst. Jamaica, Sci. Ser. 16: 42 (1967). Type:—JAMAICA. *Proctor* 9691 (Holotype: IJ; isotypes: A, BM).
Section: Amblyarrhena.
Distribution: Jamaica

Miconia nutans Donn.Sm., Bot. Gaz. 42: 296 (1906). Type:—COSTA RICA. *Pittier* 11059 (Syntype: US; isosyntype: BR).
Miconia flaviflora Gleason, Bull. Torrey Bot. Club 63: 534 (1936).
Miconia flaviflora var. *emigrans* Gleason, Brittonia 2: 320 (1937).
Section: Tamonea.
Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Ecuador, Peru.

Miconia nystroemii Ekman ex Urb., Symb. Antill. 9: 114 (1923). Type:—CUBA. *Ekman* 5453 (Holotype: S).
Section: Cremanium.
Distribution: Cuba.

Miconia obconica Gleason & Wurdack, Phytologia 16: 172 (1968). Type:—MEXICO. *Hinton* 10361 (Holotype: US; isotypes: DS, GH, K, NY, RSA).
Section: Octomeris.
Distribution: Mexico.

Miconia obliqua Gleason, Amer. J. Bot. 19: 743 (1932). Type:—BOLIVIA. *Buchtien* 1083 (Holotype: US; isotype: US).
Section: Tamonea.
Distribution: Bolivia.

Miconia oblongifolia Cogn. in Martius, Fl. bras. 14 (4): 373 pl. 75 (1887). *Acinodendron oblongifolium* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Glaziou* 13437 (Holotype: BR; isotypes: C, G, K, R).
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia obovata Triana, Trans. Linn. Soc. London 28: 123 (1871). *Acinodendron obovatum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Triana* s.n. (Holotype: BM).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia obscura (Bonpl.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 243 (1850).
Basionym: *Melastoma obscurum* Bonpl., Monogr. Melast. 1: 121 (1816). *Clidemia obscura* (Bonpl.) DC., Prodr. 3: 162 (1828). *Acinodendron obscurum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—ECUADOR. *Bonpland* s.n. (Holotype: P).
Miconia trichrona J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 183 (1929).

Section: Amblyarrhena.

Distribution: Ecuador, Peru.

Miconia obtusa (Griseb.) Triana, Trans. Linn. Soc. London 28: 103 (1871).

Basionym: *Charianthus obtusus* Griseb., Pl. Wright. 189 (1860). *Graffenrieda obtusa* (Griseb.) Griseb., Cat. Pl. Cub. 103 (1866). *Acinodendron obtusum* (Griseb.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). *Miconia confusa* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 759 (1891). *Acinodendron confusum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891). Type:—CUBA. Wright 2527 (Holotype: GOET; isotypes: BM, BR, GH, K, MO, S).

Charianthus obliquus Griseb., Pl. Wright. 186 (1860). *Calycogonium obliquum* (Griseb.) Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 947 (1891).

Miconia praecox C.Wright in Sauvalle, Anales Acad. Ci. Med. Habana 5: 436 (1869). *Tamonea praecox* (C.Wright) Jenn., Ann. Carnegie Mus. 11: 211 (1917).

Section: Octomeris.

Distribution: Cuba.

Miconia ochracea Triana, Trans. Linn. Soc. London 28: 127 (1871). *Acinodendron ochraceum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. Triana 4069 (Holotype: BM; isotypes: COL, NY).

Section: Cremanium.

Distribution: Colombia, Ecuador.

Miconia octopetala Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 754 (1891). *Acinodendron octopetalum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. Glaziou 16967 (Syntype: BR; isotypes: BR, C, K, P, R, US).

Section: Octomeris.

Distribution: Brazil Northeast, Brazil Southeast.

Miconia oellgaardii E.Cotton, Nordic J. Bot. 20: 181 (2000). Type:—ECUADOR. Ollgaard 74326 (Holotype AAU; isotypes: LOJA, MO, NY, QCA).

Section: Cremanium.

Distribution: Ecuador.

Miconia oinochrophylla Donn.Sm., Bot. Gaz. 40: 4 (1905). Type:—GUATEMALA. von Türcckheim 8684 (Holotype: US).

Section: Hartigia.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia.

Miconia oldemanii Wurdack, Phytologia 45: 326 (1980). Type:—FRENCH GUIANA. Oldeman B-637 (Holotype: CAY; isotype: MIN).

Section: Jucunda.

Distribution: French Guiana.

Miconia oligantha Wurdack, Phytologia 26: 5 (1973). Type:—ECUADOR. Lugo 22 (Holotype: S).

Section: Miconia.

Distribution: Ecuador.

Miconia oligocephala Donn.Sm., Bot. Gaz. 46: 111 (1908). Type:—GUATEMALA. von Türcckheim 1781 (Syntype: US; isotypes: BM, BR, E, F, GH, M, NY).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala.

Miconia oligotricha (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).

Basionym: *Cremanium oligotrichum* DC., Prodr. 3: 193 (1828). *Acinodendron oligotrichum* (DC.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—MEXICO. Lagasca s.n. (Holotype: BR).

Melastoma glaucocarpum Schltdl., Linnaea 13 (5): 421 (1839). *Miconia glaucocarpa* (Schltdl.) Steud., Nomencl. Bot., ed. 2 (2): 140 (1841).

Section: Cremanium.

Distribution: Mexico.

Miconia ombrophila Wurdack, Phytologia 35: 250 (1977). Type:—ECUADOR. *Little* 632 (Holotype: US).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia onaensis Wurdack, Mem. New York Bot. Gard. 16: 30 (1967). Type:—ECUADOR. *Prieto P-211* (Holotype: US; isotypes: COL, F, GH, M, NY, RB, U, UC).

Section: Cremanium.

Distribution: Ecuador.

Miconia onychocalyx Gilli, Feddes Report. 92: 682 (1981). Type:—ECUADOR. *Gilli* 252 (Holotype: W).

Section: Glossocentrum.

Distribution: Ecuador.

Miconia opacifolia J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 184 (1929). Type:—PERU. *Macbride 5197* (Holotype: F).

Section: Cremanium.

Distribution: Peru.

Miconia oraria Wurdack, Phytologia 35: 248 (1977). Type:—ECUADOR. *Sparre* 18349 (Holotype: S).

Section: Miconia.

Distribution: Costa Rica, Panama, Colombia, Ecuador.

Miconia orcheotoma Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 227 (1850). *Acinodendron orcheotomum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Linden* 1059 (Holotype: P; isotypes: GH, P).

Section: Cremanium.

Distribution: Colombia, Ecuador.

Miconia oreogena Wurdack, Brittonia 28: 139 (1976). Type:—COLOMBIA. *Barclay* 6684 (Holotype: US; isotypes: COL, MO).

Section: Cremanium.

Distribution: Colombia.

Miconia orescia L.Uribe, Caldasia 9: 461 (1966). Type:—COLOMBIA. *Uribe-Uribe* 5560 (Holotype: COL; isotypes: COL, US).

Section: Cremanium.

Distribution: Colombia, Ecuador.

Miconia organensis Gardner, London J. Bot. 2: 345 (1843). *Acinodendron organense* (Gardner) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Gardner* 390 (Holotype: BM; isotypes: BR, K).

Section: Miconia.

Distribution: Brazil Southeast.

Miconia osaensis Kriebel, Aguilar & Almeda, Proc. Calif. Acad. Sci., Ser. 4 58: 490 (2008). Type:—COSTA RICA. *Aguilar* 5145 (Holotype: INB; isotypes: CAS, CR, MO, USJ).

Section: Amblyarrhena.

Distribution: Costa Rica.

Miconia ossaeifolia Urb. & Ekman, Ark. Bot. 22A (17): 36 (1929). Type:—HAITI. *Ekman* H10350 (Holotype: S; isotype: US).

Section: Glossocentrum.

Distribution: Haiti.

Miconia ottikeri J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 185 (1929). Type:—PERU. *Macbride* 4430 (Holotype: F; isotypes: NY, US).
Section: Chaenopleura.
Distribution: Peru.

Miconia ovalifolia Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 842 (1891). *Acinodendron ovalifolium* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Glaziou* 16908 (Holotype: BR; isotypes: C, F, G, K, MO, P, R, US, Z).
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia ovata Cogn. in Martius, Fl. bras. 14 (4): 281 (1887). *Acinodendron ovatum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Riedel* 395 (Syntypes: BR, LE).
Section: Miconia.
Distribution: Brazil Southeast.

Miconia pachydonta Gleason, Phytologia 1: 41 (1933). Type:—PERU. *Klug* 2143 (Holotype: NY; isotypes: A, F, GH, K, MO, S, US).
Section: Jucunda.
Distribution: Colombia, Peru, Brazil North.

Miconia pachyphylla Cogn., Jahrb. Königl. Bot. Gart. Berlin 4: 279 (1886). *Acinodendron pachyphyllum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PUERTO RICO. *Sintenis* 1334 (Syntype: BR; isosyntypes: GH, US).
Section: Amblyarrhena.
Distribution: Puerto Rico.

Miconia paeminosa Wurdack, Phytologia 23: 479 (1972). Type:—PERU. *Dudley* 13185 (Holotype: US; isotypes: F, NA).
Section: Amblyarrhena.
Distribution: Peru.

Miconia pailasana Wurdack, Mem. New York Bot. Gard. 16: 21 (1967). Type:—ECUADOR. *Camp E-4902* (Holotype: US; isotypes: F, GH, M, MICH, MO, NY, RB, S, VEN).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia paleacea Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 757 (1891). *Acinodendron paleaceum* (Cogn.) Kuntze, Revis. Gen. 2: 952 (1891). Type:—COSTA RICA. *Wendland* s.n. (Holotype: GOET; isotype: BR).
Tococa grandifolia Standl., Publ. Field Mus. Nat. Hist., Bot. Ser. 4 (8): 319 (1929).
Section: Tamonea.
Distribution: Belize, Honduras, Nicaragua, Costa Rica, Colombia, Ecuador, Peru, Brazil North.

Miconia paludigena Wurdack, Phytologia 11: 394 (1965). Type:—PERU. *Wurdack* 1218 (Holotype: US; isotypes: COL, F, K, NY, UC, US, USM).
Section: Cremanium.
Distribution: Ecuador, Peru.

Miconia pandurata Triana, Trans. Linn. Soc. London 28: 108 (1871). *Acinodendron panduratum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Spruce* s.n. (Holotype: K).
Section: Miconia.
Distribution: Colombia, Peru.

Miconia paniculata (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).

Basionym: *Cremanium paniculatum* DC., Prodr. 3: 194 (1828). *Melastoma paniculatum* Mart. & Schrank ex DC., Prodr. 3: 194 (1828), *pro syn.* *Acinodendron paniculatum* (DC.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M).

Melastoma trinervium Vell., Fl. Flum. 4: 178: t. 113. (1825/1829).

Miconia depauperata Gardner, London J. Bot. 2: 346 (1843). *Acinodendron depauperatum* (Gardner) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Cremanium pusillum Cham. ex Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 171 (1850), *pro syn.*

Miconia dorianae Cogn. in Martius, Fl. bras. 14 (4): 376 (1887). *Acinodendron dorianum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Section: Glossocentrum.

Distribution: Brazil Northeast, Brazil Southeast, Brazil South.

Miconia papillosa (Desr.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 216 (1850).

Basionym: *Melastoma papillosum* Desr. in Lamarck, Encycl. Méth. Bot. 4: 48 (1797). *Cremanium papillosum* (Desr.) DC., Prodr. 3: 192 (1828). *Acinodendron papillosum* (Desr.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU or ECUADOR. *de Jussieu s.n.* (Holotype: P-JU).

Miconia quitensis Benth., Pl. Hartw. 182 (1845).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia paradisica Wurdack, Phytologia 11: 391 (1965). Type:—PERU. *Wurdack 1055* (Holotype: US; isotypes: F, GH, NY, USM).

Section: Cremanium.

Distribution: Peru.

Miconia paradoxa (Mart. ex DC.) Triana, Trans. Linn. Soc. London 28: 121 (1871).

Basionym: *Melastoma paradoxum* Mart. ex DC., Prodr. 3: 202 (1828). *Acinodendron paradoxum* (Mart. ex DC.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M).

Section: Amblyarrhena.

Distribution: Brazil Southeast.

Miconia paraguayensis Cogn. in Chodat & Hassler, Bull. Herb. Boissier, Ser. 2 4: 1278 (1904). Type:—PARAGUAY. *Hassler 7695* (Holotype: BR; isotypes: NY, S, UC).

Miconia paraguayensis var. *angustifolia* Cogn., Bull. Herb. Boissier, Ser. 2 4: 1279 (1904).

Section: Miconia.

Distribution: Paraguay.

Miconia parvifolia Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 905 (1891). *Acinodendron parvifolium* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Purdie s.n.* (Holotype: GH; isotype: BR).

Section: Cremanium.

Distribution: Colombia.

Miconia paspaloides Gleason, Bull. Torrey Bot. Club 57: 69 (1930). Type:—COLOMBIA. *Woronow 6436* (Holotype: LE).

Section: Miconia.

Distribution: Colombia.

Miconia pastazana Wurdack, Phytologia 26: 398 (1973). Type:—ECUADOR. *Lugo 1431* (Holotype: GB; isotypes: F, MO, NY, QCA, US).

Section: Glossocentrum.

Distribution: Ecuador.

Miconia pastoensis Triana, Trans. Linn. Soc. London 28: 126 (1871). *Acinodendron pastoense* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Jameson* 415 (Holotype: BM; isotypes: BR, K).
Section: Cremanium.
Distribution: Colombia.

Miconia paucidens DC., Prodr. 3: 186 (1828). *Melastoma paucidens* Schrank & Mart. ex DC., Prodr. 3: 186 (1828), *pro syn.* *Acinodendron paucidens* (DC.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M; isotype: G-DC).
Miconia riedelii Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 125 (1850). *Acinodendron riedelii* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Miconia langsdorffii Cogn. in Martius, Fl. bras. 14 (4): 232 (1887). *Acinodendron langsdorffii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).
Section: Jucunda.
Distribution: Brazil Central-West, Brazil Southeast, Brazil South, Paraguay.

Miconia paupercula (Naudin) Triana, Trans. Linn. Soc. London 28: 102 (1871).
Basionym: *Octomeris paupercula* Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 281 (1851). *Acinodendron pauperculum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—VENEZUELA. *Bonpland s.n.* (Holotype: P).
Section: Octomeris.
Distribution: Venezuela.

Miconia pausana Wurdack, Mem. New York Bot. Gard. 16: 11 (1967). Type:—ECUADOR. *Camp E-1603* (Holotype: US; isotype: NY).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia pavoniana Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 180 (1850). *Acinodendron pavonianum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Pavón s.n.* (Holotype: P; isotypes: BR, G).
Section: Cremanium.
Distribution: Peru.

Miconia pedicellata Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 875 (1891). *Acinodendron pedicellatum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Spruce* 4342 (Syntypes: BM, BR, K, TCD).
Section: Amblyarrhena.
Distribution: Colombia, Peru.

Miconia peltata Almeda, Proc. Calif. Acad. Sci., Ser. 4 46: 217 (1989). Type:—PANAMA. *McPherson* 7882 (Holotype: CAS; isotypes: MEXU, MO, PMA, US).
Section: Miconia.
Distribution: Panama.

Miconia pendula Umaña & Almeda, Novon 3: 7 (1993). Type:—COSTA RICA. *Umaña* 495 (Holotype: CR; isotypes: BM, BR, CAS, COL, CR, F, MEXU, MO, NY, PMA, US, USJ).
Section: Miconia.
Distribution: Costa Rica.

Miconia penduliflora Cogn. in Martius, Fl. bras. 14 (4): 414 (1888). *Acinodendron penduliflorum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Glaziou* 13845 (Syntype: BR; isosyntypes: C, K, P, R).
Section: Amblyarrhena.
Distribution: Brazil Southeast.

Miconia penicillata Gleason, Bull. Torrey Bot. Club 52: 449 (1925). Type:—COLOMBIA. *Pennell* 5808 (Holotype: NY; isotypes: GH, PH).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia penningtonii Wurdack, Phytologia 11: 392 (1965). Type:—ECUADOR. *Cazaleti* 5602 (Holotype: US; isotypes: K, NY, US).
Section: Cremanium.
Distribution: Ecuador.

Miconia pennipilis Cogn. in Martius, Fl. bras. 14 (4): 287 pl. 59 (1887). *Acinodendron pennipile* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Glaziou* 14769 (Syntype: BR; isosyntypes: C, P, R).
Section: Miconia.
Distribution: Brazil Southeast.

Miconia pentlandii Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 182 (1850). *Acinodendron pentlandii* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BOLIVIA. *Pentland* 130 (Holotype: P; isotype: BR).
Section: Miconia.
Distribution: Bolivia.

Miconia pepericarpa DC., Prodr. 3: 182 (1828). *Gallasia pepericarpa* Mart. ex DC., Prodr. 3: 182 (1828), *pro syn.* *Acinodendron pepericarpum* (DC.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Martius* 16 (Holotype: M; isotypes: BR, G-DC).
Miconia claussenii Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 143 (1850).

Miconia pepericarpa var. *grandifolia* Cogn. in Martius, Fl. bras. 14 (4): 360 (1887).

Section: Glossocentrum.
Distribution: Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia perelegans Urb., Repert. Spec. Nov. Regni Veg. 22: 226 (1926). Type:—CUBA. *Ekman* 11918 (Holotype: S; isotype: NY).
Section: Tamonea.
Distribution: Cuba.

Miconia pergamentacea Cogn., Bot. Jahrb. Syst. 8: 24 (1886). *Acinodendron pergamentaceum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Lehmann* 3830 (Holotype: K; isotypes: BR, US).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia perijensis Wurdack, Acta Bot. Venez. 13: 157 (1978). Type:—VENEZUELA. *Wood* 92 (Holotype: US; isotype: VEN).
Section: Cremanium.
Distribution: Colombia, Venezuela.

Miconia pernettifolia Triana, Trans. Linn. Soc. London 28: 127 (1871). *Acinodendron pernettifolium* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA or ECUADOR. *Turner?* s.n. (Holotype: K; isotype: BR).
Section: Cremanium.
Distribution: Ecuador.

Miconia perobscura Wurdack, Mem. New York Bot. Gard. 10 (4): 42 (1961). Type:—VENEZUELA. *Maguire* 36833 (Holotype: NY; isotypes: K, VEN).
Section: Amblyarrhena.
Distribution: Venezuela.

Miconia persicariifolia Cogn. in Britton, Bull. Torrey Bot. Club 17: 59 (1890). *Acinodendron persicariifolium* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BOLIVIA. *Rusby* 992 (Holotype: NY; isotypes: K, US).
Section: Miconia.
Distribution: Peru, Bolivia.

Miconia perturbata Wurdack, Mem. New York Bot. Gard. 10 (5): 172 (1964). Type:—VENEZUELA. *Maguire* 42474 (Holotype: US; isotypes: B, F, GH, K, MICH, MO, NY, RB, S, SP, U, UC, VEN).

Section: Glossocentrum.

Distribution: Venezuela.

Miconia peruviana Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 909 (1891). *Acinodendron peruvianum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. Lechler 2066 (Holotype: BR; isotypes: GOET, K).

Acinodendron peruvianum var. *cogniauxii* Kuntze, Revis. Gen. Pl. 3 (2): 93 (1898).

Section: Cremanium.

Distribution: Peru.

Miconia petersonii Urb., Symb. Antill. 9: 115 (1923). Type:—CUBA. Ekman 6852 (Holotype: S; isotypes: NY, S).

Section: Cremanium.

Distribution: Cuba.

Miconia petroniana Cogn. & Saldanha in Saldanha & Cogniaux, Bouquet Melast. Bras. 3 (1887). *Acinodendron petronianum* (Cogn. & Saldanha) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. Glaziou 16011 (Holotype: BR; isotypes: BR, C, K, P, R).

Miconia pseudopetroniana Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 846 (1891). *Acinodendron pseudopetronianum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia petropolitana Cogn. in Martius, Fl. bras. 14 (4): 613 (1888). *Acinodendron petropolitanum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. Glaziou 16021 (Lectotype: BR; isolectotypes: B, K, P).

Miconia petropolitana var. *macrophylla* Hoehne, Anexos Mem. Inst. Butantan, Secc. Bot. 1 (5): 145 (1922).

Miconia inconspecta var. *glabrata* Wurdack, Sellowia 14: 166 (1962).

Section: Glossocentrum.

Distribution: Brazil Southeast, Brazil South, Argentina.

Miconia phaeochaeta Wurdack, Phytologia 52: 62 (1982). Type:—ECUADOR. Harling 16581 (Holotype: GB; isotypes: MO, S, US).

Section: Tamonea.

Distribution: Colombia, Ecuador.

Miconia phaeophylla Triana, Trans. Linn. Soc. London 28: 113 (1871). *Acinodendron phaeophyllum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. Spruce 4861 (Holotype: K; isotypes: BM, BR, E, NY, P).

Miconia fendleriana Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 822 (1891). *Acinodendron fendlerianum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Section: Miconia.

Distribution: Panama, Colombia, Venezuela, Guyana, Suriname, French Guiana, Peru, Brazil North.

Miconia phaeotricha Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 193 (1850). *Acinodendron phaeotrichum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—MEXICO. Galeotti 2951 (Holotype: P; isotypes: BR, G, P, US).

Section: Amblyarrhena.

Distribution: Mexico.

Miconia phanerostila Pilg., Verh. Bot. Vereins Prov. Brandenburg 47: 173 (1905). Type:—BRAZIL. Ule 5659 (Holotype: B, destroyed; isotypes: HBG, K).

Section: Miconia.

Distribution: Colombia, Peru, Bolivia, Brazil North.

Miconia phlebodes Wurdack, Phytologia 9: 419 (1964). Type:—PERU. Wurdack 1965 (Holotype: US; isotypes: F, K, NY, S, UC, USM).

Section: Miconia.

Distribution: Ecuador, Peru.

Miconia pichinchensis Benth., Pl. Hartw. 182 (1845). *Acinodendron pichinchense* (Benth.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—ECUADOR. *Hartweg 1012* (Holotype: K; isotypes: BM, BR, P).
Section: Amblyarrhena.
Distribution: Colombia, Ecuador.

Miconia picinguabensis R.Goldenb. & A.B.Martins, Kew Bull. 54: 466 (1999). Type:—BRAZIL. *Pedroni 628* (Holotype: UEC; isotypes: MBM, K, RB, SP, US).
Section: Chaenanthera.
Distribution: Brazil Southeast.

Miconia pilaloensis Wurdack, Phytologia 52: 64 (1982). Type:—ECUADOR. *Lojnant 13947* (Holotype: AAU; isotypes: GB, QCA).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia piperifolia Triana, Trans. Linn. Soc. London 28: 109 (1871). *Acinodendron piperifolium* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Triana 4002* (Holotype: BM; isotypes: BR, COL, E, K, NY).
Section: Miconia.
Distribution: Colombia.

Miconia pisinna Wurdack, Acta Bot. Venez. 13: 153 (1978). Type:—VENEZUELA. *Wurdack 2794* (Holotype: US; isotype: NY).
Section: Amblyarrhena.
Distribution: Venezuela.

Miconia pisinniflora Wurdack, Phytologia 29: 147 (1974). Type:—ECUADOR. *Asplund 18591* (Holotype: US; isotypes: B, NY, S).
Section: Chaenanthera.
Distribution: Ecuador, Peru.

Miconia pittieri Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 1191 (1891). *Acinodendron pittieri* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COSTA RICA. *Pittier 2005* (Holotype: BR; isotypes: BR, CAS, P).
Section: Cremanium.
Distribution: Costa Rica, Panama.

Miconia platypoda Gleason, Bull. Torrey Bot. Club 72: 476 (1945). Type:—COLOMBIA. *Cuatrecasas 7185* (Holotype: NY; isotype: COL).
Section: Miconia.
Distribution: Colombia, Guyana, Suriname, Brazil North.

Miconia plena Gleason, Phytologia 1: 31 (1933). Type:—COLOMBIA. *Lawrance 152* (Holotype: NY; isotypes: A, F, K, MICH, MO, US, WIS).
Section: Cremanium.
Distribution: Colombia.

Miconia plethorica Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 227 (1850). *Acinodendron plethoricum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Bonpland s.n.* (Holotype: P; isotype: BR).
Section: Cremanium.
Distribution: Colombia.

Miconia pluknetii Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 116 (1850). *Acinodendron pluknetii* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—GUYANA. *Schomburgk II 1008* (Lectotype: P; isolectotypes: BM, K).
Diplochita sessilifolia Griseb., Fl. Brit. W. Indian Is. 252 (1860).
Section: Adenodesma.

Distribution: Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana.

Miconia plumifera Triana, Trans. Linn. Soc. London 28: 124 (1871). *Acinodendron plumiferum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU or BOLIVIA. *Pearce s.n.* (Holotype: K; isotype: K).

Miconia plumifera var. *bangii* Cogn., Bull. Torrey Bot. Club 23: 16 (1896).

Section: Amblyarrhena.

Distribution: Peru, Bolivia.

Miconia poecilantha L.Uribe, Caldasia 9: 86 (1964). Type:—COLOMBIA. *Uribe-Uribé 3791* (Holotype: COL; isotypes: COL, US).

Section: Octomeris.

Distribution: Colombia.

Miconia poeppigii Triana, Trans. Linn. Soc. London 28: 107 (1871). *Oxymeris furfuracea* Poepp. ex Triana, Trans. Linn. Soc. London 28: 107 (1871), *pro syn.* *Acinodendron poeppigii* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Poeppig 1912* (Syntypes: BR, BM, NY).

Miconia congesta Cogn., Repert. Spec. Nov. Regni Veg. 8: 2 (1910).

Miconia darienensis Pittier, Contr. U.S. Natl. Herb. 18: 248 (1917).

Miconia surinamensis Gleason, Recueil Trav. Bot. Neerl. 32: 212 (1935).

Section: Miconia.

Distribution: Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Northeast.

Miconia polita Gleason, Bull. Torrey Bot. Club 52: 379 (1925). Type:—GUYANA. *Gleason 53* (Holotype: NY).

Section: Miconia.

Distribution: Venezuela, Guyana.

Miconia polyandra Gardner, London J. Bot. 2: 346 (1843). *Acinodendron polyandrum* (Gardner) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Gardner 395* (Holotype: BM; isotypes: BM, FI, GH, K, NY).

Section: Miconia.

Distribution: Brazil Southeast.

Miconia polychaeta Wurdack, Phytologia 23: 493 (1972). Type:—PERU. *Dudley 10430* (Holotype: US)

Section: Cremanium.

Distribution: Peru.

Miconia polygama Cogn. in Britton, Bull. Torrey Bot. Club 17: 92 (1890). *Acinodendron polygamum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BOLIVIA. *Rusby 2519* (Syntype: NY; isosyntype: BR).

Section: Cremanium.

Distribution: Bolivia.

Miconia polyneura Triana, Trans. Linn. Soc. London 28: 128 (1871). *Acinodendron polyneurum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Triana 4068* (Holotype: BM; isotypes: BR, K, NY).

Section: Cremanium.

Distribution: Colombia, Peru.

Miconia polytopica Wurdack, Phytologia 23: 477 (1972). Type:—PERU. *Dudley 10346* (Holotype: US; isotype: MO).

Miconia polytopica ssp. *ayacuchensis* Wurdack, Phytologia 23: 478 (1972).

Miconia polytopica ssp. *huanucensis* Wurdack, Phytologia 23: 479 (1972).

Section: Amblyarrhena.

Distribution: Peru, Bolivia.

Miconia poortmannii (Cogn.) Wurdack, Phytologia 35: 250 (1977).

Basionym: *Tococa poortmannii* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 971 (1891). Type:—
ECUADOR. Poortmann 90 (Holotype: P; isotype: BR).

Miconia espinosana Gleason, Phytologia 3: 28 (1948).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia popayanensis Wurdack, Phytologia 41: 6 (1978). Type:—COLOMBIA. Lehmann 5473 (Holotype: K).

Section: Chaenopleura.

Distribution: Colombia.

Miconia porphyrotricha (Markgr.) Wurdack, Phytologia 22: 404 (1972).

Basionym: *Heterotrichum porphyrotrichum* Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 12: 181 (1934). Type:—
ECUADOR. Schimpff 1077 (Holotype: B, destroyed).

Miconia skutchii Gleason, Bull. Torrey Bot. Club 68: 250 (1941).

Section: Octomeris.

Distribution: Colombia, Ecuador.

Miconia pozensis Wurdack, Phytologia 5: 58 (1954). Type:—COLOMBIA. Sandeman 6114 (Holotype: K; isotypes:
COL, NY, US).

Section: Miconia.

Distribution: Colombia.

Miconia prancei Wurdack, Phytologia 24: 199 (1972). Type:—BRAZIL. Prance 14793 (Holotype: US; isotypes: F, GH,
K, M, NY, P, S, U).

Section: Adenodesma.

Distribution: Peru, Brazil North.

Miconia prasina (Sw.) DC., Prodr. 3: 188 (1828).

Basionym: *Melastoma prasinum* Sw., Prodr. 69 (1788). *Acinodendron prasinum* (Sw.) Kuntze, Revis. Gen. Pl. 1: 245
(1891). Type:—JAMAICA. Swartz s.n. (Lectotype: S).

Melastoma laevigatum Aubl., Hist. pl. Guiane 1: 412 (1775).

Melastoma parviflorum Aubl., Hist. pl. Guiane 1: 433 (1775). *Miconia parviflora* (Aubl.) Cogn. in de Candolle & de
Candolle, Monogr. Phan. 7: 805 (1891).

Melastoma pendulifolium Rich., Actes Soc. Hist. Nat. Paris 1: 109 (1792).

Melastoma montanum Spreng., Syst. Veg. 2: 296 (1825).

Conostegia parviflora DC., Prodr. 3: 175 (1828).

Miconia collina DC., Prodr. 3: 185 (1828). *Miconia prasina* var. *collina* (DC.) Triana, Trans. Linn. Soc. London 28: 109
(1871).

Miconia sepiaria DC., Prodr. 3: 185 (1828). *Melastoma sepiarium* Mart. ex DC., Prodr. 3: 185 (1828), *pro syn.*

Miconia attenuata DC., Prodr. 3: 186 (1828). *Miconia prasina* var. *attenuata* (DC.) Cogn. in Martius, Fl. bras. 14 (4):
317 (1887).

Miconia attenuata var. *subquintuplinervia* DC., Prodr. 3: 187 (1828).

Melastoma pulchrum Vell., Fl. Flum. 4: 178: t. 110 (1825/1829).

Melastoma suaveolens Raddi, Mem. Soc. Ital. Modena 20: 133 (1829).

Miconia revoluta Benth., J. Bot. 2: 313 (1840).

Miconia pteropoda Benth., J. Bot. 2: 314 (1840). *Acinodendron pteropodium* (Benth.) Kuntze, Revis. Gen. Pl. 2: 952
(1891).

Miconia aubletiana Steud., Nomencl. Bot., ed. 2 (2): 139 (1841).

Miconia fleischeriana Steud., Flora 27: 723 (1844).

Miconia macrophylla Steud., Flora 27: 723 (1844).

Miconia repandocrenata Steud., Flora 27: 724 (1844).

Miconia crispula Spruce ex Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 172 (1850), *pro syn.* *Miconia prasina* var. *crispula*
(Spruce ex Naudin) Cogn. in Martius, Fl. bras. 14 (4): 317 (1887).

Miconia cristulata Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 177 (1850).

Miconia nemoralis Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 178 (1850).

Miconia palustris Macfad., Fl. Jamaica 2: 89 (1850).

Miconia prasina var. *denticulata* Griseb., Bonplandia 6: 6 (1858).

Melastoma trinervium Salzm. ex Griseb., Fl. Brit. W. Indian Is. 257 (1860).

Melastoma quinquenervium Salzm. ex Griseb., Fl. Brit. W. Indian Is. 297 (1860).

Miconia trichotoma Bello, Anales Soc. Esp. Hist. Nat. 10: 268 (1883).

Miconia prasina var. *angustifolia* Cogn. in Martius, Fl. bras. 14 (4): 317 (1887).

Melastoma acuminatum Sessé & Moc., Fl. mexic., ed. 2: 105 (1894).

Melastoma leucanthum Sessé & Moc., Fl. mexic., ed. 2: 105 (1894).

Melastoma glabrum Sessé & Moc., Fl. mexic., ed. 2: 107 (1894).

Miconia mucronulata Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 358 (1915).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Cuba, Jamaica, Haiti, Dominican Republic, Puerto Rico, Guadeloupe, Dominica, Martinique, Santa Lucia, Grenada, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast, Brazil South, Paraguay.

Miconia prasinifolia Gleason, Phytologia 3: 348 (1950). Type:—COLOMBIA. Killip 9011 (Holotype: NY; isotype: US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia prietoi Wurdack, Mem. New York Bot. Gard. 16: 31 (1967). Type:—ECUADOR. Prieto P-188 (Holotype: US; isotypes: F, GH, MO, NY).

Section: Cremanium.

Distribution: Ecuador.

Miconia procumbens (Gleason) Wurdack, Phytologia 35: 246 (1977).

Basionym: *Clidemia procumbens* Gleason, Bull. Torrey Bot. Club 58: 255 (1931). Type:—PERU. Killip 28852 (Holotype: NY; isotypes: K, US).

Section: Miconia.

Distribution: Panama, Ecuador, Peru, Brazil North.

Miconia prominens Wurdack, Phytologia 38: 291 (1978). Type:—ECUADOR. Camp E-791 (Holotype: NY; isotype: US).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia protuberans Wurdack, Phytologia 52: 65 (1982). Type:—ECUADOR. Harling 16443 (Holotype: GB; isotype: US).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia pseudoaplostachya Cogn. in Martius, Fl. bras. 14 (4): 263 (1887). *Acinodendron pseudoaplostachyum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—VENEZUELA. Spruce 3324 (Holotype: BR; isotypes: BM, GOET, K).

Section: Miconia.

Distribution: Venezuela, Guyana, French Guiana.

Miconia pseudocapsularis Wurdack, Mem. New York Bot. Gard. 10 (4): 32 (1961). Type:—VENEZUELA. Maguire 42015 (Holotype: NY; isotypes: US, VEN).

Section: Miconia.

Distribution: Venezuela, Brazil North.

Miconia pseudocentrophora Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 893 (1891). *Acinodendron pseudocentrophorum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—ECUADOR. *Spruce 5134* (Syntypes: G-BOISS, GOET, K, LE, P; isosyntypes: BM, BR, NY, S, TCD).

Section: Amblyarrhena.

Distribution: Ecuador, Peru.

Miconia pseudoeichleri Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 844 (1891). *Acinodendron pseudoeichleri* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Glaziou 16935* (Holotype: BR; isotypes: BR, CAS, K, LIL, P, R, US).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia pseudoradula Cogn. & Gleason, Brittonia 1: 120 (1932). Type:—COLOMBIA. *Lehmann 5952* (Holotype: K; isotype: BR)

Section: Amblyarrhena.

Distribution: Colombia.

Miconia pseudorigida Proctor, Bull. Inst. Jamaica, Sci. Ser. 16: 42 (1967). Type:—JAMAICA. *Proctor 10469* (Holotype: IJ).

Section: Chaenopleura.

Distribution: Jamaica.

Miconia psychrophila Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 220 (1850). *Acinodendron psychrophilum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Goudot s.n.* (Holotype: P).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia pterocaulon Triana, Trans. Linn. Soc. London 28: 114 (1871). *Acinodendron pterocaulon* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Triana 3988* (Holotype: K; isotypes: COL, P).

Section: Miconia.

Distribution: Costa Rica, Panama, Colombia, Venezuela, Ecuador, Peru, Brazil North.

Miconia pteroclada Urb., Repert. Spec. Nov. Regni Veg. 22: 229 (1926). Type:—CUBA. *Ekman 14410* (Holotype: S; isotypes: NY, US).

Section: Chaenopleura.

Distribution: Cuba.

Miconia puberula Cogn. in Martius, Fl. bras. 14 (4): 386 (1887). *Acinodendron puberulum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Spruce s.n.* (Syntype: BR; isosyntypes: K, G, NY, P).

Section: Glossocentrum.

Distribution: Colombia, Peru, Bolivia, Brazil North.

Miconia pubicalyx Gleason, Amer. J. Bot. 19: 742 (1932). Type:—BOLIVIA. *Buchtien 999a* (Holotype: NY).

Section: Miconia.

Distribution: Bolivia.

Miconia pubipetala Miq., Stirp. Surinam. Select. 50 (1850). Type:—SURINAME. *Hostmann 896* (Holotype: U; isotype: K). *Diplochita parviflora* Benth., J. Bot. 2: 302 (1840). *Miconia parviflora* (Benth.) Cogn. in Martius, Fl. bras. 14 (4): 249 (1887). *Acinodendron parviflorum* (Benth.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Miconia verticilliflora Steud. ex Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 122 (1850).

Section: Tamonea.

Distribution: Colombia, Venezuela, Trinidad & Tobago, Guyana, French Guiana, Suriname, Peru, Brazil North.

Miconia pujana Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 14: 34 (1938). Type:—ECUADOR. *Schultze-Rhonhof 1854* (Holotype: B, destroyed).

Section: Laceraria.

Distribution: Ecuador, Peru.

Miconia pulchra Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 795 (1891). *Acinodendron pulchrum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Glaziou* 17541 (Holotype: BR; isotype: P).

Section: Miconia.

Distribution: Brazil Southeast.

Miconia pulgari J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 182 (1929). Type:—PERU. *Macbride* 4145 (Holotype: F; isotypes: NY, US).

Section: Amblyarrhena.

Distribution: Peru.

Miconia pulverulenta Ruiz & Pav., Syst. Veg. Fl. Peruv. Chil. 1: 104 (1798). *Cremanium pulverulentum* D.Don ex Triana, Trans. Linn. Soc. London 28: 126 (1871), *pro syn. Acinodendron pulverulentum* (Ruiz & Pav.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Ruiz & Pav. s.n.* (Syntypes: B, destroyed, BR, FI, G-DC).

Miconia pulverulenta var. *trifasciata* Linden & André, Ill. Hort. 20: 197 (1873).

Miconia ramosipila J.F.Macbr., Publ. Field Mus. Nat Hist., Bot. Ser. 4: 185 (1929).

Section: Cremanium.

Distribution: Peru.

Miconia pulvinata Gleason, Bull. Torrey Bot. Club 52: 386 (1925). Type:—COLOMBIA. *Smith* 1841 (Holotype: NY; isotypes: F, GH, K, US).

Section: Hartigia.

Distribution: Colombia, Venezuela.

Miconia punctata (Desr.) D.Don, Mem. Wern. Nat. Hist. Soc. 4: 316 (1823).

Basionym: *Melastoma punctatum* Desr. in Lamarck, Encycl. Méth. Bot. 4: 50 (1797). *Eurychaenia punctata* (Desr.) Griseb., Fl. Brit. W. Indian Is. 259 (1860). *Acinodendron punctatum* (Desr.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—DOMINICAN REPUBLIC. *Herb. Delétang* 59 (Holotype: P-JU).

Miconia habrolepis Standl., Publ. Field Mus. Nat. Hist., Bot. Ser. 4 (8): 319 (1929).

Miconia discolor Macfad., Fl. Jamaica 2: 85 (1850).

Eurychaenia punctata var. *latifolia* Griseb., Fl. Brit. W. Indian Is. 259 (1860). *Miconia punctata* var. *latifolia* (Griseb.) Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 794 (1891).

Miconia lepidota var. *grandifolia* Cogn. in Martius, Fl. bras. 14 (4): 293 (1887).

Miconia punctata var. *brevifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 794 (1891).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Cuba, Dominican Republic, Puerto Rico, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West.

Miconia punctibullata M.E.Morales-P., Michelangeli & F.Gonzalez, Brittonia 60: 229 (2008). Type:—COLOMBIA. *Morales* 1884 (Holotype: COL; isotypes: COL, NY).

Section: Cremanium.

Distribution: Colombia.

Miconia punicea Wurdack, Phytologia 23: 480 (1972). Type:—PERU. *Dudley* 10663 (Holotype: US).

Section: Amblyarrhena.

Distribution: Peru.

Miconia puricensis Wurdack, Phytologia 8: 170 (1962). Type:—COLOMBIA. *Cuatrecasas* 23449 (Holotype: US; isotypes: F, U).

Section: Cremanium.

Distribution: Colombia.

Miconia purulensis Donn.Sm., Bot. Gaz. 46: 111 (1908). Type:—GUATEMALA. von Türcckheim 1718 (Holotype: US; isotypes: BR, E, F, MO, NY).
Section: Cremanium.
Distribution: Mexico, Guatemala, Honduras.

Miconia pusilliflora (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 171 (1850).
Basionym: *Cremanium pusilliflorum* DC., Prodr. 3: 194 (1828). *Melastoma pusilliflorum* Mart. & Schrank ex DC., Prodr. 3: 194 (1828), *pro syn. Acinodendron pusilliflorum* (DC.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. Martius s.n. (Holotype: M; isotype: G-DC).
Melastoma hymenonervium Raddi, Mem. Soc. Ital. Modena 20: 135 (1829). *Miconia hymenonervia* (Raddi) Cogn. in Martius, Fl. bras. 14 (4): 399 (1888). *Acinodendron hymenonerve* (Raddi) Kuntze, Revis. Gen. Pl. 2: 951 (1891).
Melastoma subrepandum Vell., Fl. Flum. 4: 178 (1825/1829).
Chaenopleura parviflora Gardner, Hooker's J. Bot. Kew Gard. Misc. 2: 348 (1843). *Miconia parviflora* (Gardner) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).
Chaenopleura densiflora Gardner, Hooker's J. Bot. Kew Gard. Misc. 2: 349 (1843). *Miconia densiflora* (Gardner) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850). *Acinodendron densiflorum* (Gardner) Kuntze, Revis. Gen. Pl. 2: 951 (1891).
Chaenopleura lanceolata Gardner, Hooker's J. Bot. Kew Gard. Misc. 2: 349 (1843). *Miconia gardneriana* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850).
Miconia rigidiuscula Cogn. in Martius, Fl. bras. 14 (4): 398 (1888). *Acinodendron rigidiusculum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).
Miconia rigidiuscula var. *parvifolia* Cogn. in Martius, Fl. bras. 14 (4): 399 (1888).
Miconia rigidiuscula var. *purpurascens* Cogn. in Martius, Fl. bras. 14 (4): 399 (1888).
Miconia pusilliflora var. *intermedia* Cogn. in Martius, Fl. bras. 14 (4): 403 (1888).
Miconia pusilliflora var. *major* Cogn. in Martius, Fl. bras. 14 (4): 403 (1888).
Miconia rimalis var. *brevifolia* Cogn. in Martius, Fl. bras. 14 (4): 404 (1888).
Section: Chaenanthera.
Distribution: Brazil Southeast, Brazil South, Paraguay, Argentina.

Miconia pustulata Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 208 (1850). *Acinodendron pustulatum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. Linden 1069 (Holotype: P; isotypes: BR, G, GH, K).
Section: Amblyarrhena.
Distribution: Colombia, Ecuador.

Miconia pycnoneura Urb., Repert. Spec. Nov. Regni Veg. 17: 162 (1921). Type:—PUERTO RICO. Judd 6138 (Neotype: FLAS; isoneotypes: MO, NY, S).
Miconia grisebachii var. *reticulata* Cogn., Jahrb. Königl. Bot. Gart. Berlin 4: 280 (1886).
Section: Chaenopleura.
Distribution: Puerto Rico.

Miconia pyramidalis (Desr.) DC., Prodr. 3: 188 (1828).
Basionym: *Melastoma pyramidalis* Desr. in Lamarck, Encycl. Meth. Bot. 4: 53 (1797). Type:—"ANTILLES". Badier (Holotype: P?)
Miconia pyramidalis var. *ciliolata* DC., Prodr. 3: 188 (1828).
Miconia pyramidalis var. *sphaerocarpa* DC., Prodr. 3: 188 (1828).
Section: Miconia.
Distribution: Jamaica, Haiti, Dominican Republic, Puerto Rico.

Miconia pyrifolia Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 164 (1850). *Acinodendron pyrifolium* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. Blanchet 3412 (Holotype: P; isotypes: BR, C, F, K, NY, US).
Section: Miconia.
Distribution: Dominican Republic, Venezuela, Guyana, Suriname, French Guiana, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia quadrangularis (Sw.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 197 (1850).

Basionym: *Melastoma quadrangulare* Sw., Prodr. 69 (1788). *Cremanium quadrangulare* (Sw.) DC., Prodr. 3: 192 (1828). *Chaenopleura quadrangularis* (Sw.) Macfad., Fl. Jamaica 2: 96 (1850). *Pleurochaenia quadrangularis* (Sw.) Griseb., Abh. Konigl. Ges. Wiss. Göttingen 9: 55 (1861). *Acinodendron quadrangulare* (Sw.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—JAMAICA. *Swartz s.n.* (Holotype: S; isotypes: BM, LD, SBT).

Miconia quadrangularis f. *latifolia* Cogn., Jahrb. Königl. Bot. Gart. Berlin 4: 281 (1886).

Miconia quadrangularis f. *minor* Cogn., Jahrb. Königl. Bot. Gart. Berlin 4: 281 (1886).

Miconia quadrangularis var. *glandulosa* Proctor, Bull. Inst. Jamaica, Sci. Ser. 16: 45 (1967).

Section: Chaenopleura.

Distribution: Jamaica.

Miconia quadrialata Renner & Beck, Novon 13: 110 (2003). Type:—BOLIVIA. *Beck 25988* (Holotype: LPB; isotypes: AAU, CAS, MO, US).

Section: Cremanium.

Distribution: Bolivia.

Miconia quadrifolia Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 223 (1850). *Acinodendron quadrifolium* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Mathews 1724* (Holotype: P; isotypes: BM, BR, E, K, NY).

Section: Cremanium.

Distribution: Peru.

Miconia quadripora Wurdack, Phytologia 39: 329 (1978). Type:—ECUADOR. *Harling 3800* (Holotype: S).

Section: Cremanium.

Distribution: Ecuador, Peru.

Miconia quintuplinervia Cogn., Bot. Jahrb. Syst. 8: 26 (1886). *Acinodendron quintuplinerve* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Lehmann 32* (Holotype: K; isotypes: BR, US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia rabenii Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 832 (1891). *Acinodendron rabenii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Glaziou 17561a* (Syntype: BR; isosyntype: P).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia racemifera (DC.) Triana, Trans. Linn. Soc. London 28: 119 (1871).

Basionym: *Leandra racemifera* DC., Prodr. 3: 155 (1828). *Melastoma racemiferum* Schrank & Mart. ex DC., Prodr. 3: 155 (1828), pro syn. *Sagraea racemifera* (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 18: 91, 102 (1852). *Acinodendron racemiferum* (DC.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M).

Miconia racemifera var. *diffusa* Cogn. in Martius, Fl. bras. 14 (4): 367 (1887).

Miconia eriantha Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 845 (1891). *Acinodendron erianthum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Section: Glossocentrum.

Distribution: Brazil Southeast, Brazil South.

Miconia racemosa (Aubl.) DC., Prodr. 3: 179 (1828).

Basionym: *Melastoma racemosum* Aubl., Hist. pl. Guiane 1: 406 (1775). *Tamonea racemosa* (Aubl.) O.F.Cook & G.N.Collins, Contr. U.S. Natl. Herb. 8: 249 (1903). *Acinodendron racemosum* (Aubl.) Kuntze, Revis. Gen. Pl. 1: 245 (1891). Type:—FRENCH GUIANA. *Aublet s.n.* (Holotype: BM).

Melastoma croceum Spreng., Syst. Veg. 2: 302 (1825).

Miconia brachypoda DC., Prodr. 3: 180 (1828). *Miconia racemosa* var. *brachypoda* (DC.) Cogn. in Martius, Fl. bras. 14 (4): 411 (1888).

Hartigia oblongifolia Miq., Linnaea 18: 288 (1844).

Hartigia spectabilis Miq., Linnaea 18: 286 (1844).

Miconia racemosa var. *lanceolata* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 154 (1850).

Miconia circumsecta Macfad., Fl. Jamaica 2: 77 (1850).

Miconia verticillata Macfad., Fl. Jamaica 2: 78 (1850).

Miconia serrulata Macfad. ex Griseb., Fl. Brit. W. Indian Is. 258 (1860).

Miconia verruculosa Steud. ex Triana, Trans. Linn. Soc. London 28: 120 (1871), *pro syn.*

Melastoma venosum Sessé & Moc., Fl. mexic., ed. 2: 106 (1894).

Section: Hartigia.

Distribution: Panama, Jamaica, Haiti, Dominican Republic, Puerto Rico, Dominica, Santa Lucia, Grenada, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Brazil North, Brazil Northeast.

Miconia radula Cogn., Bull. Acad. Roy. Sci. Belgique, Cl. Sci. 14: 957 (1887). *Acinodendron radulum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—ECUADOR. André 3880 p.p. (Holotype: K; isotypes: BR, NY).

Miconia sphagnophila J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 183 (1929).

Section: Amblyarrhena.

Distribution: Ecuador, Peru.

Miconia radulifolia (Benth.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 243 (1850).

Basionym: *Clidemia radulaefolia* Benth., J. Bot. 2: 309 (1840). *Acinodendron radulifolium* (Benth.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—GUYANA. Schomburgk II 998 (Holotype: K; isotypes: BR).

Miconia scrobiculata Cogn. in Martius, Fl. bras. 14 (4): 334: pl. 66 (1887). *Acinodendron scrobiculatum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Section: Miconia.

Distribution: Colombia, Venezuela, Guyana, Ecuador, Peru, Brazil North.

Miconia ramboi Brade, Sellowia 8: 376 (1957). Type:—BRAZIL. Rambo 55924 (Syntype: PACA; isosyntype: RB).

Section: Amblyarrhena.

Distribution: Brazil South.

Miconia rava Wurdack, Phytologia 5: 129 (1955). Type:—COLOMBIA. Cuatrecasas 21705 (Holotype: NY; isotypes: F, S, U, US).

Section: Cremanium.

Distribution: Colombia.

Miconia ravenii Wurdack, Phytologia 14: 269 (1967). Type:—MEXICO. Breedlove 12928 (Holotype: US; isotypes: CAS, F, LL).

Section: Octomeris.

Distribution: Mexico.

Miconia reburrosa Wurdack, Mem. New York Bot. Gard. 16: 23 (1967). Type:—ECUADOR. Camp E-1604 (Holotype: NY).

Section: Cremanium.

Distribution: Ecuador.

Miconia reclinata (Bonpl.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 209 (1850).

Basionym: *Melastoma reclinatum* Bonpl., Monogr. Melast. 1: 112 (1815). *Cremanium reclinatum* (Bonpl.) DC., Prodr. 3: 192 (1828). *Acinodendron reclinatum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). *Tamonea reclinata* (Bonpl.) Krasser in Engler & Prantl, Nat. Pflanzenfam. 3 (7): 142 (1893). Type:—COLOMBIA. Bonpland s.n. (Holotype: P).

Section: Cremanium.

Distribution: Colombia.

Miconia recondita Wurdack, Phytologia 23: 488 (1972). Type:—BOLIVIA. Bang 2485 (Holotype: US; isotype: US).

Section: Cremanium.

Distribution: Bolivia.

Miconia reducens Triana, Trans. Linn. Soc. London 28: 106 (1871). *Acinodendron reducens* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Triana* 4011 (Syntype: BM; isotypes: COL, K, US).

Miconia curvipetiolata Cogn. & Gleason, Brittonia 1: 113 (1932).

Miconia foliosa Triana, Trans. Linn. Soc. London 28: 106 (1871). *Acinodendron foliosum* (Triana) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Miconia subnodososa Triana, Trans. Linn. Soc. London 28: 106 (1871). *Acinodendron subnodosum* (Triana) Kuntze, Revis. Gen. Pl. 2: 953 (1891).

Section: Tamonea.

Distribution: Mexico, Belize, Guatemala, El Salvador, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador.

Miconia reflexipila Wurdack, Phytologia 23: 476 (1972). Type:—PERU. *Dudley* 13244 (Holotype: US).

Section: Amblyarrhena.

Distribution: Peru.

Miconia regelii Cogn. in Martius, Fl. bras. 14 (4): 405 pl. 83. (1888). *Acinodendron regelii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Riedel* s.n. (Syntypes: BR, LE).

Section: Chaenanthera.

Distribution: Guyana, Brazil North, Brazil Central-West.

Miconia remotiflora Urb., Symb. Antill. 9: 113 (1923). Type:—CUBA. *Ekman* 7162 (Holotype: S).

Section: Glossocentrum.

Distribution: Cuba.

Miconia renneri E.Cotton, Nordic J. Bot. 20: 179 (2000). Type:—ECUADOR. *Holm-Nielsen* 24564 (Holotype: AAU; isotypes: GB, MO, NY, QCA).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia resima Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 226 (1850). *Acinodendron resimum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Linden* 816 (Holotype: P; isotypes: G, GH, P).

Section: Cremanium.

Distribution: Colombia, Bolivia.

Miconia resimoides Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 926 (1891). *Acinodendron resimoides* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—VENEZUELA. *Fendler* 414 (Holotype: BR; isotypes: GH, GOET, K, MO, NY, PH, US).

Section: Cremanium.

Distribution: Venezuela.

Miconia reticulata Triana, Trans. Linn. Soc. London 28: 123 (1871). *Acinodendron reticulatum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Triana* 4008 (Holotype: BM; isotypes: BR, CAS, COL, E, K, NY).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia retusa Pilg., Verh. Bot. Vereins Prov. Brandenburg 47: 174 (1905). Type:—PERU. *Ule* 6366 (Holotype: B, destroyed; isotypes: CORD, HBG, K).

Miconia macbridei Gleason, Bull. Torrey Bot. Club 58: 246 (1931).

Section: Amblyarrhena.

Distribution: Peru.

Miconia rhodantha Wurdack, Phytologia 8: 168 (1962). Type:—COLOMBIA. *Daniel* 1624 (Holotype: US; isotype: MEDEL).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia rhombifolia Alain, Contr. Ocas. Mus. Hist. Nat. Colegio "De La Salle" 14: 4 (1955). Type:—CUBA. Shafer 4136 (Holotype: NY).
Section: Tamonea.
Distribution: Cuba.

Miconia rhonhofiae Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 15: 379 (1941). Type:—ECUADOR. Schultze-Rhonhof 2297 (Holotype: B, destroyed).
Section: Cremanium.
Distribution: Ecuador.

Miconia rhytidophylla Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 139 (1850). *Clidemia desmantha* Benth., J. Bot. 2: 309 (1840). *Acinodendron rhytidophyllum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. Schomburgk I 884 (Holotype: K; isotypes: A, BR, E, GH, NY, P, TCD).
Section: Miconia.
Distribution: Venezuela, Brazil North.

Miconia ricardoi Kriebel & Almeda, Harvard Pap. Bot. 17: 62 (2012). Type:—COSTA RICA. Kriebel 3384 (Holotype: INB; isotypes: CAS, CR, MO, NY).
Section: Cremanium.
Distribution: Costa Rica.

Miconia rigens Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 225 (1850). *Acinodendron rigens* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. Linden 1317 (Holotype: P; isotypes: BR, G, NY, P).
Section: Cremanium.
Distribution: Colombia.

Miconia rigida (Sw.) Triana, Trans. Linn. Soc. London 28: 130 (1871).
Basonym: *Melastoma rigidum* Sw., Prodr. 69 (1788). *Cremanium rigidum* (Sw.) Macfad., Fl. Jamaica 2: 93 (1850).
Pleurochaenia rigida (Sw.) Griseb., Abh. Königl. Ges. Wiss. Göttingen 9: 55 (1861). *Acinodendron rigidum* (Sw.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—JAMAICA. Swartz s.n. (Holotype: S; isotypes: LD, S).
Section: Chaenopleura.
Distribution: Jamaica, Dominican Republic.

Miconia rigidissima Urb. & Ekman, Ark. Bot. 20A (5): 26 (1926). Type:—HAITI. Ekman H1425 (Holotype: S; isotypes: EHH, NY, US).
Section: Chaenopleura.
Distribution: Haiti.

Miconia rimachii Wurdack, Brittonia 40: 7 (1988). Type:—PERU. Rimachi 6604 (Holotype: US; isotypes: BR, F, IBE, MO, NY).
Section: Jucunda.
Distribution: Peru, Brazil North, Brazil Central-West.

Miconia rimalis Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 238 (1850). *Acinodendron rimale* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. Blanchet 3623 (Holotype: P; isotypes: BM, BR, C, G, K, US).
Section: Chaenanthera.
Distribution: Brazil Northeast, Brazil Southeast.

Miconia rimbachii Wurdack, Phytologia 26: 10 (1973). Type:—ECUADOR. Rimbach 786 (Holotype: S; isotypes: F, US)
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia riparia Triana, Trans. Linn. Soc. London 28: 117 (1871). *Acinodendron riparium* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. Spruce 4589 (Holotype: K; isotypes: BM, BR, K, NY).
Section: Miconia.
Distribution: Peru.

Miconia rivalis Wurdack, Mem. New York Bot. Gard. 16: 10 (1967). Type:—ECUADOR. Camp E-46 (Holotype: US; isotypes: F, NY, S).
Section: Miconia.
Distribution: Ecuador, Peru.

Miconia rivetii Danguy & Cherm., Bull. Mus. Hist. Nat. 28: 433 (1922). Type:—ECUADOR. Rivet 816 (Holotype: P).
Section: Amblyarrhena.
Distribution: Colombia, Ecuador, Peru.

Miconia robinsoniana Cogn., Proc. Amer. Acad. Arts 38: 183 (1902). Type:—ECUADOR. Baur 163 (Holotype: BR; isotypes: GH, K).
Section: Tamonea.
Distribution: Panama, Ecuador.

Miconia robusta Cogn. in Martius, Fl. bras. 14 (4): 270 (1887). *Miconia robustissima* Cogn. in Martius, Fl. bras. 14 (4): 611 (1888), nom. illeg. *Acinodendron robustissimum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. Glaziou 8689 (Holotype: BR; isotypes: C, F, G, K, P, R, US).
Section: Miconia.
Distribution: Brazil Southeast.

Miconia roraimensis Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 362 (1915). Type:—GUYANA. Ule 8692 (Holotype: B, destroyed; isotypes: K, U).
Section: Amblyarrhena.
Distribution: Venezuela, Guyana.

Miconia rosea Gleason, Bull. Torrey Bot. Club 68: 247 (1941). Type:—BOLIVIA. Cárdenas 762 (Holotype: NY; isotypes: K, LIL, US).
Section: Octomeris.
Distribution: Bolivia.

Miconia rotundifolia (D.Don) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 235 (1850).
Basinonym: *Cremanium rotundifolium* D.Don, Mem. Wern. Nat. Hist. Soc. 4: 311 (1823). *Acinodendron rotundifolium* (D.Don) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. Ruiz & Pavón s.n. (Holotype: B, destroyed; isotypes: BR, G, K?, MA?).

Section: Chaenopleura.

Distribution: Ecuador, Peru.

Miconia rubens (Sw.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 169 (1850).
Basinonym: *Melastoma rubens* Sw., Prodr. 71 (1788). *Cremanium rubens* (Sw.) DC., Prodr. 3: 191 (1828). *Acinodendron rubens* (Sw.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—JAMAICA. Swartz s.n. (Lectotype: S; isolectotypes: BM, LD).

Melastoma polygamum Bonpl. ex Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 195 (1850), pro syn.

Miconia microbotrya Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 195 (1850).

Miconia rubens var. *latifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 921 (1891).

Section: Cremanium.

Distribution: Jamaica, Colombia, Venezuela.

Miconia rubiginosa (Bonpl.) DC., Prodr. 3: 183 (1828).

Basionym: *Melastoma rubiginosum* Bonpl., Monogr. Melast. 1: 109 (1816). *Acinodendron rubiginosum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Humboldt & Bonpland* 372 (Holotype: P).

Melastoma cupreum Sm. in Rees, Cycl. 23: 14 (1812).

Melastoma astrolasianum Spreng., Neue Entdeck. Pflanzenk. 3: 62 (1822). *Miconia astrolasia* (Spreng.) DC., Prodr. 3: 190 (1828).

Miconia rubiginosa var. *platyura* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 166 (1850).

Cremanium caracasanum Willd. ex. Triana, Trans. Linn. Soc. London 28: 112 (1871), *pro syn.*

Miconia rubiginosa var. *cinerascens* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 824 (1891).

Miconia rubiginosa var. *glabrior* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 824 (1891).

Miconia rubiginosa var. *rufa* Huber, Bull. Soc. Bot. Genève 6: 192 (1914).

Section: Miconia.

Distribution: Costa Rica, Panama, Dominican Republic, Puerto Rico, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia rubricans Triana, Trans. Linn. Soc. London 28: 121 (1871). *Acinodendron rubricans* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Triana s.n.* (Holotype: BM; isotype: COL).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia rufa (Griseb.) Triana, Trans. Linn. Soc. London 28: 130 (1871).

Basionym: *Catachaenia rufa* Griseb., Cat. Pl. Cub. 102 (1866). *Acinodendron rufum* (Griseb.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—CUBA. *Wright* 2524 (Holotype: GOET; isotypes: BM, BR, GH, GOET, K, MO, S, UC).

Miconia plumieriifolia Britton & P. Wilson, Bull. Torrey Bot. Club 50: 45 (1923).

Section: Chaenopleura.

Distribution: Cuba.

Miconia rufescens (Aubl.) DC., Prodr. 3: 180 (1828).

Basionym: *Melastoma rufescens* Aubl., Hist. pl. Guiane 1: 408 (1775). *Acinodendron rufescens* (Aubl.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—FRENCH GUIANA. *Aublet s.n.* (Holotype: BM).

Miconia impetiginosa DC., Prodr. 3: 183 (1828). *Melastoma impetiginosum* Mart. ex DC., Prodr. 3: 183 (1828), *pro syn.*

Miconia rufescens var. *grandifolia* Cogn. in Martius, Fl. bras. 14 (4): 274 (1887).

Section: Miconia.

Distribution: Colombia, Venezuela, Suriname, Guyana, French Guiana, Peru, Bolivia, Brazil North, Brazil Central-West.

Miconia ruficalyx Gleason, Brittonia 1: 181 (1932). *Copedesma nitens* Gleason, Bull. Torrey Bot. Club 52: 331 (1925).

Type:—GUYANA. *de la Cruz* 4542 (Holotype: NY; isotypes: F, GH, K, MO, PH, US).

Section: Miconia.

Distribution: Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Bolivia, Brazil North, Brazil Northeast.

Miconia rufipila Triana, Trans. Linn. Soc. London 28: 104 (1871). *Diplochita rufipila* Steud. ex Triana, Trans. Linn. Soc. London 28: 104 (1871), *pro syn.* *Acinodendron rufipilum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Blanchet* 3952 (Syntypes: BR, K).

Section: Tamonea.

Distribution: Brazil Northeast, Brazil Southeast.

Miconia rufiramea Wurdack, Phytologia 23: 496 (1972). Type:—PERU. *Dudley* 10919 (Holotype: US; isotype: NA).

Section: Cremanium.

Distribution: Peru.

Miconia rufostellulata Pittier, J. Wash. Acad. Sci. 13 (17): 390 (1923). Type:—PANAMA. *Pittier* 6535 (Holotype: US; isotype: VEN).

Section: Miconia.

Distribution: Panama.

Miconia rugifolia Triana, Trans. Linn. Soc. London 28: 125 (1871). *Acinodendron rugifolium* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Lechler* 2196 (Holotype: K; isotype: BR).
Section: Cremanium.
Distribution: Peru.

Miconia rugosa Triana, Trans. Linn. Soc. London 28: 106 (1871). *Acinodendron rugosum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Spruce* 1718 (Holotype: K; isotypes: F, K, M, P, TCD).

Miconia plumosa Gleason, Bull. Torrey Bot. Club 52: 381 (1925).

Section: Adenodesma.

Distribution: Colombia, Venezuela, Guyana, Suriname, Peru, Brazil North.

Miconia ruizii Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 192 (1850). *Acinodendron ruizii* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Ruiz & Pavón s.n.* (Holotype: P).

Section: Amblyarrhena.

Distribution: Ecuador, Peru, Bolivia.

Miconia ruizteranii Wurdack, Phytologia 22: 413 (1972). Type:—VENEZUELA. *Ruiz-Terán* 3204 (Holotype: US; isotype: MER).

Section: Cremanium.

Distribution: Venezuela.

Miconia rupestris Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 363 (1915). Type:—GUYANA. *Ule* 8690 (Holotype: B, destroyed; isotype: K).

Section: Cremanium.

Distribution: Venezuela, Guyana.

Miconia rupicalyx Wurdack, Mem. New York Bot. Gard. 10 (4): 35 (1961). Type:—VENEZUELA. *Maguire* 42013 (Holotype: NY; isotypes: F, GH, K, U, UC, US, VEN).

Section: Laceraria.

Distribution: Venezuela, Guyana, Ecuador, Bolivia.

Miconia rusbyana Cogn. in Britton, Bull. Torrey Bot. Club 17: 93 (1890). *Acinodendron rusbyanum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BOLIVIA. *Rusby* 2259 (Holotype: NY; isotypes: CAS, F, GH, MICH, PH, US, WIS).

Section: Cremanium.

Distribution: Bolivia.

Miconia ruschiana Caddah & R.Goldenb., Syst. Bot. 37: 974 (2012). Type:—BRAZIL. *Kollmann* 3424 (Holotype: RB; isotypes: MBML, UPCB).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia russea Wurdack, Phytologia 31: 495 (1975). Type:—ECUADOR. *Harling* 12723 (Holotype: GB; isotypes: F, MO, NY, S, US).

Section: Octomeris.

Distribution: Ecuador, Peru.

Miconia rzedowskii J.R.Santiago, Brittonia 64: 144 (2012). Type:—MEXICO. *Santiago* 1843 (Holotype: FCME; isotypes: CAS, ENCB, IEB, K, MEXU, MO, NY, P, UAMIZ, US, XAL).

Section: Cremanium.

Distribution: Mexico.

Miconia sagotiana Cogn. in Martius, Fl. bras. 14 (4): 299 (1887). *Acinodendron sagotianum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—FRENCH GUIANA. *Sagot* 230 (Syntype: BR; isotypes: BM, G, K, MPU, P, S, U).

Miconia membranicalyx Gleason, Bull. Torrey Bot. Club 58: 234 (1931).

Section: Miconia.

Distribution: Guyana, Suriname, French Guiana, Brazil North, Brazil Central-West.

Miconia saldanhae Cogn. in Martius, Fl. bras. 14 (4): 372 (1887). *Acinodendron saldanhae* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Saldanha* 6821 (Syntype: BR; isotypes: R, RB).

Miconia saldanhae var. *subsessilifolia* Cogn. in Martius, Fl. bras. 14 (4): 373 (1887).

Miconia saldanhae var. *grandiflora* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 841 (1891).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia salebrosa Wurdack, Acta Bot. Venez. 13: 140 (1978). Type:—VENEZUELA. *Wurdack* 2776 (Holotype: US; isotypes: F, NY).

Section: Octomeris.

Distribution: Venezuela.

Miconia salicifolia Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 234 (1850). *Melastoma salicifolium* Bonpl. ex Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 234 (1850), *pro syn.* *Acinodendron salicifolium* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Funck* 1315 (Syntype: P; isotypes: BR, MPU).

Melastoma rosmarinifolium Pav. ex Triana, Trans. Linn. Soc. London 28: 127 (1871), *pro syn.*

Section: Chaenopleura.

Distribution: Colombia, Venezuela, Ecuador, Peru.

Miconia saltuensis Wurdack, Phytologia 14: 272 (1967). Type:—PERU. *Hutchison* 5935 (Holotype: US; isotypes: UC, USM).

Section: Amblyarrhena.

Distribution: Peru.

Miconia samanensis Urb., Ark. Bot. 22A (17): 46 (1929). Type:—DOMINICAN REPUBLIC. *Abbott* 2923 (Lectotype: GH; Isolectotypes: A, BM).

Section: Chaenopleura.

Distribution: Dominican Republic.

Miconia sanctiphilippi Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 121 (1850). *Acinodendron sanctiphilippi* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—VENEZUELA. *Funck* 680 (Holotype: P; isotypes: BM, BR, G, MPU).

Miconia myura Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 129 (1850).

Section: Tamonea.

Distribution: Venezuela.

Miconia sandemanii Wurdack, Phytologia 5: 59 (1954). Type:—COLOMBIA. *Sandeman* 5837 (Holotype: K; isotypes: COL, NY, US).

Section: Cremanium.

Distribution: Colombia.

Miconia sanguinea (D.Don) Triana, Trans. Linn. Soc. London 28: 125 (1871).

Basionym: *Tococa sanguinea* D.Don, Mem. Wern. Nat. Hist. Soc. 4: 305 (1823). *Acinodendron sanguineum* (D.Don) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Ruiz & Pavón* s.n. (Holotype: OXF?: isotype: B, destroyed).

Cremanium hispidissimum DC., Prodr. 3: 192 (1828). *Miconia hispidissima* (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 191 (1850).

Section: Amblyarrhena.

Distribution: Peru, Bolivia.

Miconia santanana Judd & Skean, Brittonia 46: 101 (1994). Type:—DOMINICAN REPUBLIC. *Ekman* H11759 (Holotype: S; isotypes: GH, NY, US).

Section: Chaenopleura.

Distribution: Dominican Republic.

Miconia santaremensis Wurdack, Phytologia 22: 402 (1972). Type:—BRAZIL. *Ginzberger s.n.* (Holotype: W).

Section: Octomeris.

Distribution: Brazil North.

Miconia santaritensis Almeda, Brittonia 35: 44 (1983). Type:—PANAMA. *Antonio 4509* (Holotype: CAS; isotypes: MO, PMA).

Section: Miconia.

Distribution: Panama.

Miconia sarmentosa Cogn. in Martius, Fl. bras. 14 (4): 323 (1887). *Acinodendron sarmentosum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—FRENCH GUIANA. *Talbot s.n.* (Syntype: BR; isosyntype: K).

Section: Miconia.

Distribution: Suriname, French Guiana, Brazil North.

Miconia sastrei Wurdack, Phytologia 45: 329 (1980). Type:—FRENCH GUIANA. *Granville 849* (Holotype: CAY; isotypes: P, U, US).

Section: Laceraria.

Distribution: French Guiana, Brazil North.

Miconia savannarum R.O.Williams, Bull. Dept. Agric. Trinidad & Tobago 1: 374 (1934). Type:—TRINIDAD. *Britton 2468* (Syntype: K).

Section: Miconia.

Distribution: Trinidad & Tobago.

Miconia saxatilis J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 186 (1929). Type:—PERU. *Macbride 4881* (Holotype: F; isotype: NY).

Section: Cremanium.

Distribution: Peru.

Miconia saxicola Brandegee, Zoe 5: 215 (1905). Type:—MEXICO. *Brandegee s.n.* (Syntype: UC).

Section: Octomeris.

Distribution: Mexico.

Miconia scabra Cogn., Bull. Acad. Roy. Bel. 14: 956 (1887). *Acinodendron scabrum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—ECUADOR. *André 5880 p.p.* (Holotype: K; isotypes: BR, NY).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia schippii Standl., Publ. Field Mus. Nat. Hist., Bot. Ser. 8 (1): 29 (1930). Type:—BELIZE. *Schipp 220* (Holotype: F; isotypes: BKL, BM, GH, K, MICH, MO, NY, S).

Section: Hartigia.

Distribution: Mexico, Belize, Guatemala, Nicaragua.

Miconia schlechtendalii Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 804 (1891). *Acinodendron schlechtendalii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—MEXICO. *Schiede 551* (Syntype: BR; isosyntypes: K, MO, US).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, El Salvador, Honduras.

Miconia schlimii Triana, Trans. Linn. Soc. London 28: 102 (1871). *Acinodendron schlimii* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Schlism 903* (Syntype: BM; isosyntypes: BR, K, MPU).

Conostegia dolichostylis Donn.Sm., Bot. Gaz. 42: 294 (1906).

Section: Amblyarrhena.

Distribution: Belize, Guatemala, El Salvador, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela.

Miconia schnellii Wurdack, Phytologia 24: 203 (1972). Type:—COSTA RICA. *Schnell 711* (Holotype: US; isotype: F).

Section: Cremanium.

Distribution: Costa Rica.

Miconia schunkei Wurdack, Phytologia 24: 200 (1972). Type:—PERU. *Schunke 5756* (Holotype: US; isotypes: F, K, RSA).

Section: Miconia.

Distribution: Ecuador, Peru, Brazil North.

Miconia schwackei Cogn. in Martius, Fl. bras. 14 (4): 342 (1887). *Acinodendron schwackei* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Schwacke III.260* (Syntype: BR; isosyntypes: GOET, RB).

Section: Miconia.

Distribution: Brazil North.

Miconia sciurea L.Uribe, Caldasia 9: 464 (1966). Type:—COLOMBIA. *Gutiérrez 2524* (Holotype: COL; isotypes: F, NY).

Section: Tamonea.

Distribution: Colombia, Ecuador.

Miconia sclerophylla Triana, Trans. Linn. Soc. London 28: 119 (1871). *Acinodendron sclerophyllum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *S.coll*, s.n. (Holotype: K).

Section: Glossocentrum.

Distribution: Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia scutata Gleason, Bull. Torrey Bot. Club 52: 388 (1925). Type:—ECUADOR. *Hitchcock 21868* (Holotype: NY; isotypes: GH, US).

Section: Amblyarrhena.

Distribution: Ecuador.

Miconia secunda R.A.Howard & E.A.Kellogg, J. Arnold Arbor. 67: 250 (1986). Type:—ST. LUCIA. *Proctor 17726* (Holotype: A; isotype: BM).

Section: Glossocentrum.

Distribution: Santa Lucia.

Miconia secundiflora Cogn. in Martius, Fl. bras. 14 (4): 285 pl. 58 (1887). *Acinodendron secundiflorum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Riedel 1548* (Syntypes: BR, LE; isosyntypes: K, P).

Section: Miconia.

Distribution: Brazil North.

Miconia secundifolia Cogn., Bot. Jahrb. Syst. 42: 145 (1909). Type:—PERU. *Weberbauer 4427* (Holotype: B, destroyed; isotype: BR).

Miconia secundifolia ssp. *malcabalensis* Wurdack, Phytologia 11: 396 (1965).

Section: Chaenopleura.

Distribution: Peru.

Miconia selleiana Urb. & Ekman, Ark. Bot. 20A (5): 27 (1926). Type:—HAITI. *Ekman H1439* (Holotype: S; isotypes: EHH, NY).

Section: Chaenopleura.

Distribution: Haiti.

Miconia sellowiana Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 206 (1850). *Cremanium sellowanum* Cham. ex Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 206 (1850), *pro syn.* *Acinodendron sellowanum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Gaudichaud* 1360 (Holotype: P; isotypes: K, P).

Miconia sellowiana var. *pubescens* Cogn. in Martius, Fl. bras. 14 (4): 401 (1888).

Miconia sellowiana var. *quintuplinervis* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 863 (1891).

Section: Chaenanthera.

Distribution: Brazil Central-West, Brazil Southeast, Brazil South.

Miconia semisterilis Gleason, Bull. Torrey Bot. Club 63: 535 (1936). Type:—PERU. *Klug* 3224 (Holotype: NY; isotypes: GH, K, MO, S, US, WIS).

Section: Glossocentrum.

Distribution: Colombia, Peru.

Miconia seposita Wurdack, Phytologia 26: 6 (1973). Type:—ECUADOR. *Asplund* 18468 (Holotype: S).

Section: Amblyarrhena.

Distribution: Colombia, Ecuador.

Miconia septentrionalis Judd & R.S. Beaman, Brittonia 40: 383 (1988). Type:—DOMINICAN REPUBLIC. *Judd* 5173 (Holotype: FLAS; isotypes: A, JBSD, MO, NY, S, US).

Section: Chaenopleura.

Distribution: Dominican Republic.

Miconia serialis DC., Prodr. 3: 182 (1828). *Acinodendron seriale* (DC.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Ferreira* s.n. (Holotype: P).

Miconia tomentella Cogn. in Martius, Fl. bras. 14 (4): 284 (1887). *Acinodendron tomentellum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891).

Section: Miconia.

Distribution: Venezuela, Guyana, Suriname, French Guiana, Brazil North, Brazil Northeast.

Miconia serrulata (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 118 (1850).

Basionym: *Diplochita serrulata* DC., Prodr. 3: 177 (1828). *Melastoma serrulatum* Rich. ex DC., Prodr. 3: 177 (1828), *pro syn.* *Miconia macrophylla* var. *serrulata* (DC.) Cogn. in Martius, Fl. bras. 14 (4): 241 (1887). Type:—BRAZIL. *Martius* s.n. (Holotype: M).

Chitonia macrophylla Pav. ex D. Don, Mem. Wern. Nat. Hist. Soc. 4: 319 (1823). *Melastoma macrophyllum* Pav. ex D. Don, Mem. Wern. Soc. 4: 319 (1823), *pro syn.* *Diplochita macrophylla* (Pav. ex D. Don) DC., Prodr. 3: 177 (1828). *Miconia platyhedra* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850). *Miconia macrophylla* (Pav. ex D. Don) Triana, Trans. Linn. Soc. London 28: 103 (1871), *nom. illeg.* *Acinodendron macrophyllum* (Pav. ex D. Don) Kuntze, Revis. Gen. Pl. 1: 244 (1891).

Diplochita leucocephala DC., Prodr. 3: 177 (1828). *Miconia leucocephala* (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850). *Miconia macrophylla* var. *leucocephala* (DC.) Cogn. in Martius, Fl. bras. 14 (4): 241 (1887).

Diplochita serrulata var. *latifolia* DC., Prodr. 3: 177 (1828). *Miconia macrophylla* var. *latifolia* (DC.) Cogn. in Martius, Fl. bras. 14 (4): 240 (1887).

Melastoma balbisianum Ser. ex DC., Prodr. 3: 177 (1828), *pro syn.*

Melastoma patens Spreng. ex DC., Prodr. 3: 177 (1828), *pro syn.*

Melastoma lasiopetalum Mart. & Schrank ex DC., Prodr. 3: 177 (1828), *pro syn.* *Tamonea lasiopetala* DC. ex Triana, Trans. Linn. Soc. London 28: 103 (1871), *pro syn.*

Melastoma strangulatum Raddi, Mem. Soc. Ital. Modena 20: 139 (1829).

Melastoma macrophyllum Vell., Fl. Flum. 4: 180 (1825/1829).

Decaraphe hostmannii Steud., Flora 27: 722 (1844). *Miconia macrophylla* var. *hostmanni* (Steud.) Cogn. in Martius, Fl. bras. 14 (4): 241 (1887).

Miconia boissieriana Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 734 (1891). *Acinodendron boissierianum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Tamonea tomentosa var. *auriculata* Jenn., Ann. Carnegie Mus. 11 (1-2): 208: t. 22 (1917).

Section: Tamonea.

Distribution: Mexico, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Cuba, Jamaica, Haiti, Dominican Republic, Puerto Rico, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast.

Miconia sessilifolia Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 181 (1850). *Acinodendron sessilifolium* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BOLIVIA. *Pentland* 217 (Holotype: P; isotypes: BR, MO).

Miconia sessiliflora C.Muell. in Walpers, Ann. Bot. Syst. 6: 1225 (1865).

Section: Miconia.

Distribution: Bolivia.

Miconia seticaulis Wurdack, Phytologia 52: 63 (1982). Type:—ECUADOR. *Harling* 17633 (Holotype: GB; isotypes: S, US).

Section: Laceraria.

Distribution: Ecuador.

Miconia setosa Wurdack, Phytologia 5: 127 (1955). Type:—COLOMBIA. *Yepes-Agrego* 548 (Holotype: F).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia setosociliata Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 843 (1891). *Acinodendron setosociliatum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Glaziou* 16934 (Holotype: BR; isotypes: C, F, G, K, NY, P, R, RB, US, Z).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia setulosa Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 898 (1891). *Acinodendron setulosum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Lechler* 2637 (Holotype: GOET ; isotypes: BR, K).

Section: Cremanium.

Distribution: Peru, Bolivia.

Miconia shaferi Cogn. in Urban, Symb. Antill. 7: 312 (1912). Type:—CUBA. *Shafer* 8199 (Holotype: NY; isotypes: A, BR).

Section: Cremanium.

Distribution: Cuba.

Miconia shattuckii Standl., Contr. Arnold Arbor. 5: 119 t. 16 (1933). Type:—PANAMA. *Shattuck* 335 (Holotype: F; isotypes: F, MO).

Section: Amblyarrhena.

Distribution: Nicaragua, Costa Rica, Panama, Colombia.

Miconia shepherdii R.Goldenb. & Reginato, Brittonia 59: 334 (2007). Type:—BRAZIL. *Goldenberg* 71 (Holotype: UEC; isotype: US).

Section: Glossocentrum.

Distribution: Brazil Southeast.

Miconia silicicola Gleason, Bull. Torrey Bot. Club 58: 428 (1931). Type:—VENEZUELA. *Tate* 212 (Holotype: NY).

Section: Adenodesma.

Distribution: Venezuela, Guyana, Brazil North.

Miconia silverstonei Wurdack, Phytologia 69: 322 (1990). Type:—COLOMBIA. *Silverstone-Sopkin* 3849 (Holotype: CUVC; isotypes: CUVC, US).

Section: Chaenopleura.

Distribution: Colombia.

Miconia simplex Triana, Trans. Linn. Soc. London 28: 104 (1871). *Acinodendron simplex* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Seemann 1030* (Holotype: K; isotype: BR).
Section: Tamonea.
Distribution: Nicaragua, Costa Rica, Panama, Colombia, Venezuela.

Miconia sintenisii Cogn., Jahrb. Königl. Bot. Gart. Berlin 4: 281 (1886). *Acinodendron sintenisii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PUERTO RICO. *Sintenis 454* (Lectotype: US; isolectotype: S).

Miconia sintenisii var. *integrifolia* Cogn., Jahrb. Königl. Bot. Gart. Berlin 4: 282 (1886).
Section: Chaenopleura.
Distribution: Puerto Rico.

Miconia skeaniana Judd, Sida 16: 226 (1994). Type:—CUBA. *Ekman 8855* (Holotype: S; isotype: NY).
Section: Chaenopleura.
Distribution: Cuba.

Miconia smaragdina Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 186 (1850). *Acinodendron smaragdinum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Goudot s.n.* (Holotype: P).
Section: Miconia.
Distribution: Mexico, Belize, Costa Rica, Panama, Colombia, Venezuela, Ecuador.

Miconia smithii Cogn. ex Gleason, Bull. Torrey Bot. Club 52: 383 (1925). Type:—COLOMBIA. *Smith 1846* (Holotype: NY; isotypes: BR, E, F, GH, K, MICH, PH, US).
Section: Cremanium.
Distribution: Colombia.

Miconia sneidernii Wurdack, Phytologia 14: 271 (1967). Type:—COLOMBIA. *von Sneidern 5525* (Holotype: S; isotypes: BM, NY, S, US, VEN).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia sodiroi Wurdack, Phytologia 38: 292 (1978). Type:—ECUADOR. *Sodiro Add. 2* (Holotype: BR; isotypes: BR, P).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia sordida Triana, Trans. Linn. Soc. London 28: 122 (1871). *Acinodendron sordidum* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Triana 4031* (Holotype: BM; isotype: COL).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia sparrei Wurdack, Phytologia 35: 245 (1977). Type:—ECUADOR. *Sparre 19290* (Holotype: S; isotype: US).
Section: Tamonea.
Distribution: Costa Rica, Ecuador, Peru.

Miconia spatellophora Gleason, Bull. Torrey Bot. Club 57: 71 (1930). Type:—COLOMBIA. *Juzepczuk 6640* (Holotype: LE).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia speciosa (A.St.-Hil. & Naudin) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 243 (1850).
Basionym: *Augustinea speciosa* A.St.-Hil. & Naudin, Ann. Sci. Nat., Bot., Ser. 3 2: 145 (1844). *Acinodendron speciosum* (A.St.-Hil. & Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). *Tamonea speciosa* (A.St.-Hil. & Naudin) Krasser in Engler & Prantl, Nat. Pflanzenfam. 3, Abt. 7: 188 (1893). Type:—BRAZIL. *St. Hilaire s.n.* (Holotype: P; isotype: BR).

Section: Octomeris.

Distribution: Brazil Northeast, Brazil Southeast.

Miconia spennerostachya Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 187 (1850). *Acinodendron spennerostachy whole* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—PERU. *Mathews 1305* (Holotype: P; isotypes: E, FI, G, K, NY, P).

Miconia pauciglandulosa Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 183 (1850). *Acinodendron pauciglandulosum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Miconia spennerostachya var. *angustifolia* Cogn. in Britton, Bull. Torrey Bot. Club 17: 59 (1890).

Miconia aspiazui J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 190 (1929).

Miconia nectaria J.F.Macbr., Publ. Field Mus. Nat Hist., Bot. Ser. 4: 190 (1929).

Section: Miconia.

Distribution: Colombia, Ecuador, Peru, Bolivia, Brazil Central-West.

Miconia sphagnicola Urb. & Ekman, Ark. Bot. 23A (11): 19 (1931). Type:—DOMINICAN REPUBLIC. *Ekman H12808* (Holotype: S; isotypes: A, GH, IJ, NY, S).

Section: Chaenopleura.

Distribution: Dominican Republic.

Miconia spicellata Bonpl. ex Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 134 (1850). *Acinodendron spicellatum* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Bonpland s.n.* (Holotype: P; isotypes: G, P).

Miconia stictophylla Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 167 (1850).

Section: Miconia.

Distribution: Colombia, Venezuela.

Miconia spichigeri Wurdack, Candollea 44: 517 (1989). Type:—PERU. *Spichiger 2000* (Holotype: US; isotype: G).

Section: Miconia.

Distribution: Colombia, Peru, Brazil North.

Miconia spinulidentata Cogn. & Gleason, Brittonia 1: 125 (1932). Type:—COLOMBIA. *Lehmann 6024* (Holotype: K; isotype: BR).

Section: Cremanium.

Distribution: Colombia, Ecuador.

Miconia spinulosa Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 195 (1850). *Acinodendron spinulosum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—VENEZUELA. *Funck 96* (Holotype: P; isotypes: BM, BR, P).

Miconia inaperta Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 210 (1850). *Acinodendron inapertum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Miconia octoscenidium Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 871 (1891). *Acinodendron octoscenidium* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Miconia spinulosa var. *subintegrifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 872 (1891).

Section: Amblyarrhena.

Distribution: Colombia, Venezuela.

Miconia spireifolia Triana, Trans. Linn. Soc. London 28: 122 (1871). *Acinodendron spireifolium* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Triana 4043* (Holotype: BM).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia splendens (Sw.) Griseb., Fl. Brit. W. Indian Is. 256 (1860).

Basionym: *Melastoma splendens* Sw., Prodr. 70 (1788). Type:—JAMAICA. *Swartz s.n.* (Holotype: S).

Miconia obovalis Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 183 (1850).

Miconia elliptica Macfad., Fl. Jamaica 2: 88 (1850). *Acinodendron ellipticum* (Macfad.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Cuba, Jamaica, Puerto Rico, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast, Brazil South.

Miconia sprucei Triana, Trans. Linn. Soc. London 28: 108 (1871). *Acinodendron sprucei* (Triana) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Spruce* 2496 (Holotype: K; isotypes: BM, BR, E, GH, K, NY).

Section: Miconia.

Distribution: Colombia, Venezuela, Brazil North.

Miconia squamulosa (Sm.) Triana, Trans. Linn. Soc. London 28: 128 (1871).

Basionym: *Melastoma squamulosum* Sm. in Rees, Cycl. 23: 45 (1812). *Acinodendron squamulosum* (Sm.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—COLOMBIA. *Mutis* s.n. (Holotype: LINN 559.14).

Maieta argentea Vent., Mem. Math. Phys. Inst. nat. Fr. 8: 13 (1807). *Huberia argentea* (Vent.) DC., Prodr. 3: 168 (1828).

Miconia otodesma Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 205 (1850).

Section: Cremanium.

Distribution: Colombia, Venezuela.

Miconia staminea (Desr.) DC., Prod. 3: 187 (1828).

Basionym: *Melastoma stamineum* Desr., Lam. Encycl. Méth. Bot. 4: 53 (1797). *Leandra staminea* (Desr.) Raddi, Mem. Soc. Ital. Modena 20: 154 (1829). *Acinodendron stamineum* (Desr.) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—BRAZIL. *Dombey* s.n. (Holotype: P-JU).

Melastoma glabrescens Schrank & Mart. ex DC., Prod. 3: 187 (1828), *pro syn.*

Melastoma reticulata Vell., Fl. Flum. 4: 178 (1825/1829).

Jucunda lhotzkyana Cham., Linnaea 9: 457 (1834). *Miconia lhotzkyana* (Cham.) Triana, Trans. Linn. Soc. London 28: 151 (1871).

Miconia teysmanniana Regel, Gartenflora 16: 67 (1867).

Miconia staminea var. *parvifolia* Cogn. in Martius, Fl. bras. 14 (4): 231 (1887).

Section: Jucunda.

Distribution: Brazil Central-West, Brazil Southeast, Paraguay, Argentina.

Miconia staphidioides (Naudin) Triana, Trans. Linn. Soc. London 28: 102 (1871).

Basionym: *Clidemia staphidioides* Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 344 (1851). *Acinodendron staphidioides* (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BOLIVIA. *Weddell* 4733 (Holotype: P; isotypes: BR, P).

Section: Amblyarrhena.

Distribution: Bolivia.

Miconia steinbachii Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 9 (90): 1158 (1927). Type:—BOLIVIA. *Steinbach* 7186 (Holotype: B, destroyed; isotypes: F, GOET, K, MO, NY, PH, S).

Section: Tamonea.

Distribution: Bolivia.

Miconia stelligera Cogn. in Martius, Fl. bras. 14 (4): 275 pl. 56. (1887). *Acinodendron stelligerum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. *Poeppig* 2377 p.p. (Syntype: BR; isosyntype: MO).

Miconia clavistila Gleason, Bull. Torrey Bot. Club 58: 229 (1931).

Section: Miconia.

Distribution: Ecuador, Peru, Bolivia, Brazil North.

Miconia stellulata Gleason, Bull. Torrey Bot. Club 68: 247 (1941). Type:—BOLIVIA. *Krukoff* 11376 (Holotype: NY; isotypes: A, K, LP, MO, NY, RSA, S, U, UC, US, WIS).

Section: Jucunda.

Distribution: Bolivia, Brazil North.

Miconia stenobotrys (Rich.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 240 (1850).

Basionym: *Melastoma stenobotrys* Rich. in Bonpland, Monogr. Melast. 1: 66 (1811). *Chaenopleura stenobotrys* (Rich.) DC., Prodr. 3: 197 (1828). *Acinodendron stenobotryum* (Rich.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—HISPAÑOLA. Richard s.n. (Holotype: P?).

Miconia buchii Cogn. in Urban, Symb. Antill. 5: 448 (1908).

Miconia leptoneura Urb. & Ekman, Ark. Bot. 22A (17): 49 (1929).

Miconia artibonensis Urb. & Ekman, Ark. Bot. 22A (17): 43 (1929).

Miconia azuensis Urb. & Ekman, Ark. Bot. 22A (17): 50 (1929).

Section: Chaenopleura.

Distribution: Haiti.

Miconia stenocardia Cogn., Meded. Rijks-Herb. 19: 79 (1913). Type:—BOLIVIA. Herzog 1549 (Holotype: BR; isotypes: JE, S).

Section: Miconia.

Distribution: Bolivia.

Miconia stenophylla Wurdack, Mem. New York Bot. Gard. 16: 33 (1967). Type:—ECUADOR. Prieto P-312 (Holotype: US; isotypes: F, GH, MO, NY).

Section: Chaenopleura.

Distribution: Ecuador.

Miconia stenostachya DC., Prodr. 3: 181 (1828). *Melastoma stenostachyum* Schrank & Mart. ex DC., Prodr. 3: 181 (1828), pro syn. *Acinodendron stenostachyum* (DC.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. Martius s.n. (Holotype: M).

Miconia hypargyrea Miq., Linnaea 18: 622 (1844).

Melastoma fastigiatum Mart. ex Triana, Trans. Linn. Soc. London 28: 116 (1871), pro syn.

Melastoma holosericeum Steud. ex Triana, Trans. Linn. Soc. London 28: 116 (1871), pro syn.

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Trinidad & Tobago, Guyana, Suriname, Peru, Bolivia, Brazil North, Brazil Central-West, Brazil Northeast, Brazil Southeast, Brazil South, Paraguay.

Miconia stenouraea Triana, Trans. Linn. Soc. London 28: 104 (1871). *Acinodendron stenourum* (Triana) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—COLOMBIA. Triana s.n. (Holotype: K).

Section: Tamonea.

Distribution: Colombia.

Miconia stephananthera Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 359 (1915). Type:—BRAZIL. Ule 7609 (Holotype: B, destroyed; isotypes: K, NY).

Miconia palmetorum Pittier, Bol. Soc. Venez. Ci. Nat. 8: 314 (1942).

Section: Miconia.

Distribution: Colombia, Venezuela, Guyana, Suriname, Bolivia, Brazil North.

Miconia sterilis Gleason, Madroño 8 (6): 190 (1946). Type:—EL SALVADOR. Tucker 1100 (Holotype: UC; isotypes: F, K, LL, MICH, PH).

Section: Cremanium.

Distribution: El Salvador.

Miconia stevensiana Almeda, Brittonia 35: 46 (1983). Type:—NICARAGUA. Stevens 12148 (Holotype: CAS; isotypes: BM, HNMN, MEXU, MO, SI, U, US).

Section: Miconia.

Distribution: Honduras, Nicaragua, Costa Rica, Panama, Colombia.

Miconia steyermarkii Gleason in Steyermark, Fieldiana, Bot. 28 (2): 435 (1952). Type:—VENEZUELA. *Steyermark* 58262 (Holotype: F).
Section: Miconia.
Distribution: Venezuela.

Miconia stipitata Gleason, Bull. Torrey Bot. Club 72: 476 (1945). Type:—COLOMBIA. *Cuatrecasas* 13713 (Holotype: NY; isotypes: COL, F).
Section: Miconia.
Distribution: Colombia, Ecuador.

Miconia stipularis Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 194 (1850). *Acinodendron stipulare* (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—COLOMBIA. *Bonpland* s.n. (Lectotype: P; isolectotype: P).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia striata (Vahl) Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 765 (1891).
Basionym: *Melastoma striatum* Vahl, Eclog. Amer. 3: 14 (1807). Type:—MONTSERRAT. *Ryan* 5 (Holotype: C).
Melastoma acuminatum Desr. in Lamarck, Encycl. Méth. Bot. 4: 54 (1797). *Acinodendron acuminatum* (Desr.) Kuntze, Revis. Gen. Pl. 2: 949 (1891).
Melastoma elongatum Vahl, Eclog. Amer. 2: 16 (1798).
Melastoma ebracteatum F. Dietr., Nachtr. vollst. Lex. Gärtn. 4: 637 (1818).
Melastoma calyptatum Sieber ex Triana, Trans. Linn. Soc. London 28: 102 (1871), *pro syn.*
Miconia antiguensis Urb., Repert. Spec. Nov. Regni Veg. 17: 162 (1920).
Melastoma adpressum C. Presl ex Griseb., Fl. Brit. W. Indian Is. 255 (1860).
Section: Laceraria.
Distribution: Saint Kitts & Nevis, Antigua, Montserrat, Guadeloupe, Dominica, Martinique, Santa Lucia.

Miconia strigosa (Triana) Wurdack, Phytologia 22: 404 (1972).
Basionym: *Heterotrichum strigosum* Triana, Trans. Linn. Soc. London 28: 134 (1871). *Staphidium strigosum* Berg ex Triana, Trans. Linn. Soc. London 28: 134 (1871), *pro syn.* Type:—BRAZIL. *Spruce* 2047 (Holotype: K; isotypes: BM, BR, M).
Section: Miconia.
Distribution: Brazil North, Venezuela.

Miconia suaveolens Wurdack, Acta Bot. Venez. 13: 142 (1978). Type:—VENEZUELA. *Ruiz-Terán* 9204a (Holotype: US; isotypes: MO, PORT).
Section: Octomeris.
Distribution: Venezuela.

Miconia subalpina Gleason, Bull. Torrey Bot. Club 52: 447 (1925). Type:—COLOMBIA. *Pennell* 10504 (Holotype: NY).
Section: Amblyarrhena.
Distribution: Colombia.

Miconia subandicola Wurdack, Phytologia 23: 474 (1972). *Miconia glandulifera* var. *boliviensis* Cogn. in Rusby, Mem. Torrey Bot. Club 6: 38 (1896). Type:—BOLIVIA. *Bang* 1173 (Holotype: BR; isotypes: BR, CAS, MO, NY, US).
Section: Tamonea.
Distribution: Peru, Bolivia, Brazil North.

Miconia subcompressa Urb., Ark. Bot. 17A (7): 45 (1921). Type:—HAITI. *Ekman* H206 (Holotype: S).
Miconia plumerii Urb. & Ekman, Ark. Bot. 22A (17): 38 (1929).
Section: Chaenopleura.
Distribution: Haiti, Dominican Republic.

Miconia subcordata Cogn. in Martius, Fl. bras. 14 (4): 244 (1887). *Acinodendron subcordatum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. *Blanchet 1101* (Syntype: G; isosyntype: LE).
Section: Tamonea.
Distribution: Brazil Northeast.

Miconia subglabra Cogn. in Martius, Fl. bras. 14 (4): 412 (1888). *Acinodendron subglabrum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—PERU. *Spruce 4842* (Syntype: BR; isosyntypes: BM, E, G, GH, GOET, K, LD, LE, NY, P, TCD).
Section: Amblyarrhena.
Distribution: Peru.

Miconia submacrophylla Gleason, Bull. Torrey Bot. Club 57: 65 (1930). Type:—COLOMBIA. *Woronow 6084* (Holotype: LE; isotype: NY).
Section: Tamonea.
Distribution: Colombia.

Miconia suborbicularis Cogn., Bull. Acad. Belg., Ser. 3 14: 958 (1887). *Acinodendron suborbiculare* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—ECUADOR. *André 3989 bis* (Holotype: K; isotype: BR).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia subsessilifolia Wurdack, Phytologia 55: 141 (1984). Type:—COLOMBIA. *Hernandez 61* (Holotype: HUA; isotypes: COL, US).
Section: Miconia.
Distribution: Colombia.

Miconia subsimplex Pilg., Verh. Bot. Vereins Prov. Brandenburg 47: 174 (1905). Type:—BRAZIL. *Ule 6155* (Holotype: B, destroyed; isotypes: HBG, K).
Section: Miconia.
Distribution: Brazil North.

Miconia subspicata Wurdack, Phytologia 41: 1 (1978). Type:—ECUADOR. *Madison 5313* (Holotype: US; isotypes: F, QCA, SEL, US).
Section: Miconia.
Distribution: Ecuador, Peru, Brazil North.

Miconia subulipetala Wurdack, Phytologia 48: 245 (1981). Type:—PERU. *Rimachi 3421* (Holotype: US; isotype: MISSA).
Section: Miconia.
Distribution: Peru.

Miconia subverncosa Cogn. in Martius, Fl. bras. 14 (4): 374 (1887). *Acinodendron subvernicosum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. *Glaziou 590* (Syntype: BR; isosyntypes: C, P, R).
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia sulcata J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 186 (1929). Type:—PERU. *Schunke 423* (Holotype: F).
Section: Cremanium.
Distribution: Peru.

Miconia summa Cuatrec., Trab. Mus. Nac. Ci. Nat., Ser. Bot. 26: 25 (1933). Type:—COLOMBIA. *Cuatrecasas 2510* (Holotype: MA; isotype: F).
Section: Cremanium.
Distribution: Colombia.

Miconia superba Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 356 (1915). Type:—GUYANA. *Ule 8695* (Holotype: B, destroyed; isotype: K).
Miconia pachypoda Gleason in Steyermark, Fieldiana, Bot. 28 (2): 435 (1952).
Section: Tamonea.
Distribution: Venezuela, Guyana.

Miconia superposita Wurdack, Phytologia 5: 130 (1955). Type:—COLOMBIA. *Cuatrecasas 23324* (Holotype: NY; isotypes: F, US).
Section: Chaenopleura.
Distribution: Colombia.

Miconia sylvatica (Schltdl.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 243 (1850).
Basionym: *Melastoma sylvaticum* Schltdl., Linnaea 13 (5): 422 (1839). Type:—MEXICO. *Schiede s.n.* (Holotype: HAL; isotypes: BR, G-DEL).
Acinodendron glandulosum Kuntze, Revis. Gen. Pl. 2: 951 (1891). *Miconia glandulosa* (Kuntze) R. Knuth, Fedde Repert. Beih. 43: 531 (1928).
Section: Miconia.
Distribution: Mexico, Guatemala, El Salvador, Honduras, Venezuela.

Miconia symplocoidea Triana, Trans. Linn. Soc. London 28: 108 (1871). *Acinodendron symplocoideum* (Triana) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—COLOMBIA. *Triana 4017* (Holotype: BM?; isotypes: G-DEL, K, NY, US).
Section: Miconia.
Distribution: Colombia.

Miconia tabayensis Wurdack, Phytologia 21: 359 (1971). *Miconia micrantha* Pittier, Bol. Soc. Venez. Ci. Nat. 11: 27 (1947), nom. illeg.. Type:—VENEZUELA. *Gehringer 595* (Holotype: VEN; isotypes: F, US).
Section: Cremanium.
Distribution: Venezuela.

Miconia tacanensis Wurdack, Phytologia 16: 181 (1968). Type:—MEXICO. *Matuda 2939* (Holotype: MICH; isotypes: A, CAS, K, LL, MICH).
Section: Amblyarrhena.
Distribution: Mexico, Guatemala.

Miconia tachirensis Wurdack, Acta Bot. Venez. 4: 63 (1970). Type:—VENEZUELA. *Steyermark 101038* (Holotype: US; isotypes: NY, VEN).
Section: Amblyarrhena.
Distribution: Venezuela.

Miconia talamanicensis Almeda, Proc. Calif. Acad. Sci., Ser. 4 52: 48 (2000). Type:—COSTA RICA. *Almeda 4777* (Holotype: CAS; isotypes: CR, INB, MO).
Section: Cremanium.
Distribution: Costa Rica, Panama.

Miconia tamana Wurdack, Acta Bot. Venez. 4: 64 (1970). Type:—VENEZUELA. *Steyermark 100878* (Holotype: US; isotypes: COL, NY, VEN).
Section: Cremanium.
Distribution: Colombia, Venezuela.

Miconia tenensis Markgr., Notizbl. Bot. Gart. Berlin-Dahlem 12: 178 (1934). Type:—ECUADOR. Heinrichs 288 (Holotype: B, destroyed; isotypes: M, NY).
Section: Miconia.
Distribution: Ecuador, Peru.

Miconia tentaculifera Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 241 (1850). *Acinodendron tentaculiferum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. Claussen 54 (Lectotype: P; isolectotypes: F, G, L).
Section: Chaenanthera.

Distribution: Brazil Southeast.

Miconia tenuis Triana, Trans. Linn. Soc. London 28: 120 (1871). *Acinodendron tenue* (Triana) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—COLOMBIA. Triana 4058 (Holotype: BM).
Section: Chaenanthera.
Distribution: Colombia.

Miconia teotepicensis J.R.Santiago, Acta Bot. Mex. 50: 22 (2000). Type:—MEXICO. Santiago 167 (Holotype: FCME; isotypes: CAS, CHAPA, ENCB, IBUG, IEB, K, MEXU, MO, US).
Section: Amblyarrhena.
Distribution: Mexico.

Miconia tephrodes Wurdack, Mem. New York Bot. Gard. 16: 32 (1967). Type:—ECUADOR. Camp E-4872 (Holotype: US; isotypes: F, NY).
Section: Chaenopleura.
Distribution: Ecuador.

Miconia tepicana Standl., Contr. U.S. Natl. Herb. 23 (4): 1064 (1924). Type:—MEXICO. Rose 2058 (Holotype: US; isotypes: GH, NY).
Section: Amblyarrhena.
Distribution: Mexico.

Miconia terborghii Wurdack, Phytologia 23: 483 (1972). Type:—PERU. Dudley 10890 (Holotype: US; isotype: NA).
Section: Amblyarrhena.
Distribution: Peru.

Miconia terera Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 196 (1850). *Acinodendron terera* (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—PERU. Dombey s.n. (Holotype: P).
Section: Amblyarrhena.
Distribution: Peru.

Miconia ternatifolia Triana, Trans. Linn. Soc. London 28: 118 (1871). *Acinodendron ternatifolium* (Triana) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—PERU. Mathews 1720 (Syntypes: K, L, NY).
Section: Glossocentrum.
Distribution: Colombia, Ecuador, Peru, Bolivia, Brazil North.

Miconia tetragona Cogn. in Martius, Fl. bras. 14 (4): 387 (1887). *Acinodendron tetragonum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. Spruce 2747 (Syntype: BR; isosyntypes: BM, G, K, LE, NY, TCD).
Section: Glossocentrum.
Distribution: Peru, Brazil North.

Miconia tetrandra (Sw.) D.Don in Loudon, Hort. Brit. 1: 174 (1830).
Basionym: *Melastoma tetrandrum* Sw., Prodr. 72 (1788). *Tetrazygia tetrandra* (Sw.) DC., Prodr. 3: 172 (1828).
Cremanium tetrandrum (Sw.) Griseb., Pl. Wright. 1: 186 (1860). *Acinodendron tetrandrum* (Sw.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—JAMAICA. Swartz s.n. (Lectotype: S; isolectotypes: BM, C).

Melastoma farinulentum Sw. ex Wikstr., Kongl. Vetensk. Akad. Handl. 1827: 65 (1827).

Section: Cremanium.

Distribution: Cuba, Jamaica, Haiti, Dominican Republic, Puerto Rico Guadeloupe, Dominica, Grenada.

Miconia tetrasperma Gleason, Bull. Torrey Bot. Club 58: 237 (1931). Type:—PERU. *Williams* 3757 (Holotype: NY).

Section: Miconia.

Distribution: Colombia, Peru, Bolivia, Brazil North.

Miconia tetraspermoides Wurdack, Phytologia 18: 155 (1969). Type:—VENEZUELA. *Lasser* 1874 (Holotype: NY; isotype: F).

Section: Miconia.

Distribution: Venezuela, Guyana, Suriname, French Guiana, Brazil North, Brazil Central-West.

Miconia tetrastoma Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 236 (1850). *Acinodendron tetrastomum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—CUBA. *Linden* 2002 (Holotype: P; isotype: BR).

Section: Cremanium.

Distribution: Cuba, Haiti, Dominican Republic, Puerto Rico.

Miconia tetrazygioides Urb. & Ekman, Ark. Bot. 22A (17): 37 (1929). Type:—HAITI. *Ekman* H10684 (Holotype: S; isotypes: K, NY, US).

Seção: Chaenanthera

Distribution: Haiti.

Miconia thaminantha Wurdack, Phytologia 23: 494 (1972). Type:—PERU. *Dudley* 10531 (Holotype: US; isotype: NA).

Miconia thaminantha ssp. *huanucensis* Wurdack, Phytologia 23: 495 (1972).

Section: Cremanium.

Distribution: Peru.

Miconia theizans (Bonpl.) Cogn. in Martius, Fl. bras. 14 (4): 419 (1888).

Basionym: *Melastoma theizans* Bonpl., Monogr. Melast. 1: 17 (1807). *Cremanium theizans* (Bonpl.) DC., Prodr. 3: 194 (1828). *Acinodendron theizans* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—COLOMBIA. *Humboldt & Bonpland* 2026 (Holotype: P).

Cremanium glaberrimum DC., Prodr. 3: 193 (1828). *Melastoma dimorphum* Schrank & Mart. ex DC., Prodr. 3: 193 (1828), pro syn. *Miconia dimorpha* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850). *Miconia theizans* var. *glaberrima* (DC.) Cogn. in Martius, Fl. bras. 14 (4): 420 (1888).

Cremanium milleflorum DC., Prodr. 3: 193 (1828). *Melastoma milleflorum* Schrank & Mart. ex DC., Prodr. 3: 193 (1828), pro syn. *Hartigia milleflora* (DC.) Miq., Linnaea 18: 286 (1844). *Miconia milleflora* (DC.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 237 (1850). *Miconia theizans* var. *milleflora* (DC.) Cogn. in Martius, Fl. bras. 14 (4): 422 (1888).

Cremanium minutiflorum DC., Prodr. 3: 193 (1828). *Miconia hartigoides* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850). *Miconia theizans* var. *minutiflora* (DC.) Cogn. in Martius, Fl. bras. 14 (4): 420 (1888).

Cremanium paludosum Gardner, Hooker's J. Bot. Kew Gard. Misc. 2: 347 (1843). *Miconia paludosa* (Gardner) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 245 (1850). *Miconia theizans* var. *paludosa* (Gardner) Cogn. in Martius, Fl. bras. 14 (4): 420 (1888).

Cremanium ambiguum Benth., Pl. Hartw. 181 (1845).

Miconia multinervulosa Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 926 (1891). *Acinodendron multinervulosum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Miconia theizans var. *setulosa* Hoehne, Anexos Mem. Inst. Butantan, Secc. Bot. 1 (5): 154 (1922).

Miconia theizans ssp. *viridis* Cogn. in Martius, Fl. bras. 14 (4): 420 (1888).

Miconia theizans var. *genuina* Cogn. in Martius, Fl. bras. 14 (4): 420 (1888).

Miconia theizans ssp. *flavescens* Cogn. in Martius, Fl. bras. 14 (4): 421 (1888).

Miconia theizans var. *cuneata* Cogn. in Martius, Fl. bras. 14 (4): 421 (1888).

Miconia theizans var. *glazioviana* Cogn. in Martius, Fl. bras. 14 (4): 421 (1888).

Miconia theizans var. *lanceolata* Cogn. in Martius, Fl. bras. 14 (4): 421 (1888).

Miconia theizans var. *longifolia* Cogn. in Martius, Fl. bras. 14 (4): 421 (1888).

Miconia theizans var. *spinulosa* Cogn. in Martius, Fl. bras. 14 (4): 421 (1888).

Miconia theizans var. *subtriplinervia* Cogn. in Martius, Fl. bras. 14 (4): 421 (1888).

Miconia theizans var. *tetragona* Cogn. in Martius, Fl. bras. 14 (4): 421 (1888).

Miconia theizans var. *vulgaris* Cogn. in Martius, Fl. bras. 14 (4): 421 (1888).

Miconia theizans var. *integrifolia* Cogn. in Martius, Fl. bras. 14 (4): 422 (1888).

Miconia theizans var. *parvifolia* Cogn. in Martius, Fl. bras. 14 (4): 422 (1888).

Miconia theizans var. *saldanhae* Cogn. in Martius, Fl. bras. 14 (4): 422 (1888).

Miconia theizans var. *triplinervia* Cogn. in Martius, Fl. bras. 14 (4): 422 (1888).

Section: Cremanium.

Distribution: Honduras, Costa Rica, Panama, Jamaica, Colombia, Venezuela, Guyana, Ecuador, Peru, Bolivia, Brazil Central-West, Brazil Northeast, Brazil Southeast, Brazil South, Paraguay.

Miconia thomasiana DC., Prod. 3: 189 (1828). *Acinodendron thomasianum* (DC.) Kuntze, Revis. Gen. Pl. 2: 953: 1 (1891). *Tamonea thomasiana* (DC.) O.F.Cook & G.N.Collins, Contr. U. S. Natl. Herb. 8: 249 (1903). Type:—UNKNOWN. *S.coll.*, s.n. (Holotype: G-DC).

Melastoma coriaceum Juss. ex DC., Prodr. 3: 190 (1828), *pro syn.*

Miconia vernicosa Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 191 (1850).

Melastoma coriaceum Sessé & Moc., Fl. mexic., ed. 2: 106 (1894), *nom. illeg.*

Section: Octomeris.

Distribution: Puerto Rico, Virgin Islands.

Miconia thyrsiflora (D.Don) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850).

Basionym: *Cremanium thyrsiflorum* D.Don, Mem. Wern. Nat. Hist. Soc. 4: 312 (1823). Type:—PERU. *Pavón* s.n. (Holotype: G-BOISS; isotype: OXF). *Miconia thyrsoides* Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850).

Acinodendron thyrsoides (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891).

Miconia integrifolia Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 936 (1891). *Acinodendron cogniauxii* Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Section: Chaenopleura.

Distribution: Peru.

Miconia thysanophylla Wurdack, Brittonia 40: 13 (1988). Type:—PERU. *Kayap 1414* (Holotype: US; isotype: MO).

Section: Miconia.

Distribution: Peru.

Miconia tiliifolia Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 151 (1850). *Acinodendron tiliifolium* (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BOLIVIA. *Orbigny 870* (Holotype: P).

Section: Miconia.

Distribution: Peru, Bolivia, Brazil North, Brazil Central-West.

Miconia tillettii Wurdack, Mem. New York Bot. Gard. 10 (5): 167 (1964). Type:—GUYANA. *Tillett 44977* (Holotype: US; isotypes: K, GH, NY).

Section: Miconia.

Distribution: Venezuela, Guyana, Suriname, French Guiana, Brazil North.

Miconia tinifolia Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 225 (1850). *Acinodendron tinifolium* (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—VENEZUELA. *Linden 438* (Holotype: P; isotypes: F, FI, G, K, MPU, NY, P, TCD).

Miconia tinifolia var. *parvifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 917 (1891).

Miconia tinifolia var. *caudata* Wurdack, Phytologia 22: 408 (1972).

Miconia tinifolia var. *roraimensis* Wurdack, Phytologia 22: 408 (1972).

Section: Cremanium.

Distribution: Colombia, Ecuador, Venezuela, Guyana.

Miconia tiri Triana, Trans. Linn. Soc. London 28: 127 (1871). *Acinodendron tiri* (Triana) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—PERU. *Dombey s.n.* (Syntypes: BR, P).

Section: Amblyarrhena.

Distribution: Peru.

Miconia titanophylla Gleason, Bull. Torrey Bot. Club 57: 67 (1930). Type:—COLOMBIA. Woronow 5844 (Holotype: NY; isotypes: NY, LE)
Section: Tamonea.
Distribution: Colombia.

Miconia tixicensis Standl. & Steyermark, Publ. Field Mus. Nat. Hist., Bot. Ser. 23 (4): 181 (1944). Type:—GUATEMALA. Steyermark 31585 (Holotype: F).
Section: Octomeris.
Distribution: Guatemala.

Miconia tomentosa (Rich.) D.Don ex DC., Prodr. 3: 183 (1828).
Basionym: *Melastoma tomentosum* Rich., Actes Soc. Hist. Nat. Paris 1: 109 (1792). *Jucunda tomentosa* (Rich.) Benth., J. Bot. 2: 302 (1840). *Diplochita tomentosa* (Rich.) Griseb., Fl. Brit. W. Indian Is. 252 (1860). *Acinodendron tomentosum* (Rich.) Kuntze, Revis. Gen. Pl. 1: 245 (1891). Type:—FRENCH GUIANA. *Richard s.n.* (Holotype: P).
Melastoma megalophyllum Mart. ex DC., Prod. 3: 183 (1828), *pro syn.*
Pogonorhynchus amplexans Crueg., Linnaea 20: 107 (1847). *Miconia amplexans* (Crueg.) Cogn. in Martius, Fl. bras. 14 (4): 256 (1887). *Acinodendron amplexans* (Crueg.) Kuntze, Revis. Gen. Pl. 2: 950 (1891).
Pogonorhynchus sessilifolius Crueg., Linnaea 20: 107 (1847).
Miconia umbrifera Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 116 (1850).
Miconia tomentosa var. *ovata* Cogn. in Martius, Fl. bras. 14 (4): 255 (1887).
Miconia symplectocaulos Pilg., Verh. Bot. Vereins Prov. Brandenburg 47: 175 (1905).
Section: Adenodesma.
Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Cuba, Colombia, Venezuela, Trinidad-Tobago, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil North, Brazil Northeast, Brazil West-central.

Miconia tonduzii Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 1191 (1891). *Acinodendron tonduzii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—COSTA RICA. *Tonduz* 2115 (Neotype: BR; isoneotype: US).
Miconia tonduzii var. *cuneata* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 1191 (1891).
Miconia tonduzii var. *furfuracea* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 1191 (1891).
Miconia tonduzii var. *latifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 1191 (1891).
Miconia tonduzii var. *oblongifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 1191 (1891).
Miconia tonduzii var. *parvifolia* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 1191 (1891).
Miconia tonduzii var. *serrulata* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 1191 (1891).
Miconia tonduzii var. *hirtella* Almeda, Novon 15: 514 (2005).
Section: Cremanium.
Distribution: Nicaragua, Costa Rica, Panama.

Miconia toroi Gleason, Bull. Torrey Bot. Club 54: 617 (1927). Type:—COLOMBIA. *Toro* 278 (Holotype: NY).
Section: Hartigia.
Distribution: Colombia.

Miconia tovarensis Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 917 (1891). *Acinodendron tovarensense* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—VENEZUELA. *Fendler* 419 (Holotype: BR; isotypes: GH, K, MO, NY, PH, TCD).
Section: Cremanium.
Distribution: Colombia, Venezuela.

Miconia traillii Cogn. in Martius, Fl. bras. 14 (4): 242 (1887). *Acinodendron traillii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. *Trail* 267 (Holotype: K; isotype: GH).
Section: Tamonea.
Distribution: Colombia, Venezuela, Guyana, Suriname, French Guiana, Peru, Brazil North, Brazil Central-West.

Miconia transversa Gleason, Bull. Torrey Bot. Club 72: 475 (1945). Type:—COLOMBIA. Cuatrecasas 17459 (Holotype: NY; isotypes: BC, COL, F, MO, US).
Section: Tamonea.
Distribution: Colombia, Ecuador.

Miconia trianae Cogn. in Martius, Fl. bras. 14 (4): 394 (1887). *Acinodendron trianae* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. Martius 943 (Lectotype: BR; isolectotypes: BM, CAS, FI, G, K, L, LE, M, MO, P).

Miconia brevipes var. *longifolia* Cogn. in Martius, Fl. bras. 14 (4): 345 (1887).

Miconia hispida Cogn. in Martius, Fl. bras. 14 (4): 395 (1887). *Acinodendron hispidum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 951 (1891).

Miconia regnellii Cogn. in Martius, Fl. bras. 14 (4): 396 (1887). *Acinodendron regnellii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Section: Chaenanthera.

Distribution: Brazil Northeast, Brazil Southeast.

Miconia triangularis Gleason, Bull. Torrey Bot. Club 59: 369 (1932). Type:—COLOMBIA. Klug 1697 (Holotype: NY; isotypes: GH, MO).

Section: Tamonea.

Distribution: Colombia, Ecuador, Peru.

Miconia tricaudata Wurdack, Brittonia 28: 140 (1976). Type:—COLOMBIA. Barclay 6579 (Holotype: US; isotypes: COL, MO).

Section: Cremanium.

Distribution: Colombia.

Miconia trichogona J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 187 (1929). Type:—PERU. Macbride 5178 (Holotype: F; isotypes: NY, US).

Section: Cremanium.

Distribution: Peru.

Miconia trichophora Gleason, Bull. Torrey Bot. Club 57: 68 (1930). Type:—COLOMBIA. Killip 14848 (Holotype: NY; isotypes: A, GH, K, MO).

Section: Miconia.

Distribution: Colombia.

Miconia trichotoma (Desr.) DC., Prodr. 3: 188 (1828).

Basionym: *Melastoma trichotomum* Desr. in Lamarck, Encycl. Méth. Bot. 4: 53 (1797). *Acinodendron trichotomum* (Desr.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—"ANTILLES". *Herb. Surian* 774 (Holotype: P-JU)

Melastoma nigricans Vahl, Eclog. Amer. 3: 26 (1807).

Melastoma matuba Spreng., Neue Entdeck. Pflanzenk. 3: 61 (1822).

Miconia rivoeriae Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 170 (1850). *Cremanium rivoeriae* (Naudin) Griseb., Fl. Brit. W. Indian Is. 262 (1860).

Section: Glossocentrum.

Distribution: Antigua, Guadeloupe, Dominica.

Miconia trimera Wurdack, Phytologia 45: 327 (1980). Type:—FRENCH GUYANA. Granville 2181 (Holotype: US; isotypes: CAY, RB).

Section: Miconia.

Distribution: Suriname, French Guiana, Brazil North.

Miconia trinervia (Sw.) D.Don ex Loudon, Hort. Brit. 1: 174 (1830).

Basionym: *Melastoma trinervium* Sw., Prodr. 69 (1788). *Acinodendron trinervium* (Sw.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—JAMAICA. Swartz s.n. (Lectotype: S).

Melastoma scorpioides Schlehd. & Cham., Linnaea 5: 564 (1830). *Miconia scorpioides* (Schlehd. & Cham.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 243 (1850). *Acinodendron scorpioides* (Schlehd. & Cham.) Kuntze, Revis. Gen. Pl. 2: 952 (1891).

Miconia anceps Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 150 (1850).

Section: Miconia.

Distribution: Mexico, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Jamaica, Colombia, Venezuela, French Guiana, Ecuador, Peru, Bolivia, Brazil North.

Miconia triplinervis Ruiz & Pav., Syst. Veg. Fl. Peruv. Chil. 1: 105 (1798). *Acinodendron triplinerve* (Ruiz & Pav.) Kuntze, Revis. Gen. Pl. 1: 245 (1891). Type:—PERU. Ruiz & Pavón s.n. (Holotype: not located).

Cremanium trinerve Macfad., Fl. Jamaica 2: 94 (1850). *Miconia trinervis* (Macfad.) Griseb., Fl. Brit. W. Indian Is. 257 (1860).

Miconia spicata Macfad. ex Griseb., Fl. Brit. W. Indian Is. 257 (1860).

Miconia hexaptera Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 129 (1850).

Miconia haplobotrya Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 131 (1850).

Miconia angustispica S.F.Blaeke, Contr. U.S. Natl. Herb. 24: 15 (1922).

Miconia triplinervis ssp. *exalluvia* Wurdack, Mem. New York Bot. Gard. 16: 10 (1967).

Section: Miconia.

Distribution: Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Jamaica, Colombia, Venezuela, Ecuador, Peru, Bolivia, Brazil North.

Miconia tristis Spring, Flora 20 (2, Beibl.): 76 (1837). *Acinodendron triste* (Spring) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. Martius s.n. (Holotype: M).

Miconia tristis ssp. *australis* Wurdack, Sellowia 14: 60 (1962).

Section: Glossocentrum.

Distribution: Brazil Northeast, Brazil Southeast, Brazil South, Paraguay.

Miconia trujillensis Wurdack, Phytologia 22: 406 (1972). Type:—VENEZUELA. Steyermark 104813 (Holotype: US; isotype: VEN).

Section: Amblyarrhena.

Distribution: Venezuela.

Miconia truncata Triana, Trans. Linn. Soc. London 28: 102 (1871). *Acinodendron truncatum* (Triana) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. Spruce 1952 (Holotype: K; isotypes: BR, K).

Section: Jucunda.

Distribution: Colombia, Venezuela, Brazil North.

Miconia tschudyooides Cogn. in Martius, Fl. bras. 14 (4): 327 (1887). *Tschudya robusta* Sagot, Ann. Sci. Nat., Bot., Ser. 6 15: 329 (1883). *Miconia robusta* (Sagot) Cogn. in Martius, Fl. bras. 14 (4): 611 (1888), nom. illeg. *Acinodendron robustum* (Sagot) Kuntze, Revis. Gen. Pl. 2: 952 (1891). Type:—FRENCH GUIANA. Sagot 220 (Holotype: P; isotypes: BR, G, US).

Clidemia trichodes Sagot ex Cogn. in Martius, Fl. bras. 14 (4): 612 (1888).

Section: Miconia.

Distribution: Suriname, French Guiana.

Miconia tuberculata (Naudin) Triana, Trans. Linn. Soc. London 28: 102 (1871).

Basionym: *Octomeris tuberculata* Naudin, Ann. Sci. Nat., Bot., Ser. 3 4: 53 (1845). *Acinodendron tuberculatum* (Naudin) Kuntze, Revis. Gen. Pl. 1: 245 (1891). Type:—VENEZUELA. Linden 293 (Holotype: P; isotype: P).

Octomeris rostrata Naudin, Ann. Sci. Nat., Bot., Ser. 3 4: 53 (1845). *Miconia rostrata* (Naudin) Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 752 (1891). *Acinodendron rostratum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 952 (1891). *Heterotrichum rostratum* (Naudin) Gleason, Bull. Torrey Bot. Club 52: 378 (1925).

Section: Octomeris.

Distribution: Colombia, Venezuela.

Miconia tubulosa Gleason, Phytologia 1: 32 (1933). Type:—COLOMBIA. Lawrence 484 (Holotype: NY; isotypes: S, US, WIS).

Section: Tamonea.

Distribution: Colombia.

Miconia tuckeri Gleason, Madroño 8 (6): 190 (1946). Type:—EL SALVADOR. Tucker 998 (Holotype: UC; isotypes: F, K, LL, MICH, NY, PH, US).

Section: Cremanium.

Distribution: El Salvador.

Miconia tuerckheimii Cogn. in Smith, Bot. Gaz. 16: 5 (1891). *Acinodendron tuerckheimii* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—GUATEMALA. von Türckheim 581 (Holotype: BR; isotypes: K, NY, US).

Clidemia glandulifera Cogn. in Smith, Bot. Gaz. 20: 289 (1895).

Section: Octomeris.

Distribution: Mexico, Guatemala.

Miconia turgida Gleason, Bull. Torrey Bot. Club 52: 452 (1925). Type:—COLOMBIA. Pennell 9302 (Holotype: NY; isotypes: GH, PH, US).

Section: Chaenopleura.

Distribution: Colombia.

Miconia turquensis Urb. & Ekman, Repert. Spec. Nov. Regni Veg. 22: 228 (1926). Type:—CUBA. Ekman 14535 (Lectotype: S).

Section: Chaenopleura.

Distribution: Cuba.

Miconia ulmariooides Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 211 (1850). *Acinodendron ulmariooides* (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—VENEZUELA. Linden 396 (Holotype: P; isotypes: BR, G, MPU).

Miconia aegrotans Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 211 (1850). *Acinodendron aegrotans* (Naudin) Kuntze, Revis. Gen. Pl. 2: 950 (1891).

Miconia ulmariooides var. *fendleri* Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 873 (1891).

Section: Amblyarrhena.

Distribution: Colombia, Venezuela.

Miconia umbriensis Wurdack, Phytologia 18: 153 (1969). Type:—COLOMBIA. Klug 1926 (Holotype: US; isotypes: BM, K, MO, S).

Section: Miconia.

Distribution: Colombia, Panama.

Miconia umbrosa Cogn. in Martius, Fl. bras. 14 (4): 315 pl. 64 (1887). *Acinodendron umbrosum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. Riedel 1461 (Syntypes: BR, LE; isosyntypes: K, P).

Section: Miconia.

Distribution: Brazil North.

Miconia undata Triana, Trans. Linn. Soc. London 28: 107 (1871). *Acinodendron undatum* (Triana) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. Spruce 2000 (Holotype: K; isotypes: BR, K, M, TCD).

Miconia undata var. *robusta* Cogn., Bull. Torrey Bot. Club 23: 277 (1896).

Miconia undata var. *boliviensis* Cogn. in Rusby, Mem. Torrey Bot. Club 6: 39 (1896).

Section: Miconia.

Distribution: Venezuela, Bolivia, Brazil North.

Miconia uninervis Alain, Contr. Ocas. Mus. Hist. Nat. Colegio "De La Salle" 14: 4 (1955). Type:—CUBA. Alain 1008 (Holotype: HAC; isotype: NY).

Section: Amblyarrhena.

Distribution: Cuba.

Miconia urbaniana Cogn., Bot. Jahrb. Syst. 42: 139 (1909). Type:—PERU. *Weberbauer 5012* (Holotype: B, destroyed; isotype: BR).

Section: Miconia.

Distribution: Peru.

Miconia urceolata Urb., Symb. Antill. 9: 116 (1923). *Tetrazygia urceolata* (Urb.) Borhidi, Acta Bot. Hung. 23: 37 (1977). Type:—CUBA. *Ekman 9401* (Holotype: S; isotypes: A, F, K, NY).

Section: Tamonea.

Distribution: Cuba.

Miconia uribei Wurdack, Phytologia 50: 303 (1982). Type:—COLOMBIA. *Uribe-Uribe 6580* (Holotype: COL; isotypes: COL, US).

Section: Amblyarrhena.

Distribution: Colombia.

Miconia urophylla DC., Prodr. 3: 186 (1828). *Rhexia caudata* Schrank & Mart. ex DC., Prod. 3: 186 (1828), *pro syn.* *Acinodendron urophyllum* (DC.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. *Martius s.n.* (Holotype: M; isotype: G-DC).

Section: Chaenanthera.

Distribution: Brazil Southeast.

Miconia urticoides Triana, Trans. Linn. Soc. London 28: 123 (1871). *Acinodendron urticoides* (Triana) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—COLOMBIA. *Triana 6258/64* (Holotype: BM; isotype: COL).

Section: Amblyarrhena.

Distribution: Panama, Colombia.

Miconia uvida Wurdack, Phytologia 9: 418 (1964). Type:—PERU. *Wurdack 2153* (Holotype: US; isotypes: F, GH, K, MICH, NY, S, UC, USM).

Section: Miconia.

Distribution: Peru.

Miconia uvifera Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 222 (1850). *Acinodendron uviferum* (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BOLIVIA. *Weddell 4284* (Holotype: P; isotypes: BR, F).

Section: Cremanium.

Distribution: Bolivia.

Miconia vaccinioides (Bonpl.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 233 (1850).

Basionym: *Melastoma vaccinioides* Bonpl., Monogr. Melast. 1: 15 (1807). *Cremanium vaccinioides* (Bonpl.) D.Don, Mem. Wern. Nat. Hist. Soc. 4: 312 (1823). *Acinodendron vaccinioides* (Bonpl.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—PERU. *Bonpland s.n.* (Holotype: P; isotype: P).

Section: Cremanium.

Distribution: Ecuador, Peru.

Miconia valerioana (Standl.) Wurdack, Phytologia 31: 497 (1975).

Basionym: *Ossaea valerioana* Standl., Field Mus. Publ. Bot. 18 (3): 836 (1938). Type:—COSTA RICA. *Brenes 4525* (Holotype: F; isotype: NY).

Clidemia trichopoda Gleason, Brittonia 3: 133 (1939). *Miconia trichopoda* (Gleason) Wurdack, Phytologia 26: 4 (1973).

Section: Laceraria.

Distribution: Costa Rica, Panama.

Miconia valida Cogn., Bull. Torrey Bot. Club 23: 278 (1896). *Acinodendron validum* (Cogn.) Kuntze, Revis. Gen. Pl. 3 (2): 94 (1898). Type:—BOLIVIA. *Bang 2630* (Holotype: BR; isotype: NY).

Section: Cremanium.

Distribution: Bolivia.

Miconia vallensis Wurdack, Phytologia 8: 169 (1962). Type:—COLOMBIA. *Cuatrecasas* 21667 (Holotype: US; isotypes: BC, NY, U).
Section: Cremanium.
Distribution: Colombia.

Miconia valtheri Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 155 (1850). *Acinodendron valtheri* (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. *Vauthier* 109 (Holotype: P; isotype: US).
Miconia valtheri var. *saldanhae* Cogn. in Martius, Fl. bras. 14 (4): 362 (1887).
Section: Glossocentrum.
Distribution: Brazil Southeast, Brazil South.

Miconia vargasii Wurdack, Phytologia 23: 498 (1972). Type:—PERU. *Vargas* 2942 (Holotype: US).
Section: Cremanium.
Distribution: Peru.

Miconia velutina Triana, Trans. Linn. Soc. London 28: 113 (1871). *Acinodendron velutinum* (Triana) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—COLOMBIA. *Schlimgen* 545 (Holotype: BM; isotypes: BR, K, MPU, US).
Section: Amblyarrhena
Distribution: Colombia, Venezuela.

Miconia venulosa Wurdack, Phytologia 9: 420 (1964). Type:—PERU. *Wurdack* 2044 (Holotype: US, isotypes: S, UC, USM).
Section: Miconia.
Distribution: Ecuador, Peru.

Miconia verrucosa Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 920 (1891). *Acinodendron verrucosum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—COLOMBIA. *Schlimgen* 435 (Holotype: BR; isotypes: BM, BR, K, MPU).
Section: Cremanium.
Distribution: Colombia.

Miconia versicolor Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 189 (1850). *Acinodendron versicolor* (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—COLOMBIA. *Bonpland* s.n. (Lectotype: P; isolectotypes: G, P).
Miconia stricta Cogn., Bot. Jahrb. Syst. 8: 27 (1886). *Acinodendron strictum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891).
Section: Miconia.
Distribution: Colombia, Venezuela, Ecuador.

Miconia vesca Wurdack, Phytologia 41: 5 (1978). Type:—ECUADOR. *Steyermark* 53582 (Holotype: F; isotypes: NY, US).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia vestita Almeda, Proc. Calif. Acad. Sci., Ser. 4 52: 51 (2000). Type:—COSTA RICA. *Almeda* 7399 (Holotype: CAS; isotype: CR).
Section: Cremanium.
Distribution: Costa Rica.

Miconia victorinii Alain, Contr. Ocas. Mus. Hist. Nat. Colegio "De La Salle" 14: 5 (1955). Type:—CUBA. *Leon* 20670 (Holotype: HAC; isotype: NY).
Section: Octomeris.
Distribution: Cuba.

Miconia vilhenensis Wurdack, Phytologia 50: 299 (1982). Type:—BRAZIL. Vieira 626 (Holotype: MG; isotypes: NY, US).
Section: Glossocentrum.
Distribution: Bolivia, Brazil North, Brazil Central-West.

Miconia villonacensis Wurdack, Phytologia 39: 322 (1978). Type:—ECUADOR. Harling 13485 (Holotype: GB; isotypes: MO, S, US).
Section: Cremanium.
Distribution: Ecuador.

Miconia violacea Cogn., Bot. Jahrb. Syst. 8: 28 (1886). *Acinodendron violaceum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—COLOMBIA. Lehmann 3842 (Holotype: K; isotypes: BR, US).
Section: Cremanium.
Distribution: Colombia.

Miconia viscidula Urb. & Cogn., Symb. Antill. 7: 527 (1913). Type:—DOMINICAN REPUBLIC. Fuertes 1753 (Lectotype: NY; isolectotype: A).
Section: Chaenopleura.
Distribution: Dominican Republic.

Miconia vismiooides Triana, Trans. Linn. Soc. London 28: 124 (1871). *Acinodendron vismiooides* (Triana) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. *S.coll.*, s.n. (Holotype: BM; isotype: BR).
Section: Amblyarrhena.
Distribution: Brazil Southeast (probably).

Miconia vitiflora J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 187 (1929). Type:—PERU. Macbride 5165 (Holotype: F; isotypes: NY, US).
Section: Cremanium.
Distribution: Peru.

Miconia vittata (Linden & André) Cogn. in de Candolle & de Candolle, Monogr. Phan. 7: 729 (1891).
Basinonym: *Clidemia vittata* Linden & André, Ill. Hort. 22: 152 (1875). *Acinodendron vittatum* (Linden & André) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—PERU. *S.coll.*, s.n. (Holotype: BR).
Miconia capitata Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 360 (1915).
Section: Jucunda.
Distribution: Peru.

Miconia voronovii Gleason, Bull. Torrey Bot. Club 57: 66 (1930). Type:—COLOMBIA. Woronow 4529 (Holotype: LE).
Section: Tamonea.
Distribution: Colombia.

Miconia wagneri J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 4: 179 (1929). Type:—PERU. Macbride 5718 (Holotype: F; isotypes: NY, US).
Section: Laceraria.
Distribution: Peru, Bolivia, Brazil North.

Miconia walterjuddii Bécquer & Michelangeli, Brittonia 63: 425 (2011). Type:—CUBA. Bécquer HFC-85487 (Holotype: HAJB; isotypes: B, HAJB, NY).
Section: Chaenopleura.
Distribution: Cuba.

Miconia warmingiana Cogn. in Martius, Fl. bras. 14 (4): 282 (1887). *Acinodendron warmingianum* (Cogn.) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. Warming s.n. (Holotype: C; isotypes: BR, K, P).
Section: Glossocentrum.
Distribution: Brazil Southeast.

Miconia weberbaueri Cogn., Bot. Jahrb. Syst. 42: 144 (1909). Type:—PERU. *Weberbauer* 3396 (Holotype: B, destroyed; isotype: BR).
Section: Cremanium.
Distribution: Peru.

Miconia weddelli Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 144 (1850). *Acinodendron weddelli* (Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. *Weddell* 3001 (Holotype: P; isotype: BR).
Section: Miconia.
Distribution: Brazil Central-West.

Miconia willdenowii Klotzsch ex Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 199 (1850). *Acinodendron willdenowii* (Klotzsch ex Naudin) Kuntze, Revis. Gen. Pl. 2: 953 (1891). Type:—BRAZIL. *Sello* s.n. (Holotype: P; isotypes: K, US).
Section: Glossocentrum.
Distribution: Brazil Southeast, Brazil South.

Miconia wittii Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 367 (1915). *Miconia micrantha* Pilg., Verh. Bot. Vereins Prov. Brandenburg 47: 173 (1905), *nom. illegit.* Type:—BRAZIL. *Ule* 5090 (Holotype: B, destroyed; isotype: HBG).
Section: Miconia.
Distribution: Colombia, Venezuela, Brazil North.

Miconia wolfei Wurdack, Phytologia 23: 486 (1972). Type:—PERU. *Wolfe* 12429 (Holotype: US; isotype: NA).
Section: Amblyarrhena.
Distribution: Peru.

Miconia woytkowskii Wurdack, Phytologia 16: 178 (1968). Type:—PERU. *Woytkowski* 34407 (Holotype: UC; isotypes: F, LIL, S, UC, US).
Section: Miconia.
Distribution: Peru, Brazil North, Brazil West-Central.

Miconia wurdackii L.Uribe, Caldasia 10: 287 (1969). Type:—COLOMBIA. *Uribe-Uribe* 5683 (Holotype: COL; isotypes: COL, US).
Section: Cremanium.
Distribution: Colombia.

Miconia xenotricha Urb. & Ekman, Ark. Bot. 22A (17): 32 (1929). Type:—HAITI. *Ekman* H7447 (Holotype: S; isotypes: NY, S, US).
Section: Uncertain.
Distribution: Haiti.

Miconia yatuensis Wurdack, Mem. New York Bot. Gard. 10 (4): 33 (1961). Type:—VENEZUELA. *Maguire* 36695 (Holotype: NY; isotype: VEN).
Section: Miconia.
Distribution: Venezuela.

Miconia zamorensis Gleason, Phytologia 2: 301 (1947). Type:—ECUADOR. *Steyermark* 54701 (Holotype: F).
Section: Amblyarrhena.
Distribution: Ecuador.

Miconia zanonii Judd, Skean & R.S.Beaman, Brittonia 40: 208 (1988). Type:—DOMINICAN REPUBLIC. *Judd* 5158 (Holotype: FLAS; isotypes: A, F, JBSD, MO, NY, S, US).
Section: Chaenopleura.
Distribution: Dominican Republic.

Miconia zarucchii Wurdack, Phytologia 48: 244 (1981). Type:—COLOMBIA. *Zarucchi* 1632 (Holotype: US; isotypes: GH, HUA, K).

Section: Miconia.

Distribution: Colombia.

Miconia zemurrayana Standl. & L.O.Williams, Ceiba 1: 42 (1950). Type:—HONDURAS. *Williams* 15828 (Holotype: F; isotypes: EAP, US).

Tococa parvifolia Donn.Sm., Bot. Gaz. 27: 335 (1899).

Section: Jucunda.

Distribution: Mexico, Guatemala, El Salvador, Honduras.

Miconia zubenetana J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (4): 464 (1941). Type:—PERU. *Klug* 2208 (Holotype: F; isotypes: BM, K, NY, US).

Section: Miconia.

Distribution: Ecuador, Peru.

LIST 2. Species of *Miconia* organized by section.

Section Uncertain (7 spp)

- Miconia albiviridis*
Miconia alloeotricha
Miconia ancistrophora
Miconia carvalhoana
Miconia dura
Miconia johnwurdackiana
Miconia xenotricha

Miconia caesia

- Miconia cajanumana*

Miconia calignosa

Miconia calocoma

Miconia capitellata

Miconia carpishana

Miconia castrensis

Miconia cataractae

Miconia centrophora

Miconia centrosperma

Miconia clivorum

Miconia clypeata

Miconia codonostigma

Miconia collayensis

Miconia colliculosa

Miconia coloradensis

Miconia condylata

Miconia cookii

Miconia cordifolia

Miconia coronifera

Miconia costaricensis

Miconia crassifolia

Miconia crinita

Miconia cruenta

Miconia dapsiliflora

Miconia denticulata

Miconia difficilis

Miconia dissitiflora

Miconia dissitinervia

Miconia dunstervillei

Miconia erosa

Miconia espinosae

Miconia expansa

Miconia ferreyrae

Miconia floribunda

Miconia fosbergii

Miconia frontinoana

Miconia gibba

Miconia glabrata

Miconia glandulistylo

Miconia glutinosa

Miconia goniostigma

Miconia gossypina

Miconia grandiflora

Miconia grayana

Miconia hadrophylla

Miconia hamata

Miconia hematostemon

Miconia heterothrix

Miconia hexamera

Miconia hirta

Miconia huanucensis

Adenodesma (6 spp)

- Miconia biglandulosa*
Miconia plukenetii
Miconia prancei
Miconia rugosa
Miconia silicicola
Miconia tomentosa

Amblyarrhena (212 spp)

- Miconia acalephoides*
Miconia adrieni
Miconia aequatorialis
Miconia aggregata
Miconia alberti
Miconia alborosea
Miconia aligera
Miconia alypifolia
Miconia amabilis
Miconia amblyandra
Miconia andreana
Miconia androsaemifolia
Miconia anisophylla
Miconia annulata
Miconia antioquiensis
Miconia arbutifolia
Miconia asclepiadea
Miconia aspergillaris
Miconia asperrima
Miconia asplundii
Miconia atropilis
Miconia aymardii
Miconia bailloniana
Miconia bangii
Miconia barbeyana
Miconia barbipilis
Miconia bella
Miconia beneolens
Miconia benoistii
Miconia bipatrialis
Miconia bolivarensis
Miconia boxii
Miconia cacumina
Miconia caesariata

| | |
|-----------------------------------|----------------------------------|
| <i>Miconia huigrensis</i> | <i>Miconia pailasana</i> |
| <i>Miconia idiogena</i> | <i>Miconia papillosa</i> |
| <i>Miconia igniaria</i> | <i>Miconia paradoxa</i> |
| <i>Miconia imitans</i> | <i>Miconia pausana</i> |
| <i>Miconia inamoena</i> | <i>Miconia pedicellata</i> |
| <i>Miconia inanis</i> | <i>Miconia penduliflora</i> |
| <i>Miconia incachacana</i> | <i>Miconia penicillata</i> |
| <i>Miconia incurva</i> | <i>Miconia pergamentacea</i> |
| <i>Miconia innata</i> | <i>Miconia perobscura</i> |
| <i>Miconia koepckeana</i> | <i>Miconia phaeotricha</i> |
| <i>Miconia kuntzei</i> | <i>Miconia pichinchensis</i> |
| <i>Miconia lambayequeensis</i> | <i>Miconia pilaloensis</i> |
| <i>Miconia lamprarrhena</i> | <i>Miconia pisinna</i> |
| <i>Miconia larensis</i> | <i>Miconia plumifera</i> |
| <i>Miconia lasiocalyx</i> | <i>Miconia polytopica</i> |
| <i>Miconia lasiostyla</i> | <i>Miconia poortmannii</i> |
| <i>Miconia laxa</i> | <i>Miconia prasinifolia</i> |
| <i>Miconia leandroides</i> | <i>Miconia prominens</i> |
| <i>Miconia lennartanderssonii</i> | <i>Miconia protuberans</i> |
| <i>Miconia licrophora</i> | <i>Miconia pseudocentrophora</i> |
| <i>Miconia lithophila</i> | <i>Miconia pseudoradula</i> |
| <i>Miconia livida</i> | <i>Miconia psychrophila</i> |
| <i>Miconia longisepala</i> | <i>Miconia pulgari</i> |
| <i>Miconia loreyoides</i> | <i>Miconia punicea</i> |
| <i>Miconia lucida</i> | <i>Miconia pustulata</i> |
| <i>Miconia lutescens</i> | <i>Miconia quintuplinervia</i> |
| <i>Miconia luteynii</i> | <i>Miconia radula</i> |
| <i>Miconia macbrydeana</i> | <i>Miconia ramboi</i> |
| <i>Miconia macrantha</i> | <i>Miconia reflexipila</i> |
| <i>Miconia madisonii</i> | <i>Miconia renneri</i> |
| <i>Miconia madrensis</i> | <i>Miconia reticulata</i> |
| <i>Miconia majalis</i> | <i>Miconia retusa</i> |
| <i>Miconia mapirensis</i> | <i>Miconia rhodantha</i> |
| <i>Miconia mazatecana</i> | <i>Miconia rimbachii</i> |
| <i>Miconia mcvaughii</i> | <i>Miconia rivetii</i> |
| <i>Miconia militis</i> | <i>Miconia roraimensis</i> |
| <i>Miconia modica</i> | <i>Miconia rubricans</i> |
| <i>Miconia monzonensis</i> | <i>Miconia ruizii</i> |
| <i>Miconia mulleola</i> | <i>Miconia saltuensis</i> |
| <i>Miconia multiplinervia</i> | <i>Miconia sanguinea</i> |
| <i>Miconia myrtillifolia</i> | <i>Miconia scabra</i> |
| <i>Miconia namandensis</i> | <i>Miconia schlimii</i> |
| <i>Miconia nasella</i> | <i>Miconia scutata</i> |
| <i>Miconia neurotricha</i> | <i>Miconia seposita</i> |
| <i>Miconia nitidissima</i> | <i>Miconia setosa</i> |
| <i>Miconia nubicola</i> | <i>Miconia shattuckii</i> |
| <i>Miconia obovata</i> | <i>Miconia sneidernii</i> |
| <i>Miconia obscura</i> | <i>Miconia sodiroi</i> |
| <i>Miconia ombrophila</i> | <i>Miconia sordida</i> |
| <i>Miconia osaensis</i> | <i>Miconia spatellophora</i> |
| <i>Miconia pachyphylla</i> | <i>Miconia spinulosa</i> |
| <i>Miconia paeminoa</i> | <i>Miconia spireifolia</i> |

Miconia staphidioides
Miconia stipularis
Miconia subalpina
Miconia subglabra
Miconia suborbicularis
Miconia tacanensis
Miconia tachirensis
Miconia teotepecensis
Miconia tepicana
Miconia terborghii
Miconia terera
Miconia tiri
Miconia trujillensis
Miconia ulmariooides
Miconia uninervis
Miconia uribei
Miconia urticoides
Miconia velutina
Miconia vesca
Miconia vismiooides
Miconia wolfei
Miconia zamorensis

Chaenanthera (17 spp)

Miconia chrysophylla
Miconia dolichorrhyncha
Miconia elaeodendron
Miconia hypoleuca
Miconia inaequidens
Miconia latecrenata
Miconia picinguabensis
Miconia pisinniflora
Miconia pusilliflora
Miconia regelii
Miconia rimalis
Miconia sellowiana
Miconia tentaculifera
Miconia tenuis
Miconia tetrazygioides
Miconia trianae
Miconia urophylla

Chaenopleura (86 spp)

Miconia adenocalyx
Miconia alainii
Miconia alpina
Miconia alternifolia
Miconia apiculata
Miconia ascendens
Miconia barkeri
Miconia basilensis
Miconia bifaria
Miconia bullata
Miconia calycina
Miconia campanensis
Miconia campii
Miconia cauingia
Miconia chionophila
Miconia concinna
Miconia confertiflora
Miconia coniophora
Miconia corazonica
Miconia corymbiformis
Miconia cubensis
Miconia dielsiana
Miconia domingensis
Miconia favosa
Miconia ferruginea
Miconia fictilis
Miconia firma
Miconia flavescens
Miconia foveolata
Miconia fruticulosa
Miconia fuertesii
Miconia grandidentata
Miconia griffisii
Miconia grisea
Miconia harlingii
Miconia haughtii
Miconia howardiana
Miconia hymenanthera
Miconia hypoides
Miconia jimenezii
Miconia krugii
Miconia lachnoclada
Miconia lanceolata
Miconia latifolia
Miconia longibracteata
Miconia longisetosa
Miconia luteola
Miconia macayana
Miconia mansfeldiana
Miconia markgrafii
Miconia melanotricha
Miconia moensis
Miconia multiglandulosa
Miconia nigricans
Miconia nitida
Miconia ottikeri
Miconia popayanensis
Miconia pseudorigida
Miconia pteroclada
Miconia pycnoneura
Miconia quadrangularis

| | |
|--------------------------------|---------------------------------|
| <i>Miconia rigida</i> | <i>Miconia brenesii</i> |
| <i>Miconia rigidissima</i> | <i>Miconia brevis</i> |
| <i>Miconia rotundifolia</i> | <i>Miconia brevistylis</i> |
| <i>Miconia rufa</i> | <i>Miconia brevitheca</i> |
| <i>Miconia salicifolia</i> | <i>Miconia brittonii</i> |
| <i>Miconia samanensis</i> | <i>Miconia buxifolia</i> |
| <i>Miconia santanana</i> | <i>Miconia caelata</i> |
| <i>Miconia secundifolia</i> | <i>Miconia caerulea</i> |
| <i>Miconia selleana</i> | <i>Miconia carnea</i> |
| <i>Miconia septentrionalis</i> | <i>Miconia castillensis</i> |
| <i>Miconia silverstonei</i> | <i>Miconia cernua</i> |
| <i>Miconia sintenisii</i> | <i>Miconia chiriquiensis</i> |
| <i>Miconia skeaniana</i> | <i>Miconia choriophylla</i> |
| <i>Miconia sphagnicola</i> | <i>Miconia chrysocoma</i> |
| <i>Miconia stenobotrys</i> | <i>Miconia cladonia</i> |
| <i>Miconia stenophylla</i> | <i>Miconia clathrantha</i> |
| <i>Miconia subcompressa</i> | <i>Miconia cleefii</i> |
| <i>Miconia superposita</i> | <i>Miconia cocoensis</i> |
| <i>Miconia tephrodes</i> | <i>Miconia coelestis</i> |
| <i>Miconia thrysiflora</i> | <i>Miconia comosa</i> |
| <i>Miconia turgida</i> | <i>Miconia compressicaulis</i> |
| <i>Miconia turquinensis</i> | <i>Miconia coriacea</i> |
| <i>Miconia viscidula</i> | <i>Miconia cosangensis</i> |
| <i>Miconia walterjuddii</i> | <i>Miconia crassistigma</i> |
| <i>Miconia zanonii</i> | <i>Miconia crebribullata</i> |
| Cremanium (223 spp) | |
| <i>Miconia acanthocoryne</i> | <i>Miconia cremadena</i> |
| <i>Miconia aenigmatica</i> | <i>Miconia cremophylla</i> |
| <i>Miconia aguirrei</i> | <i>Miconia crocata</i> |
| <i>Miconia alpestris</i> | <i>Miconia crocea</i> |
| <i>Miconia amilcariana</i> | <i>Miconia cuatrecasae</i> |
| <i>Miconia amplinodis</i> | <i>Miconia cundinamarcensis</i> |
| <i>Miconia angelana</i> | <i>Miconia cuprea</i> |
| <i>Miconia aprica</i> | <i>Miconia cuspidatissima</i> |
| <i>Miconia aspratilis</i> | <i>Miconia cyanocarpa</i> |
| <i>Miconia atrofusca</i> | <i>Miconia cyathantha</i> |
| <i>Miconia auritinoda</i> | <i>Miconia danielii</i> |
| <i>Miconia ayacuchensis</i> | <i>Miconia dasyclada</i> |
| <i>Miconia barclayana</i> | <i>Miconia demissifolia</i> |
| <i>Miconia bernardii</i> | <i>Miconia densifolia</i> |
| <i>Miconia berryi</i> | <i>Miconia desportesii</i> |
| <i>Miconia biacuta</i> | <i>Miconia dichroa</i> |
| <i>Miconia biappendiculata</i> | <i>Miconia diegogomezii</i> |
| <i>Miconia biformis</i> | <i>Miconia dielsii</i> |
| <i>Miconia bilopezii</i> | <i>Miconia dioica</i> |
| <i>Miconia biperulifera</i> | <i>Miconia dissimulans</i> |
| <i>Miconia boliviensis</i> | <i>Miconia dissita</i> |
| <i>Miconia bordoncilloana</i> | <i>Miconia divergens</i> |
| <i>Miconia brachyanthera</i> | <i>Miconia dodsonii</i> |
| <i>Miconia brachygyna</i> | <i>Miconia dolichopoda</i> |
| <i>Miconia bracteolata</i> | <i>Miconia dumetosa</i> |
| | <i>Miconia egregia</i> |
| | <i>Miconia elaeoides</i> |

- Miconia elongata*
Miconia elvirae
Miconia floccosa
Miconia fuliginosa
Miconia galactantha
Miconia glaberrima
Miconia gleasoniana
Miconia globuliflora
Miconia glomerulifera
Miconia glyptophylla
Miconia gonioclada
Miconia hemenostigma
Miconia herzogii
Miconia hildeana
Miconia hirsutivena
Miconia hirtella
Miconia hutchisonii
Miconia hyemalis
Miconia hygrophila
Miconia inconspicua
Miconia insueta
Miconia ioneura
Miconia jahnii
Miconia jentacularum
Miconia jorgensenii
Miconia kappellei
Miconia laciniata
Miconia laetivirens
Miconia latistigma
Miconia laurina
Miconia ledifolia
Miconia lehmannii
Miconia ligustrina
Miconia lilacina
Miconia limitaris
Miconia loxensis
Miconia lugubris
Miconia lymanii
Miconia machinazana
Miconia mandonii
Miconia manicata
Miconia martinicensis
Miconia media
Miconia mediocris
Miconia medusa
Miconia mesmeana
Miconia micayana
Miconia micrantha
Miconia micropetala
Miconia milesmorganii
Miconia mimica
Miconia minuta

Miconia miocarpa
Miconia molinopampana
Miconia monciona
Miconia mornicola
Miconia mutisiana
Miconia myrtiformis
Miconia neblinensis
Miconia nerifolia
Miconia nodosa
Miconia nystroemii
Miconia ochracea
Miconia oellgaardii
Miconia oligotricha
Miconia onaensis
Miconia opacifolia
Miconia orcheotoma
Miconia oreogena
Miconia orescia
Miconia paludigena
Miconia paradisica
Miconia parvifolia
Miconia pastoensis
Miconia pavoniana
Miconia penningtonii
Miconia perijensis
Miconia pernettifolia
Miconia peruviana
Miconia petersonii
Miconia pittieri
Miconia plena
Miconia plethorica
Miconia polychaeta
Miconia polygama
Miconia polyneura
Miconia prietoi
Miconia pulverulenta
Miconia punctibullata
Miconia puracensis
Miconia purulensis
Miconia quadrialata
Miconia quadrifolia
Miconia quadripora
Miconia rava
Miconia reburrosa
Miconia reclinata
Miconia recondita
Miconia resima
Miconia resimoides
Miconia rhonhofiae
Miconia ricardoi
Miconia rigens
Miconia rubens

| | |
|-------------------------------|--------------------------------|
| <i>Miconia rufiramea</i> | <i>Miconia brunnea</i> |
| <i>Miconia rugifolia</i> | <i>Miconia buddlejoides</i> |
| <i>Miconia ruizteranii</i> | <i>Miconia cabucu</i> |
| <i>Miconia rupestris</i> | <i>Miconia castaneiflora</i> |
| <i>Miconia rusbyana</i> | <i>Miconia chartacea</i> |
| <i>Miconia rzedowskii</i> | <i>Miconia chlorocarpa</i> |
| <i>Miconia sandemanii</i> | <i>Miconia cinerascens</i> |
| <i>Miconia saxatilis</i> | <i>Miconia cinnamomifolia</i> |
| <i>Miconia schnellii</i> | <i>Miconia collatata</i> |
| <i>Miconia setulosa</i> | <i>Miconia corallina</i> |
| <i>Miconia shaferi</i> | <i>Miconia cubatanensis</i> |
| <i>Miconia smithii</i> | <i>Miconia discolor</i> |
| <i>Miconia spinulidentata</i> | <i>Miconia divaricata</i> |
| <i>Miconia squamulosa</i> | <i>Miconia dorsaliporosa</i> |
| <i>Miconia sterilis</i> | <i>Miconia egensis</i> |
| <i>Miconia sulcata</i> | <i>Miconia eichleri</i> |
| <i>Miconia summa</i> | <i>Miconia ernstii</i> |
| <i>Miconia tabayensis</i> | <i>Miconia fasciculata</i> |
| <i>Miconia talamancensis</i> | <i>Miconia flammea</i> |
| <i>Miconia tamana</i> | <i>Miconia fluminensis</i> |
| <i>Miconia tetrandra</i> | <i>Miconia formosa</i> |
| <i>Miconia tetrastoma</i> | <i>Miconia gigantea</i> |
| <i>Miconia thaminantha</i> | <i>Miconia gilva</i> |
| <i>Miconia theizans</i> | <i>Miconia guaiquinimae</i> |
| <i>Miconia tinifolia</i> | <i>Miconia herrerae</i> |
| <i>Miconia tonduzii</i> | <i>Miconia icosandra</i> |
| <i>Miconia tovarensis</i> | <i>Miconia klotzschii</i> |
| <i>Miconia tricaudata</i> | <i>Miconia klugii</i> |
| <i>Miconia trichogona</i> | <i>Miconia kollmannii</i> |
| <i>Miconia tuckeri</i> | <i>Miconia kriegeriana</i> |
| <i>Miconia uvifera</i> | <i>Miconia laeta</i> |
| <i>Miconia vaccinoides</i> | <i>Miconia lagunensis</i> |
| <i>Miconia valida</i> | <i>Miconia leptantha</i> |
| <i>Miconia vallensis</i> | <i>Miconia ligustroides</i> |
| <i>Miconia vargasii</i> | <i>Miconia longicuspis</i> |
| <i>Miconia verrucosa</i> | <i>Miconia longifolia</i> |
| <i>Miconia vestita</i> | <i>Miconia lurida</i> |
| <i>Miconia villonacensis</i> | <i>Miconia mattogrossensis</i> |
| <i>Miconia violacea</i> | <i>Miconia maximowicziana</i> |
| <i>Miconia vitiflora</i> | <i>Miconia mellina</i> |
| <i>Miconia weberbaueri</i> | <i>Miconia minutiflora</i> |
| <i>Miconia wurdackii</i> | <i>Miconia molesta</i> |
| Glossocentrum (82 spp) | |
| <i>Miconia acunae</i> | <i>Miconia myriantha</i> |
| <i>Miconia alborufescens</i> | <i>Miconia oblongifolia</i> |
| <i>Miconia altissima</i> | <i>Miconia onychocalyx</i> |
| <i>Miconia argyraea</i> | <i>Miconia ossaeifolia</i> |
| <i>Miconia augusti</i> | <i>Miconia ovalifolia</i> |
| <i>Miconia baumgratziana</i> | <i>Miconia paniculata</i> |
| <i>Miconia bisulcata</i> | <i>Miconia pastazana</i> |
| <i>Miconia brasiliensis</i> | <i>Miconia pepericarpa</i> |
| | <i>Miconia perturbata</i> |
| | <i>Miconia petroniana</i> |

Miconia petropolitana
Miconia pseudoeichleri
Miconia puberula
Miconia rabenii
Miconia racemifera
Miconia remotiflora
Miconia ruschiana
Miconia saldanhae
Miconia sclerophylla
Miconia secunda
Miconia semisterilis
Miconia setosociliata
Miconia shepherdii
Miconia subvernucosa
Miconia ternatifolia
Miconia tetragona
Miconia trichotoma
Miconia tristis
Miconia valtheri
Miconia vilhenensis
Miconia warmingiana
Miconia willdenowii

Hartigia (7 spp)

Miconia ciliata
Miconia lacera
Miconia oinochrophylla
Miconia pulvinata
Miconia racemosa
Miconia schippii
Miconia toroi

Jucunda (23 spp)

Miconia acuminata
Miconia amacurensis
Miconia ampla
Miconia boomii
Miconia comptifolia
Miconia correae
Miconia decurrens
Miconia fissa
Miconia francavillana
Miconia gratissima
Miconia holosericea
Miconia jucunda
Miconia lasseri
Miconia melinonis
Miconia oldemanii
Miconia pachydonta
Miconia paucidens
Miconia rimachii
Miconia staminea

Miconia stellulata
Miconia truncata
Miconia vittata
Miconia zemurrayana

Laceraria (19 spp)

Miconia aliquantula
Miconia centrodesma
Miconia centrodesmoides
Miconia cornifolia
Miconia cuneata
Miconia diaphanea
Miconia divisoriana
Miconia dorsiloba
Miconia duckei
Miconia fanshawei
Miconia mariae
Miconia mexicana
Miconia pujana
Miconia rupticalyx
Miconia sastrei
Miconia seticaulis
Miconia striata
Miconia valerioana
Miconia wagneri

Miconia (245 spp)

Miconia abbreviata
Miconia abeggii
Miconia abysmophila
Miconia acinodendron
Miconia acutipetala
Miconia adinantha
Miconia aeruginosa
Miconia affinis
Miconia alata
Miconia albicans
Miconia alternans
Miconia amissa
Miconia amnicola
Miconia amoena
Miconia amplexicaulis
Miconia andersonii
Miconia apostachya
Miconia appendiculata
Miconia arboricola
Miconia argentea
Miconia argyrophylla
Miconia aristata
Miconia astroplocama
Miconia astrotricha
Miconia aulocalyx

- Miconia barbinervis*
Miconia biglomerata
Miconia borjensis
Miconia brachybotrya
Miconia bracteata
Miconia brevipes
Miconia burchellii
Miconia calvescens
Miconia campestris
Miconia cannabina
Miconia capixaba
Miconia carassana
Miconia caucana
Miconia caudigera
Miconia cautis
Miconia cazaletii
Miconia ceramicarpa
Miconia chaetodon
Miconia chamissois
Miconia chrysoneura
Miconia cinerea
Miconia cionotricha
Miconia cipoensis
Miconia commutata
Miconia compressa
Miconia conformis
Miconia cordata
Miconia cowanii
Miconia crassinervia
Miconia crassipes
Miconia cretacea
Miconia curta
Miconia cuspidata
Miconia cutucuensis
Miconia decipiens
Miconia desmantha
Miconia dichrophylla
Miconia dipsacea
Miconia dispar
Miconia dudleyi
Miconia elata
Miconia elegans
Miconia emendata
Miconia eriocalyx
Miconia erioclada
Miconia eriodonta
Miconia eugeniooides
Miconia falcata
Miconia fallax
Miconia ferruginata
Miconia filamentosa
Miconia fosteri
Miconia fragilis
Miconia fragrans
Miconia friedmaniorum
Miconia fulvostellata
Miconia glaucescens
Miconia glazioviana
Miconia glomerata
Miconia goudotii
Miconia gracilis
Miconia grandifoliata
Miconia grayumii
Miconia grossidentata
Miconia guayaquilensis
Miconia heliotropoides
Miconia herpetica
Miconia heterochaeta
Miconia heteromera
Miconia heterotricha
Miconia histothrix
Miconia hookeriana
Miconia hyperprasina
Miconia ibaguensis
Miconia ibarreae
Miconia idroboi
Miconia impetiolaris
Miconia inaequalifolia
Miconia insularis
Miconia irwinii
Miconia iteophylla
Miconia jefensis
Miconia kappleri
Miconia kavanayensis
Miconia laevigata
Miconia lamprophylla
Miconia lanata
Miconia lappacea
Miconia lateriflora
Miconia lechleri
Miconia lepidota
Miconia leucocarpa
Miconia liebmannii
Miconia ligulata
Miconia lonchophylla
Miconia longiracemosa
Miconia longicupidata
Miconia longispicata
Miconia loretensis
Miconia lourteigiana
Miconia lugonis
Miconia macrothyrsa
Miconia magdalenaе
Miconia maguirei

| | |
|----------------------------------|--------------------------------|
| <i>Miconia malatestae</i> | <i>Miconia pyrifolia</i> |
| <i>Miconia manauara</i> | <i>Miconia radulaefolia</i> |
| <i>Miconia marginata</i> | <i>Miconia rhytidophylla</i> |
| <i>Miconia maroana</i> | <i>Miconia riparia</i> |
| <i>Miconia martiniana</i> | <i>Miconia rivalis</i> |
| <i>Miconia matthaei</i> | <i>Miconia robusta</i> |
| <i>Miconia membranacea</i> | <i>Miconia rubiginosa</i> |
| <i>Miconia mendoncae</i> | <i>Miconia rufescens</i> |
| <i>Miconia mocquerysii</i> | <i>Miconia ruficalyx</i> |
| <i>Miconia mollicula</i> | <i>Miconia rufostellulata</i> |
| <i>Miconia mollis</i> | <i>Miconia sagotiana</i> |
| <i>Miconia molybdea</i> | <i>Miconia santaritensis</i> |
| <i>Miconia mourae</i> | <i>Miconia sarmentosa</i> |
| <i>Miconia multiflora</i> | <i>Miconia savannarum</i> |
| <i>Miconia multispicata</i> | <i>Miconia schlechtendalii</i> |
| <i>Miconia mutabilis</i> | <i>Miconia schunkei</i> |
| <i>Miconia nambyquarae</i> | <i>Miconia schwackei</i> |
| <i>Miconia navioensis</i> | <i>Miconia secundiflora</i> |
| <i>Miconia nematophora</i> | <i>Miconia serialis</i> |
| <i>Miconia nervosa</i> | <i>Miconia sessilifolia</i> |
| <i>Miconia nobilis</i> | <i>Miconia smaragdina</i> |
| <i>Miconia oligantha</i> | <i>Miconia spennerostachya</i> |
| <i>Miconia oligocephala</i> | <i>Miconia spicellata</i> |
| <i>Miconia oraria</i> | <i>Miconia spichigeri</i> |
| <i>Miconia organensis</i> | <i>Miconia splendens</i> |
| <i>Miconia ovata</i> | <i>Miconia sprucei</i> |
| <i>Miconia pandurata</i> | <i>Miconia stelligera</i> |
| <i>Miconia paraguayensis</i> | <i>Miconia stenocardia</i> |
| <i>Miconia paspaloides</i> | <i>Miconia stenostachya</i> |
| <i>Miconia peltata</i> | <i>Miconia stephananthera</i> |
| <i>Miconia pendula</i> | <i>Miconia stevensiana</i> |
| <i>Miconia pennipilis</i> | <i>Miconia steyermarkii</i> |
| <i>Miconia pentlandii</i> | <i>Miconia stipitata</i> |
| <i>Miconia persicariaefolia</i> | <i>Miconia strigosa</i> |
| <i>Miconia phaeophylla</i> | <i>Miconia subsessilifolia</i> |
| <i>Miconia phanerostila</i> | <i>Miconia subsimplex</i> |
| <i>Miconia phlebodes</i> | <i>Miconia subspicata</i> |
| <i>Miconia piperifolia</i> | <i>Miconia subulipetala</i> |
| <i>Miconia platypoda</i> | <i>Miconia sylvatica</i> |
| <i>Miconia poeppigii</i> | <i>Miconia symplocoidea</i> |
| <i>Miconia polita</i> | <i>Miconia tenensis</i> |
| <i>Miconia polyandra</i> | <i>Miconia tetrasperma</i> |
| <i>Miconia pozensis</i> | <i>Miconia tetraspermoides</i> |
| <i>Miconia prasina</i> | <i>Miconia thysanophylla</i> |
| <i>Miconia procumbens</i> | <i>Miconia tiliifolia</i> |
| <i>Miconia pseudoaplostachya</i> | <i>Miconia tillettii</i> |
| <i>Miconia pseudocapsularis</i> | <i>Miconia trichophora</i> |
| <i>Miconia pterocaulon</i> | <i>Miconia trimera</i> |
| <i>Miconia pubicalyx</i> | <i>Miconia trinervia</i> |
| <i>Miconia pulchra</i> | <i>Miconia triplinervis</i> |
| <i>Miconia punctata</i> | <i>Miconia tschudyooides</i> |
| <i>Miconia pyramidalis</i> | <i>Miconia umbriensis</i> |

| | |
|------------------------------|-------------------------------|
| <i>Miconia umbrosa</i> | <i>Miconia notabilis</i> |
| <i>Miconia undata</i> | <i>Miconia novemnervia</i> |
| <i>Miconia urbaniana</i> | <i>Miconia obconica</i> |
| <i>Miconia uvida</i> | <i>Miconia obtusa</i> |
| <i>Miconia venulosa</i> | <i>Miconia octopetala</i> |
| <i>Miconia versicolor</i> | <i>Miconia paupercula</i> |
| <i>Miconia weddellii</i> | <i>Miconia poecilantha</i> |
| <i>Miconia wittii</i> | <i>Miconia porphyrotricha</i> |
| <i>Miconia wojtkowskii</i> | <i>Miconia ravenii</i> |
| <i>Miconia yatuensis</i> | <i>Miconia rosea</i> |
| <i>Miconia zarucchii</i> | <i>Miconia russea</i> |
| <i>Miconia zubenetana</i> | <i>Miconia salebrosa</i> |
| Octomeris (59 spp) | <i>Miconia santaremensis</i> |
| <i>Miconia anisotricha</i> | <i>Miconia saxicola</i> |
| <i>Miconia araguensis</i> | <i>Miconia speciosa</i> |
| <i>Miconia avia</i> | <i>Miconia suaveolens</i> |
| <i>Miconia axinaeoides</i> | <i>Miconia thomasiana</i> |
| <i>Miconia baracoensis</i> | <i>Miconia tixicensis</i> |
| <i>Miconia benthamiana</i> | <i>Miconia tuberculata</i> |
| <i>Miconia brachycalyx</i> | <i>Miconia tuerckheimii</i> |
| <i>Miconia breteleri</i> | <i>Miconia victorinii</i> |
| <i>Miconia bucherae</i> | Tamonea (71 spp) |
| <i>Miconia canaguensis</i> | <i>Miconia acuminifera</i> |
| <i>Miconia celaquensis</i> | <i>Miconia acutifolia</i> |
| <i>Miconia centronioides</i> | <i>Miconia aguitensis</i> |
| <i>Miconia cerasiflora</i> | <i>Miconia amambayensis</i> |
| <i>Miconia chinantlana</i> | <i>Miconia amazonica</i> |
| <i>Miconia contrerasii</i> | <i>Miconia aponeura</i> |
| <i>Miconia delicatula</i> | <i>Miconia archeri</i> |
| <i>Miconia explicita</i> | <i>Miconia aurea</i> |
| <i>Miconia funckii</i> | <i>Miconia aureoides</i> |
| <i>Miconia gentryi</i> | <i>Miconia blakeifolia</i> |
| <i>Miconia globulifera</i> | <i>Miconia bubalina</i> |
| <i>Miconia guatemalensis</i> | <i>Miconia cacatin</i> |
| <i>Miconia hexapetala</i> | <i>Miconia caudata</i> |
| <i>Miconia imbricata</i> | <i>Miconia cercophora</i> |
| <i>Miconia ingens</i> | <i>Miconia chapensis</i> |
| <i>Miconia javorkaeana</i> | <i>Miconia ciliaris</i> |
| <i>Miconia jitotolana</i> | <i>Miconia cinchonifolia</i> |
| <i>Miconia labiakiana</i> | <i>Miconia coronata</i> |
| <i>Miconia laevipilis</i> | <i>Miconia dodecandra</i> |
| <i>Miconia leiotricha</i> | <i>Miconia donaeana</i> |
| <i>Miconia lenticellata</i> | <i>Miconia eremita</i> |
| <i>Miconia luciana</i> | <i>Miconia flaccida</i> |
| <i>Miconia lundellii</i> | <i>Miconia furfuracea</i> |
| <i>Miconia macrodon</i> | <i>Miconia glandulifera</i> |
| <i>Miconia megalantha</i> | <i>Miconia hondurensis</i> |
| <i>Miconia meridensis</i> | <i>Miconia hospitalis</i> |
| <i>Miconia moorei</i> | <i>Miconia huberi</i> |
| <i>Miconia multinervia</i> | <i>Miconia hylophila</i> |
| <i>Miconia nigripes</i> | <i>Miconia iluensis</i> |

- Miconia intricata*
Miconia juruensis
Miconia kraenzlinii
Miconia liesneri
Miconia littlei
Miconia macrotis
Miconia mazanana
Miconia metallica
Miconia michelangeliana
Miconia mirabilis
Miconia mituana
Miconia morii
Miconia muricata
Miconia napoana
Miconia nutans
Miconia obliqua
Miconia paleacea
Miconia perelegans
Miconia phaeochaeta
Miconia pubipetala
Miconia reducens
- Miconia rhombifolia*
Miconia robinsoniana
Miconia rufipila
Miconia sanctiphilippi
Miconia sciurea
Miconia serrulata
Miconia simplex
Miconia sparrei
Miconia steinbachii
Miconia stenourea
Miconia subandicola
Miconia subcordata
Miconia submacrophylla
Miconia superba
Miconia titanophylla
Miconia traillii
Miconia transversa
Miconia triangularis
Miconia tubulosa
Miconia urceolata
Miconia voronovii

LIST 3. Species of *Miconia* by country, with respective vouchers.

Mexico (79 spp)

- Miconia aeruginosa* (Breedlove 49887, CAS)
Miconia affinis (Cowan 3070, CAS)
Miconia albicans (Breedlove 57791, CAS)
Miconia ampla (Croat 63258, CAS)
Miconia anisotricha (Breedlove 45894, CAS)
Miconia argentea (Cowan 2592, CAS)
Miconia bubalina (Wendt 2672, CAS)
Miconia calvescens (Breedlove 58010, CAS)
Miconia caudata (Breedlove 56910, CAS)
Miconia chamissois (Breedlove 57803, CAS)
Miconia chinantlana (Breedlove 52655, CAS)
Miconia chrysoneura (Breedlove 64741, CAS)
Miconia chrysophylla (Breedlove 57693, CAS)
Miconia ciliata (Martínez 7059, CAS)
Miconia costaricensis (Breedlove 56748, CAS)
Miconia decurrens (Breedlove 57649, CAS)
Miconia desmantha (Breedlove 57601, CAS)
Miconia dodecandra (Breedlove 57714, CAS)
Miconia elata (Breedlove 57611, CAS)
Miconia fulvostellata (Breedlove 58054, CAS)
Miconia glaberrima (Breedlove 68645, CAS)
Miconia glabrata (Purpus 6783, US)
Miconia globulifera (Seler 2938, DS)
Miconia gracilis (Wendt 3080, CAS)
Miconia guatemalensis (Breedlove 48187, CAS)
Miconia hemenostigma (Breedlove 64644, CAS)
Miconia heterothrix (Breedlove 64492, CAS)
Miconia holosericea (Martínez 18858, CAS)
Miconia hondurensis (Martínez 18398, CAS)
Miconia hyperprasina (Breedlove 56794, CAS)
Miconia ibaguensis (Breedlove 58067, CAS)
Miconia ibarreae (Ibarra 3374, CAS)
Miconia impetiolaris (Breedlove 57218, CAS)
Miconia insularis (Nee 30015, CAS)
Miconia jititolana (Breedlove 49303, CAS)
Miconia lacera (Martínez 9276, CAS)
Miconia laevigata (Breedlove 60181, CAS)
Miconia lateriflora (Martínez 13496, CAS)
Miconia liebmannii (Martin 621, CAS)
Miconia livida (Breedlove 49357, CAS)
Miconia lonchophylla (Croat 65557, CAS)
Miconia longifolia (Valdivia 2261, CAS)
Miconia madrensis (Breedlove 45106, CAS)
Miconia matthaei (Breedlove 57651, CAS)
Miconia mazatecana (de Santiago 2076, MEXU)
Miconia mcvaughii (Boutin 2583, CAS)
Miconia mexicana (Daniel 8400, CAS)
Miconia militis (Breedlove 65022, CAS)
Miconia minutiflora (Breedlove 57856, CAS)
Miconia mirabilis (Cowan 4931, CAS)
Miconia moorei (Breedlove 59660, CAS)

Miconia nutans (Breedlove 57502, CAS)

- Miconia obconica* (López 641, CAS)
Miconia oinochrophylla (Martínez 13531, CAS)
Miconia oligocephala (Breedlove 55149, CAS)
Miconia oligotricha (Hernández 5634, CAS)
Miconia phaeotricha (Lorenz 4219, CAS)
Miconia prasina (Breedlove 57855, CAS)
Miconia punctata (Hernández PV-20, CAS)
Miconia purulensis (Breedlove 56718, CAS)
Miconia ravenii (Breedlove 57058, CAS)
Miconia reducens (Martínez 15460, CAS)
Miconia rzedowskii (Tenorio 2659, CAS)
Miconia saxicola (Breedlove 45139, CAS)
Miconia schippii (Martínez 13497, CAS)
Miconia schlechtendalii (Ventura 648, CAS)
Miconia serrulata (Calzada 4092, CAS)
Miconia smaragdina (Breedlove 58445, CAS)
Miconia splendens (Boege 1120, DS)
Miconia stenostachya (Breedlove 57792, CAS)
Miconia sylvatica (Breedlove 25361, CAS)
Miconia tacanensis (Breedlove 66149, CAS)
Miconia teotepicensis (Tenorio 1364, MEXU)
Miconia tepicana (González 414, CAS)
Miconia tomentosa (Martínez 18301, CAS)
Miconia trinervia (Wendt 3167, CAS)
Miconia triplinervis (Breedlove 57337, CAS)
Miconia tuerckheimii (Breedlove 60000, CAS)
Miconia zemurrayana (Daniel 8409, CAS)

Belize (46 spp)

- Miconia affinis* (Davidse 36736A, CAS)
Miconia albicans (Davidse 37019, CAS)
Miconia ampla (Gentle 4804, CAS)
Miconia argentea (Gentle 7883, CAS)
Miconia bubalina (Davidse 36635, CAS)
Miconia calvescens (Almeda 4190, CAS)
Miconia centrodesma (Davidse 36772, CAS)
Miconia chamissois (Bartlett 11657, CAS)
Miconia chrysophylla (Gentle 8172, CAS)
Miconia ciliata (Bartlett 11299, CAS)
Miconia desmantha (Holst 6006, CAS)
Miconia dodecandra (Davidse 36066, CAS)
Miconia elata (Schipp 150, F)
Miconia fulvostellata (Davidse 35896, CAS)
Miconia glaberrima (Hawkins 1519, CAS)
Miconia gracilis (Holst 5901, CAS)
Miconia holosericea (Holst 5794, CAS)
Miconia hondurensis (Holst 5136, CAS)
Miconia hyperprasina (Gentle 5109, CAS)
Miconia ibaguensis (Nee 46804, CAS)
Miconia impetiolaris (Holst 4414, CAS)
Miconia insularis (Gentle 7956, CAS)

Miconia lacera (Gentle 7469, CAS)
Miconia laevigata (Daniel 7043, CAS)
Miconia lateriflora (Gentle 8198, CAS)
Miconia longifolia (Gentle 4886, CAS)
Miconia matthaei (Gentle 9313, CAS)
Miconia minutiflora (Schipp S-554, F)
Miconia mirabilis (Holst 4061, CAS)
Miconia nervosa (Gentle 4291, CAS)
Miconia nutans (Davidse 36771, CAS)
Miconia oinochrophylla (Hawkins 1482, CAS)
Miconia oligocephala (Lundell 6608, NY)
Miconia paleacea (Schipp 1253, NY)
Miconia poeppigii (Cited by Almeda 2009)
Miconia prasina (Hawkins 1471, CAS)
Miconia punctata (Holst 5272, CAS)
Miconia reducens (Bartlett s.n., NY)
Miconia schippii (Gentle 9244, CAS)
Miconia schlechtendalii (Gentle 1354, NY)
Miconia schlimii (Davidse 36557, CAS)
Miconia smaragdina (Davidse 36725, CAS)
Miconia splendens (Gentle 8174, CAS)
Miconia stenostachya (Davidse 32448, CAS)
Miconia tomentosa (Balick 2608, US)
Miconia triplinervis (Davidse 36990, CAS)

Guatemala (62 spp)

Miconia aeruginosa (Lundell 20995, CAS)
Miconia affinis (Martínez 22941, CAS)
Miconia albicans (Contreras 9793, CAS)
Miconia alpestris (Breedlove 8541, CAS)
Miconia ampla (Contreras 8040, CAS)
Miconia argentea (Contreras 9618, CAS)
Miconia barbinervis (Cited by Standley & Williams 1963)
Miconia bupalina (Contreras 7936, CAS)
Miconia calvescens (Contreras 7970, CAS)
Miconia caudata (Kellerman 6677, US)
Miconia centrodesma (Esteban 22933, MO)
Miconia chamissois (Cited by Standley & Williams 1963)
Miconia chinantlana (Steyermark 48831, F)
Miconia chrysophylla (Contreras 11136, CAS)
Miconia ciliata (Contreras 520, CAS)
Miconia contrerasii (Lundell 20896, CAS)
Miconia desmantha (Contreras 11007, CAS)
Miconia dodecandra (Lundell 20993, US)
Miconia donaeana (Contreras 4531, CAS)
Miconia fulvostellata (Steyermark 49143a, F)
Miconia glaberrima (Contreras 10888, CAS)
Miconia globulifera (Breedlove 58106, US)
Miconia guatemalensis (Contreras 10913, CAS)
Miconia hemenostigma (Cited by Standley & Williams 1963)
Miconia holosericea (Lundell 19493, CAS)
Miconia hondurensis (Contreras 4259, DS)

Miconia hyperprasina (Contreras 9391, CAS)
Miconia ibaguensis (Contreras 2874, CAS)
Miconia impetiolaris (Bartlett 12634, CAS)
Miconia insularis (Contreras 9458, CAS)
Miconia lacera (Contreras 7926, CAS)
Miconia laevigata (Contreras 8230, CAS)
Miconia lateriflora (Contreras 7966, CAS)
Miconia livida (Contreras 10899, CAS)
Miconia longifolia (Contreras 2889, CAS)
Miconia lundellii (Lundell 18958, CAS)
Miconia matthaei (Contreras 4483, CAS)
Miconia mexicana (Lundell 20451, CAS)
Miconia minutiflora (Contreras 6601, CAS)
Miconia nervosa (Contreras 7587, CAS)
Miconia nutans (Contreras 4782, DS)
Miconia oinochrophylla (Contreras 7925, CAS)
Miconia oligocephala (Tuerckheim II 1781, NY)
Miconia poeppigii (Cited by Wurdack et al. 1993)
Miconia prasina (Lundell 19489, CAS)
Miconia punctata (von Türkheim 7866, NY)
Miconia purulensis (Croat 41183, CAS)
Miconia reducens (Schipp 1244, NY)
Miconia schippii (Contreras 7571, CAS)
Miconia schlechtendalii (Iltis G-136, CAS)
Miconia schlimii (Lundell 16259, CAS)
Miconia serrulata (Contreras 7936, CAS)
Miconia splendens (Contreras 11411, CAS)
Miconia stenostachya (Contreras 2228, CAS)
Miconia sylvatica (Rojas 341, US)
Miconia tacanensis (Breedlove 64794, CAS)
Miconia tixicensis (Lundell 19719, CAS)
Miconia tomentosa (Cited by Standley & Williams 1963, as *M. amplexans*)
Miconia trinervia (Lundell 19534, CAS)
Miconia triplinervis (Contreras 7899, CAS)
Miconia tuerckheimii (Lundell 18957, CAS)
Miconia zemurrayana (Heyde 6137, US)

Honduras (54 spp)

Miconia aeruginosa (Nelson 2334, CAS)
Miconia affinis (Saunders 354, CAS)
Miconia albicans (Caballero 149, CAS)
Miconia ampla (Liesner 26140, MO)
Miconia argentea (Nelson 5101, CAS)
Miconia barbinervis (MacDougal 3285, CAS)
Miconia bupalina (Nelson 9197, CAS)
Miconia calvescens (Schnell 982, US)
Miconia caudata (Mejía 149, CAS)
Miconia celaquensis (Thomas 79, CAS)
Miconia centrodesma (MacDougal 3403, CAS)
Miconia chamissois (Williams 10360, CAS)
Miconia chrysophylla (Nelson 5504, CAS)
Miconia ciliata (Nelson 4116, CAS)

Miconia desmantha (Muñoz 29, CAS)
Miconia dodecandra (Valladares 193, US)
Miconia donaeana (Liesner 26407, CAS)
Miconia elata (Nelson 8595, CAS)
Miconia glaberrima (Thomas 657, MO)
Miconia gracilis (Evans 1060, CAS)
Miconia guatemalensis (Liesner 26519, CAS)
Miconia hondurensis (Saunders 1213, CAS)
Miconia hyperprasina (Molina 1975)
Miconia ibaguensis (Nelson 5063, CAS)
Miconia impetiolaris (Evans 1598, CAS)
Miconia lacera (Martínez 129, CAS)
Miconia laevigata (Nelson 6742, CAS)
Miconia lateriflora (Evans 1637, CAS)
Miconia livida (Evans 1360, CAS)
Miconia lonchophylla (Croat 64249, CAS)
Miconia longifolia (Edwards 669, US)
Miconia matthaei (Evans 1678, CAS)
Miconia mexicana (Pilz 1376, CAS)
Miconia minutiflora (Saunders 896, CAS)
Miconia nutans (Cited by Molina 1975)
Miconia oinochrophylla (MacDougal 3280, CAS)
Miconia paleacea (Croat 64593, CAS)
Miconia poeppigii (Saunders 1224, CAS)
Miconia prasina (Saunders 634, CAS)
Miconia punctata (Saunders 1307, CAS)
Miconia purulensis (Meigs 1909, CAS)
Miconia reducens (Nelson 4992, CAS)
Miconia schlechtendalii (Davidse 35165, CAS)
Miconia schlimii (Hawkins 813, CAS)
Miconia serrulata (Saunders 1095, CAS)
Miconia splendens (Nelson & Vargas 5011, CAS)
Miconia stenostachya (Saunders 650, CAS)
Miconia stevensiana (Saunders 1223, CAS)
Miconia sylvatica (Rodriguez 152, CAS)
Miconia theizans (Daniel 9438, CAS)
Miconia tomentosa (Saunders 1158, CAS)
Miconia trinervia (Aguilar 4082, CAS)
Miconia triplinervis (Croat 42602, CAS)
Miconia zemurrayana (Davidse 35014, CAS)

El Salvador (19 spp)

Miconia aeruginosa (Villacorta 2068, CAS)
Miconia albicans (Sidwell 759, CAS)
Miconia argentea (Sandoval 1156, CAS)
Miconia caudata (Cited by Winkler 1965)
Miconia chamissois (Sidwell 749, CAS)
Miconia dodecandra (Tucker 804, NY)
Miconia glaberrima (Cited by Winkler 1965)
Miconia guatemalensis (Sandoval 1236, CAS)
Miconia ibaguensis (Sidwell 763, CAS)
Miconia laevigata (Chinchilla s.n., CAS)
Miconia mexicana (Villacorta 2149, CAS)

Miconia prasina (Sidwell 764, CAS)
Miconia reducens (Winkler 1963, as *M. subnodosa*)
Miconia schlechtendalii (Cited by Winkler 1965)
Miconia schlimii (Sandoval 1142, CAS)
Miconia sterilis (Tucker 1100, US)
Miconia sylvatica (Croat 42283, CAS)
Miconia tuckeri (Tucker 998, UC)
Miconia zemurrayana (Croat 42315, CAS)

Nicaragua (62 spp)

Miconia aeruginosa (Moreno 23497, CAS)
Miconia affinis (Sandino 3252, CAS)
Miconia albicans (Neill 7276, CAS)
Miconia ampla (De la Cerda 3489, CAS)
Miconia appendiculata (Rueda 3409, CAS)
Miconia argentea (Moreno 336, CAS)
Miconia astroplocama (Stevens 21002, CAS)
Miconia barbinervis (Stevens 16584, CAS)
Miconia bubalina (Neill 4372, CAS)
Miconia calvescens (Narvaez 3063, CAS)
Miconia caudata (Moreno 23496, CAS)
Miconia centrodesma (Martinez 2118, CAS)
Miconia chamissois (Stevens 6515, CAS)
Miconia chrysophylla (Stevens 8484, CAS)
Miconia ciliata (Marshall 6572, CAS)
Miconia decurrens (Sandino 130, CAS)
Miconia desmantha (Wilbur 16459, CAS)
Miconia dodecandra (Sandino 4695, CAS)
Miconia donaeana (Neill 299, CAS)
Miconia dorsiloba (Riviere 338, CAS)
Miconia elata (Stevens 7564, CAS)
Miconia fulvostellata (Sandino 225, CAS)
Miconia gracilis (Sandino 225, CAS)
Miconia gratissima (Holdridge 6624, CAS)
Miconia grayumii (Araquistain 3069, CAS)
Miconia guatemalensis (Moreno 1880, CAS)
Miconia holosericea (Proctor 26971, CAS)
Miconia hondurensis (Pipoly 4412, CAS)
Miconia hyperprasina (Stevens 6341, CAS)
Miconia ibaguensis (Pipoly 4010c, CAS)
Miconia impetiolaris (Stevens 7949, CAS)
Miconia lacera (Neill 2575, CAS)
Miconia laevigata (Sandino 2499, CAS)
Miconia lamprophylla (Neill 1902, MO)
Miconia lateriflora (Stevens 20691, CAS)
Miconia ligulata (Grijalva 440, CAS)
Miconia longifolia (Rueda 2478, CAS)
Miconia matthaei (Pipoly 3741, CAS)
Miconia mexicana (Atwood AN161, CAS)
Miconia minutiflora (Moreno 14630, CAS)
Miconia mirabilis (Little 25251, CAS)
Miconia multispicata (Stevens 18784, CAS)
Miconia nervosa (Sandino 161, CAS)

- Miconia nutans* (Stevens 11697, CAS)
Miconia oinochrophylla (Seymour 5649, CAS)
Miconia paleacea (Moreno 27231, CAS)
Miconia poeppigii (Little 25294, CAS)
Miconia prasina (Miller 1211, CAS)
Miconia punctata (Pipoly 3554, CAS)
Miconia reducens (Holdridge 6623, CAS)
Miconia schippii (Pipoly 4152, CAS)
Miconia schlimirii (Moreno 24275, CAS)
Miconia serrulata (Miller 1190, CAS)
Miconia shattuckii (Araquistain 3075, CAS)
Miconia simplex (Martínez 2121, CAS)
Miconia splendens (Moreno 23234, CAS)
Miconia stenostachya (Stevens 19608, CAS)
Miconia stevensiana (Stevens 8398, CAS)
Miconia tomentosa (Neill 3998, CAS)
Miconia tonduzii (Moreno 510, CAS)
Miconia trinervia (Sandino 4804, CAS)
Miconia triplinervis (Moreno 936, CAS)

Costa Rica (111 spp)

- Miconia aeruginosa* (Almeda 3652, CAS)
Miconia affinis (Aguilar 3285, NY)
Miconia albicans (Almeda 4062, CAS)
Miconia ampla (Almeda 3414, CAS)
Miconia amplinodis (Almeda 4999, CAS)
Miconia aponeura (Herrera 4661, CAS)
Miconia appendiculata (Araya 355, CAS)
Miconia arboricola (Grayum 8489, CAS)
Miconia argentea (Almeda 5540, CAS)
Miconia astroplocama (Almeda 5452, CAS)
Miconia barbinervis (Almeda 7027, CAS)
Miconia benthamiana (Almeda 3263, CAS)
Miconia biperulifera (Almeda 2209, CAS)
Miconia brenesii (Almeda 5050, CAS)
Miconia brevitheca (Almeda & Nakai 3866, CAS)
Miconia bubalina (Almeda 5126, CAS)
Miconia calocoma (Folsom 9990, CAS)
Miconia carnea (Almeda 4806, CAS)
Miconia caudata (Stork 4789, CAS)
Miconia centrodesma (Hammel 12403, CAS)
Miconia chamissois (Morales 4209, CAS)
Miconia chionophila (Davidse 25967, CAS)
Miconia chiriquiensis (Almeda 5322, CAS)
Miconia chrysophylla (Gomez-Laurito 10416, CAS)
Miconia ciliata (Almeda 3325, CAS)
Miconia cocoensis (Trusty 270, CAS)
Miconia colliculosa (Herrera 3129, CAS)
Miconia commutata (Folsom 9767, CAS)
Miconia confertiflora (Almeda 4516, CAS)
Miconia coriacea (Almeda 4667, CAS)
Miconia correae (Chacon 1347, USJ)
Miconia costaricensis (Almeda 3945, CAS)
Miconia cremadrena (Barringer 3824, CAS)

- Miconia cuspidatissima* (Almeda 5428, CAS)
Miconia danielii (Utley 2466, CAS)
Miconia desmantha (Grayum 8091, CAS)
Miconia diegogomezii (Lépiz 347, CAS)
Miconia dissita (Davidse 23199, CAS)
Miconia dissitiflora (Herrera 4335, CAS)
Miconia dissitinervia (Croat 59849, CAS)
Miconia dodecandra (Almeda 2772, CAS)
Miconia dolichopoda (Almeda 3815, CAS)
Miconia dolichorrhyncha (Almeda 3059, CAS)
Miconia donaeana (Almeda 6720, CAS)
Miconia dorsiloba (McDowell 143, CAS)
Miconia elata (Almeda 4139, CAS)
Miconia friedmaniorum (Almeda 5416, CAS)
Miconia globuliflora (Haber 2356, CAS)
Miconia gracilis (Almeda 6951, CAS)
Miconia grandidentata (Almeda 5251, CAS)
Miconia grayumii (Smith 152, CAS)
Miconia hildeana (Kriebel 3667, CAS)
Miconia holosericea (Skutch 5061, CAS)
Miconia hondurensis (Almeda 4088, CAS)
Miconia hyperprasina (Burger 11749, CAS)
Miconia ibaguensis (Almeda 4268, CAS)
Miconia impetiolaris (Hammel 8453, CAS)
Miconia incurva (Bello 1544, CAS)
Miconia insularis (Grayum 5844, CAS)
Miconia intricata (Gomez 21092, CAS)
Miconia kappellei (Herrera 5911, CAS)
Miconia lacera (Almeda 3314, CAS)
Miconia laevigata (Schubert 658, US)
Miconia lamprophylla (Burger 10625, CAS)
Miconia lateriflora (Almeda 7021, CAS)
Miconia ligulata (Stevens 24356, CAS)
Miconia livida (Grayum 7756, CAS)
Miconia lonchophylla (Almeda 7147, CAS)
Miconia longibracteata (Mora 530, CAS)
Miconia longifolia (Davidse 31228, CAS)
Miconia loreyoides (Herrera 2829, CAS)
Miconia matthaei (Almeda 6570, CAS)
Miconia melanotricha (Almeda 2912, CAS)
Miconia mexicana (Almeda 2836, CAS)
Miconia minutiflora (Almeda 3329, CAS)
Miconia mirabilis (Herrera 4390, CAS)
Miconia multiplinervia (Almeda 5225, CAS)
Miconia multispicata (Almeda 3181, CAS)
Miconia nervosa (Almeda 5561, CAS)
Miconia nutans (Almeda 3910, CAS)
Miconia oinochrophylla (Almeda 5551, CAS)
Miconia oraria (Vargas 116, CAS)
Miconia osaensis (Aguilar 6208, CAS)
Miconia paleacea (Almeda 7015, CAS)
Miconia pendula (Almeda 7245, CAS)
Miconia pittieri (Almeda 3830, CAS)

Miconia poeppigii (Hammel 18427, CAS)
Miconia prasina (Almeda 4087, CAS)
Miconia pterocaulon (Burger 4614, CAS)
Miconia punctata (Almeda 6567, CAS)
Miconia reducens (Stevens 25081, TEFH)
Miconia ricardoi (Kriebel 3384, CAS)
Miconia rubiginosa (Almeda 3328, CAS)
Miconia schlimii (Almeda 3391, CAS)
Miconia schnellii (Almeda 6786, CAS)
Miconia serrulata (Almeda 5126, CAS)
Miconia shattuckii (Jiménez 877, CAS)
Miconia simplex (Almeda 7022, CAS)
Miconia smaragdina (Almeda 5465, CAS)
Miconia sparrei (Zamora 2451, CAS)
Miconia splendens (Villalobos 170, CAS)
Miconia stenostachya (Almeda 4069, CAS)
Miconia stevensiana (Folsom 9633, CAS)
Miconia talamancensis (Almeda 4777, CAS)
Miconia theizans (Almeda 4182, CAS)
Miconia tomentosa (Zamora 1248, CAS)
Miconia tonduzii (Almeda 7366, CAS)
Miconia trinervia (Almeda 6721, CAS)
Miconia triplinervis (Herrera 2192, CAS)
Miconia valerioana (Vargas 111, CAS)
Miconia vestita (Almeda 7399, CAS)

Panama (107 spp)

Miconia aeruginosa (Woodson 1143, MO)
Miconia affinis (Almeda 5974, CAS)
Miconia albicans (Galdames 2019, CAS)
Miconia ampla (Foster 14690, CAS)
Miconia appendiculata (Antonio 3819, CAS)
Miconia arboricola (Almeda 7524, CAS)
Miconia argentea (Almeda 5954, CAS)
Miconia astroplocama (Almeda 7643, CAS)
Miconia barbinervis (Correa 16130, CAS)
Miconia benthamiana (Hernández 746, CAS)
Miconia biperulifera (Gomez 21929, CAS)
Miconia brachybotrya (De Nevers 8301, CAS)
Miconia brevitheca (Folsom 4320, CAS)
Miconia bubalina (Almeda 6025, CAS)
Miconia calvescens (Hammel 14812, CAS)
Miconia carnea (Almeda 6143, CAS)
Miconia caudata (Allen 3734, MO)
Miconia centrodesma (Aranda 1570, CAS)
Miconia centronioides (McPherson 12517, CAS)
Miconia centrosperma (McPherson 10004, CAS)
Miconia chamissois (Von Wedel 528, CAS)
Miconia chionophila (Gomez 22662, CAS)
Miconia chiriquiensis (Aranda 1397, CAS)
Miconia chrysophylla (Almeda 5961, CAS)
Miconia ciliata (Croat 67050, CAS)

Miconia cionotricha (Sullivan 707, CAS)
Miconia colliculosa (Almeda 6514, CAS)
Miconia coloradensis (Folsom 4733, CAS)
Miconia commutata (Correa 4014, CAS)
Miconia concinna (Herrera 3520, CAS)
Miconia correae (Dressler 5806, CAS)
Miconia costaricensis (Almeda 7527, CAS)
Miconia crocata (Aranda 2849, CAS)
Miconia cuspidatissima (Van der Werff 6349, CAS)
Miconia desmantha (Cited by Standley & Williams 1963)
Miconia dissita (Almeda 6226, CAS)
Miconia dissitinervia (Liesner 201, CAS)
Miconia dodecandra (Galdames 2459, CAS)
Miconia dolichorrhyncha (Folsom 3974, CAS)
Miconia donaeana (McPherson 11750, CAS)
Miconia dorsiloba (Hammel 4809, CAS)
Miconia elata (Almeda 6013, CAS)
Miconia emendata (Almeda 5996, CAS)
Miconia friedmaniorum (Almeda 7618, CAS)
Miconia gentryi (McPherson 14062, CAS)
Miconia globuliflora (Croat 66416, CAS)
Miconia goniostigma (Almeda 3468, CAS)
Miconia gracilis (Almeda 7588, CAS)
Miconia grandidentata (McPherson 7718, CAS)
Miconia gratissima (Dwyer 4365, US)
Miconia hildeana (McPherson 11090, CAS)
Miconia holosericea (Correa 10104, CAS)
Miconia hondurensis (Almeda 6020, CAS)
Miconia ibaguensis (Hammel 1774, CAS)
Miconia impetiolaris (De Nevers 4947, CAS)
Miconia insularis (Hamilton 1429, CAS)
Miconia intricata (McPherson 12780, CAS)
Miconia iteophylla (Almeda 7668, CAS)
Miconia jefensis (Almeda 7696, CAS)
Miconia lacera (Almeda 5819, CAS)
Miconia lateriflora (Almeda 6333, CAS)
Miconia ligulata (Aranda 2246, CAS)
Miconia livida (McPherson 10385, CAS)
Miconia lonchophylla (Almeda 6101, CAS)
Miconia longifolia (Almeda 6026, CAS)
Miconia loreyoides (Almeda 6366, CAS)
Miconia matthaei (McPherson 8559, CAS)
Miconia melanotricha (Almeda 3487, CAS)
Miconia minutiflora (Almeda 7682, CAS)
Miconia morii (Mori 3773, CAS)
Miconia multiplinervia (Almeda 7536, CAS)
Miconia multispicata (Almeda 5882, CAS)
Miconia nervosa (Wilbur 11329, CAS)
Miconia nutans (Almeda 6317, CAS)
Miconia oinochrophylla (Aranda 2252, CAS)
Miconia oraria (de Nevers 5463, CAS)
Miconia peltata (Aranda 185, CAS)

Miconia phaeophylla (Sytsma 1995, CAS)
Miconia pittieri (Schmalzel 1673, CAS)
Miconia poeppigii (Nee 11085, CAS)
Miconia prasina (Almeda 6494, CAS)
Miconia procumbens (Hammel 16388, CAS)
Miconia pterocaulon (Almeda 5965, CAS)
Miconia punctata (Almeda 5828, CAS)
Miconia racemosa (Correa 4790, CAS)
Miconia reducens (Almeda 5909, CAS)
Miconia robinsoniana (Cited by Gleason 1958)
Miconia rubiginosa (Schmalzel 1546, CAS)
Miconia rufostellulata (D'Arcy 5216, CAS)
Miconia santaritensis (Liesner 1271, CAS)
Miconia schlimii (Galdames 2765, CAS)
Miconia serrulata (de Nevers 3569, CAS)
Miconia shattuckii (Herrera 1095, CAS)
Miconia simplex (Croat 60137, CAS)
Miconia smaragdina (de Nevers 8496, CAS)
Miconia splendens (Almeda 6027, CAS)
Miconia stenostachya (Carcamo 268, CAS)
Miconia stevensiana (McPherson 21014, CAS)
Miconia talamancensis (McPherson 9487, CAS)
Miconia theizans (Folsom 4320, CAS)
Miconia tomentosa (Almeda 5963, CAS)
Miconia tonduzii (Almeda 6318, CAS)
Miconia trinervia (McPherson 10855, CAS)
Miconia triplinervis (Hamilton 1125, CAS)
Miconia umbriensis (Almeda 5850, CAS)
Miconia urticoides (McPherson 14071, CAS)
Miconia valerioana (McPherson 14080, CAS)

Cuba (46 spp)

Miconia acunae (Stahlick 27233 photo in NY)
Miconia albicans (Acevedo-Rodriguez 6425, US)
Miconia alternifolia (Ekman 5389, US)
Miconia ancistrophora (Britton 5866, NY)
Miconia androsaemifolia (Morton 10059, US)
Miconia baracoensis (Howard 5875, US)
Miconia bisulcata (Ekman 14707, US)
Miconia bucherae (Cited in description)
Miconia calycina (León 10540, NY)
Miconia cerasiflora (Morton 9054, US)
Miconia cubensis (Jack 7425, US)
Miconia delicatula (Killip 44726, US)
Miconia dodecandra (Shafer 8123, US)
Miconia elata (Bro. Hioram 6061, US)
Miconia ibaguensis (Killip 13558, US)
Miconia impetiolaris (Liogier 3459, US)
Miconia javorkaeana (Cited in description)
Miconia laevigata (Fre. Leon 8532, US)
Miconia lanata (Cited by Liogier 1963)
Miconia lenticellata (Liogier 3746, NY)
Miconia matthaei (Britton 5251, NY)

Miconia minutiflora (Morton 4367, US)
Miconia mirabilis (Webster 3814, US)
Miconia moensis (Shafer 8038, US)
Miconia nystroemii (Ekman 7084, US)
Miconia obtusa (Marie-Victorin 21626, US)
Miconia perelegans (Britton 14443, US)
Miconia petersonii (López 119, US)
Miconia prasina (Britton 7192, US)
Miconia pteroclada (Ekman 14410, NY)
Miconia punctata (Bro. Leon 72336, US)
Miconia remotiflora (Ekman H-14814, US)
Miconia rhombifolia (Shafer 4136, NY)
Miconia rufa (Wright 2524, NY)
Miconia serrulata (Bro. Leon 5099, US)
Miconia shaferi (Liogier 3712, US)
Miconia skeaniana (Ekman 8855, NY)
Miconia splendens (Ekman 13850, US)
Miconia tetrandra (Ekman 9910, US)
Miconia tetrastoma (Bro. Clemente 7345, US)
Miconia tomentosa (Liogier 4301, US)
Miconia turquicensis (Léon 10939, US)
Miconia uninervis (Liogier 1008, US)
Miconia urceolata (Ekman 15644, US)
Miconia victorinii (Léon 20670, US)
Miconia walterjuddii (Bécquer 85487, NY)

Jamaica (26 spp)

Miconia albicans (Wurdack 2622, US)
Miconia ampla (Proctor 11345, US)
Miconia chrysophylla (Prior s.n., NY)
Miconia ciliata (Proctor 11303, US)
Miconia dodecandra (Proctor 22614, US)
Miconia elata (Wurdack 2628, US)
Miconia furfuracea (Cited by Proctor 1972)
Miconia impetiolaris (Harris 8558, US)
Miconia laevigata (Howard 14577, US)
Miconia matthaei (Cited by Proctor 1972)
Miconia mirabilis (Andrews 773, US)
Miconia multispicata (Proctor 30014, US)
Miconia nubicola (Proctor 9691, US)
Miconia prasina (Wurdack 2617, US)
Miconia pseudorigida (Proctor 10469, US)
Miconia pyramidalis (Judd 8322, FLAS)
Miconia quadrangularis (Proctor 6592, US)
Miconia racemosa (Harris 11154, NY)
Miconia rigida (Skean Jr. 1939, NY)
Miconia rubens (Harris 6940, US)
Miconia serrulata (Yuncker 17921, US)
Miconia splendens (Proctor 10082, US)
Miconia tetrandra (Proctor 10092, US)
Miconia theizans (Kapos 1560, US)
Miconia trinervia (Proctor 18430, US)
Miconia triplinervis (Proctor 26537, US)

Haiti (39 spp)

- Miconia abeggii* (Cited in description)
Miconia adenocalyx (Ekman 2836, US)
Miconia albiviridis (Ekman 1400, US)
Miconia alloetricha (Judd 6956a, NY)
Miconia apiculata (Ekman H-10641, US)
Miconia barkeri (Ekman 5329, US)
Miconia basileensis (Ekman 9663, US)
Miconia calycina (Cited by Liogier 2000)
Miconia coniophora (Ekman H-8067, US)
Miconia dodecandra (Ekman H5914)
Miconia domingensis (Ekman H3147)
Miconia favosa (Judd 4455, NY)
Miconia ferruginea (Skean Jr. 3383, NY)
Miconia fuertesii (Ekman H2498, NY)
Miconia hypoides (Ekman 7552, US)
Miconia impetiolaris (Bro. Augusto 588, US)
Miconia krugii (Ekman H9662, NY)
Miconia laevigata (Zanoni 33337, NY)
Miconia lanceolata (Howard 12587, US)
Miconia leptantha (Cited in description)
Miconia luteola (Ekman H-1407, US)
Miconia macayana (Skean Jr. 1428, NY)
Miconia markgraffii (Cited in description)
Miconia mirabilis (Ekman H4700, NY)
Miconia multiglandulosa (Ekman H4347, NY)
Miconia nematophora (Ekman 6218, US)
Miconia ossaeifolia (Ekman 10350, US)
Miconia prasina (Cited by Liogier 2000)
Miconia pyramidalis (Leonard 9511, NY)
Miconia racemosa (Zanoni 35811, NY)
Miconia rigidissima (Ekman 1425, US)
Miconia selleana (Judd 4528, NY)
Miconia serrulata (Nash 220, NY)
Miconia stenobotrys (Cited by Judd & Karpoor 1993)
Miconia subcompressa (Zanoni 25668, NY)
Miconia tetrandra (Ekman H10585, NY)
Miconia tetrastoma (Skean Jr. 1677, NY)
Miconia tetrazygioides (Ekman 10684, NY)
Miconia xenotricha (Ekman H7447, NY)

Dominican Republic (41 spp)

- Miconia adenocalyx* (Judd 6688, NY)
Miconia affinis (Zanoni 40863, US)
Miconia alainii (Cited in description)
Miconia bifaria (Cited in description)
Miconia calycina (Zanoni 39773, NY)
Miconia campanensis (Cited in description)
Miconia coniophora (Skean Jr. 1821, NY)
Miconia desportesii (Fuertes Lorén 1491, NY)
Miconia dielsiana (Ekman H13229, NY)
Miconia dodecandra (Judd 8093, NY)
Miconia domingensis (Liogier 16818, NY)

Miconia ferruginea (Judd 8117, NY)

- Miconia fuertesii* (Fuertes 1905, NY)
Miconia howardiana (Judd 5178, NY)
Miconia impetiolaris (Cited by Liogier 2000)
Miconia jimenezii (Judd 5192, NY)
Miconia krugii (Judd 6645, NY)
Miconia laevigata (Cited by Liogier 2000)
Miconia lanceolata (Zanoni 33820, NY)
Miconia luteola (Zanoni 33793, NY)
Miconia mansfeldiana (Ekman H11737, NY)
Miconia mirabilis (Judd 6507, NY)
Miconia monciona (Ekman 12807, US)
Miconia nematophora (Cited in description)
Miconia prasina (Zanoni 41404, NY)
Miconia punctata (Zanoni 43939, US)
Miconia pyramidalis (Judd 6576, NY)
Miconia pyrifolia (Cited by Wurdack 1973 and 1993, as *M. ottoschultzii*)
Miconia racemosa (García 1959, US)
Miconia rigida (Cited by Liogier 2000)
Miconia rubiginosa (Jimenez 1098, US)
Miconia samanensis (Abbott 2233, NY)
Miconia santanana (Ekman H11759, NY)
Miconia septentrionalis (Zanoni 42455, NY)
Miconia serrulata (Judd 5138, NY)
Miconia sphagnicola (Ekman H12808, NY)
Miconia subcompressa (Zanoni 18892, NY)
Miconia tetrandra (Zanoni 41155, NY)
Miconia tetrastoma (Judd 5179, NY)
Miconia viscidula (Fuertes Lorén 1753, NY)
Miconia zanonii (Judd 5158, US)

Puerto Rico (20 spp)

- Miconia affinis* (Sintenis 2738, US)
Miconia calycina (Sintenis 5972, NY)
Miconia foveolata (King 10603, US)
Miconia impetiolaris (Sintenis 4518, US)
Miconia laevigata (Britton 2755, US)
Miconia lanata (Cited by Liogier & Martorell 1982)
Miconia mirabilis (Mejia 90, US)
Miconia pachyphylla (Howard 15743, US)
Miconia prasina (Liogier 37060, US)
Miconia punctata (Sintenis 6426, US)
Miconia pycnoneura (Howard 16826, US)
Miconia pyramidalis (Judd 6087, NY)
Miconia racemosa (Boom 10065, US)
Miconia rubiginosa (Liogier 36798, US)
Miconia serrulata (Liogier 32761, US)
Miconia sintenisii (Boom 10255, US)
Miconia splendens (Cited by Liogier & Martorell 1982)
Miconia tetrandra (Proctor 44546, US)
Miconia tetrastoma (Woodbury 17890, NY)
Miconia thomasiana (Axelrod 5208, US)

Virgin Islands (2 spp)

Miconia laevigata (Little 26075, US)
Miconia thomasiana (Fishlock 93, US)

Saba (Dutch W. Indies; 1 sp)

Miconia laevigata (Howard 18132, NY)

Saint Eustatius (Dutch W. Indies; 2 spp)

Miconia impetiolaris (Boldinch 469, NY)
Miconia laevigata (Cited by Howard 1989)

Saint Kitts & Nevis (3 spp)

Miconia laevigata (Proctor 19542, US)
Miconia mirabilis (Britton 384 (NY))
Miconia striata (Britton 708 (NY))

Antigua (4 spp)

Miconia laevigata (Box 1526, US)
Miconia mirabilis (Cited by Howard 1989)
Miconia striata (Smith 10383, NY)
Miconia trichotoma (Duss 47, NY)

Montserrat (3 spp)

Miconia laevigata (Howard 19668, US)
Miconia mirabilis (Shafer 524, NY)
Miconia striata (Howard 18942, NY)

Guadeloupe (12 spp)

Miconia coriacea (Holdridge 431)
Miconia cornifolia (Barrier 3735, NY)
Miconia furfuracea (Barrier 2359, NY)
Miconia globuliflora (Howard 19794, US)
Miconia impetiolaris (Cited by Howard 1989)
Miconia laevigata (Fairchild 3780, US)
Miconia martinicensis (Cited by Howard 1989)
Miconia mirabilis (Raynal-Roques AR.19598, NY)
Miconia prasina (Duchassaing s.n., US)
Miconia striata (Duss 2253 (NY))
Miconia tetrandra (Duss 3469, US)
Miconia trichotoma (Duss 2254, US)

Dominica (15 spp)

Miconia coriacea (Wasshausen 410, US)
Miconia cornifolia (Cowan 1598, NY)
Miconia dodecandra (Whitefoord 7230, US)
Miconia ernstii (Ernst 1767, US)
Miconia furfuracea (Whitefoord 4248, US)
Miconia globuliflora (Hodge 1398, NY)
Miconia impetiolaris (Hill 25452, US)
Miconia laevigata (Webster 13278, US)
Miconia mirabilis (Stijfhoorn 739, US)
Miconia mornicola (Lack 2, US)
Miconia prasina (Hodge 508 (NY))

Miconia racemosa (Howard 1168, US)

Miconia striata (Hill 25689, US)
Miconia tetrandra (Penneys 1300, NY)
Miconia trichotoma (Imray 219, NY)

Martinique (9 spp)

Miconia acinodendron (Cited by Howard 1989)
Miconia cornifolia (Barrier 4522, NY)
Miconia furfuracea (Duss 1175, NY)
Miconia globuliflora (Egler 39-101, NY)
Miconia laevigata (Stehlé 6906, US)
Miconia martinicensis (Duss 4047, US)
Miconia mirabilis (Hahn 453, NY)
Miconia prasina (Duss 4412, NY)
Miconia striata (Duss 116 (NY))

Santa Lucia (9 spp)

Miconia cornifolia (Smith 988, NY)
Miconia furfuracea (Howard 19909, US)
Miconia laevigata (Howard 19982, US)
Miconia luciana (Webster 9404, US)
Miconia mirabilis (Howard 19842, NY)
Miconia prasina (Proctor 17726, NY)
Miconia racemosa (Howard 19873, NY)
Miconia secunda (Proctor 17726, A)
Miconia striata (Howard 11445, NY)

Saint Vincent (2 spp)

Miconia andersonii (Beard s.n., US)
Miconia laevigata (Morton 4864, US)

Barbados (2 spp)

Miconia cornifolia (Cited by Howard 1989)
Miconia laevigata (Waby 92, US)

Grenada (7 spp)

Miconia andersonii (Morton 5874, US)
Miconia cornifolia (Broadway 1771, NY)
Miconia laevigata (Proctor 16817, US)
Miconia mirabilis (Broadway s.n., NY)
Miconia prasina (Broadway s.n., NY)
Miconia racemosa (Broadway s.n., US)
Miconia tetrandra (Broadway 7894, US)

Colombia (338 spp)

Miconia abbreviata (Sneidern K-1188, US)
Miconia acalephoides (Cuatrecasas 22278, US)
Miconia acanthocoryne (Sneidern 5364, US)
Miconia acinodendron (Callejas 4633, US)
Miconia acuminifera (Murcia 73, US)
Miconia acutipetala (Cited by Cogniaux 1891)
Miconia aenigmatica (Cuatrecasas 21752, US)
Miconia aeruginosa (Roldán 1402, US)

- Miconia affinis* (Callejas 5799, US)
Miconia aggregata (Gentry 47716, US)
Miconia aguirrei (Aguirre 428, NY)
Miconia alata (Gonzalez 3034, COL)
Miconia alberti (Killip 15896, US)
Miconia albicans (Fernández 2752, US)
Miconia alborosea (Uribe-Uribe 6082, US)
Miconia alborufescens (Echeverry 4758, COL)
Miconia alternans (Fonnegra 4500, US)
Miconia alypifolia (Mutis 5163, US)
Miconia amazonica (Rudas 1651, US)
Miconia ambyandra (Cuatrecasas 12259, US)
Miconia amnicola (Little 9682, US)
Miconia ampla (Idrobo 705, US)
Miconia andreana (Arbelaez 2248, US)
Miconia anisophylla (Triana s.n., US)
Miconia annulata (Uribe-Uribe 5484, US)
Miconia antioquiensis (Uribe-Uribe 2697, US)
Miconia apostachya (Garcia-Barriga 14101, US)
Miconia aponeura (Quiñones 1330, US)
Miconia appendiculata (Sneidern A-1322, US)
Miconia araguensis (Garcia-Barriga 20546, US)
Miconia arboricola (Betancur 4780, COL)
Miconia archeri (Archer 768, US)
Miconia argentea (Diaz 231, COL)
Miconia argyrophylla (Cited by Cogniaux 1891)
Miconia asclepiadea (Cuatrecasas 26880, US)
Miconia asperrima (Uribe-Uribe 6236, US)
Miconia astroplocama (Almeda 10668, COL)
Miconia astrotricha (Schultes 15591, US)
Miconia atropilis (Lehmann 6216, photo in US)
Miconia aurea (Idrobo 890, US)
Miconia aureoides (Schultes 18988, US)
Miconia barbinervis (Sneidern 5069, US)
Miconia bella (Garcia-Barriga 16122, US)
Miconia benthamiana (Forero 2346, US)
Miconia biappendiculata (Cleef 7387, US)
Miconia biglandulosa (Vester 97, US)
Miconia blakeifolia (Cuatrecasas 15181, US)
Miconia bordoncilloana (Muñoz 60, COL)
Miconia borjensis (Córdoba 7526, FMB)
Miconia brachycalyx (Gentry 17281, US)
Miconia brachygyna (Killip 15266, US)
Miconia bracteolata (Cuatrecasas 2654, US)
Miconia brevitheca (Cuatrecasas 11412, COL)
Miconia bupalina (Quiñones 1574, US)
Miconia buxifolia (Killip 18186, US)
Miconia caesia (Barclay 3352, US)
Miconia calvescens (Fernández Alonso 8799, US)
Miconia cannabina (Pipoly 15144, US)
Miconia capitellata (Lehmann 8519, photo in US)
Miconia carassana (Schultes 16132, US)
Miconia cataractae (Pennell 2121, US)

Miconia caucana (Holton 909, US)
Miconia caudata (Little 7618, US)
Miconia cautis (Schultes 10129, US)
Miconia cazaletii (Ewan 16743, US)
Miconia centrodesma (Forero 4057, US)
Miconia centrodesmoides (Robinson 136, US)
Miconia centronioides (Murphy 536, US)
Miconia chamissois (Allen 596, MO)
Miconia chionophila (Cuatrecasas 25622, US)
Miconia chlorocarpa (Cuatrecasas 9234, US)
Miconia chrysocoma (Rangel 2100, US)
Miconia chrysophylla (Uribe-Uribe 3445, US)
Miconia ciliata (Sastre 2204, US)
Miconia cionotricha (Fuchs 22348, US)
Miconia cladonia (Uribe-Uribe 6117, US)
Miconia clathrantha (Mendoza 14124, FMB)
Miconia cleefii (Cleef 1204, US)
Miconia clypeata (Lutelyn 7325, US)
Miconia codonostigma (Cuatrecasas 22468, US)
Miconia compressicaulis (Core 1284, US)
Miconia cordifolia (D'Arcy 15631, US)
Miconia coronata (Archer 578, US)
Miconia coronifera (Cuatrecasas 11636, US)
Miconia corymbiformis (Espinal 3458, US)
Miconia costaricensis (Uribe-Uribe 1927, US)
Miconia crinita (Stein 3441, US)
Miconia crocea (López-Palacios 3735, US)
Miconia cruenta (Triana s.n., US)
Miconia cuatrecasae (Cuatrecasas 2509, US)
Miconia cundinamarcensis (Orozco 2665, US)
Miconia dapsiliflora (Mendoza 8641, FMB)
Miconia decipiens (Cited by Cogniaux 1891)
Miconia decurrens (Schultes 3047, US)
Miconia denticulata (Barclay 3371, US)
Miconia desmantha (Espinal 2874, US)
Miconia diaphanea (Pipoly 15108, US)
Miconia dichrophylla (Cuatrecasas 8882, COL)
Miconia difficilis (Spruce 4972, photo in US)
Miconia divergens (Triana 3999, photo in US)
Miconia dodecandra (Escobar 8346, US)
Miconia dolichopoda (Kirkbride 1998, US)
Miconia dolichorrhyncha (Denslow 2238, US)
Miconia donaeana (Hoyos 91, HUA)
Miconia dorsiloba (Forero 3321, US)
Miconia dunstervillei (Almeda 10541, COL)
Miconia elaeoides (Melampy 1044, US)
Miconia elata (Zarucchi 487, US)
Miconia elviraе (Cuatrecasas 12722, COL)
Miconia emendata (Vasquez 12534, US)
Miconia eremita (Uribe-Uribe 4444, NY)
Miconia erioclada (Uribe-Uribe 4693, US)
Miconia erosa (Zarucchi 5969, US)
Miconia eugenoides (Schultes 13188, US)

- Miconia filamentosa* (Pipoly 15155, US)
Miconia fissa (Garcia-Barriga 13780, US)
Miconia floribunda (Lawrance 270, US)
Miconia fosteri (Vester 843, US)
Miconia frontinoana (Escobar 2295, US)
Miconia gentryi (Gentry 28624, US)
Miconia gibba (Mendoza 14222, FMB)
Miconia glaberrima (Dryander 2073, US)
Miconia gleasoniana (Pennell 9366, US)
Miconia goniostigma (Forero 2453, US)
Miconia gossypina (Triana s.n., US)
Miconia goudotii (Cited by Cogniaux 1891)
Miconia gracilis (Zarucchi 5074, US)
Miconia grandiflora (Lozano 5938, US)
Miconia gratissima (Franco 6059, COL)
Miconia hadrophylla (Garcia-Barriga 12946, US)
Miconia harlingii (Espinol 996, US)
Miconia haughtii (Haught 5201, US)
Miconia hematostemon (Triana 6258, US)
Miconia heterotricha (Sandeman 6042, photo in US)
Miconia holosericea (Schultes 16399, US)
Miconia hookeriana (Klug 1899, US)
Miconia hymenanthera (Restrepo 557, US)
Miconia ibaguensis (Uribe-Uribe 2649, US)
Miconia idroboi (Idrobo 837, US)
Miconia imbricata (Pennell 10503, US)
Miconia impetiolaris (Zarucchi 6513, US)
Miconia inaequalifolia (Schultes 16685, US)
Miconia ingens (Hilty AX-22, US)
Miconia insueta (Barclay 6790, US)
Miconia intricata (Léon 1502, US)
Miconia jahnii (Uribe-Uribe 5936, US)
Miconia jentaculorum (Cuatrecasas 13530, US)
Miconia juruensis (Little 9649, US)
Miconia kraenzlinii (Lehmann 3025, US)
Miconia lacera (Quiñones 1576, US)
Miconia laetivirens (Uribe-Uribe 5564, US)
Miconia laevigata (Kirkbride 2546, US)
Miconia lamprarrhena (Triana 404, US)
Miconia lamprophylla (Cuatrecasas 11233, US)
Miconia lanata (Uribe-Uribe 3110, US)
Miconia lateriflora (Shepherd 513, US)
Miconia latifolia (Cleef 7132, US)
Miconia lasiocalyx (Almeda 10297, COL)
Miconia lehmannii (Escobar 3582, US)
Miconia lepidota (Cited by Cogniaux 1891)
Miconia licrophora (Fosberg 19266, US)
Miconia liesneri (Cuatrecasas 13092, US)
Miconia ligulata (Juncosa 1837, US)
Miconia ligustrina (Uribe-Uribe 5099, US)
Miconia limitaris (Cuatrecasas 25228, US)
Miconia lithophila (King 6178, US)
Miconia lonchophylla (Croat 56455, US)

Miconia longifolia (Forero 4008, US)
Miconia longispicata (Zarucchi 1608, US)
Miconia loreyoides (Forero 5364, US)
Miconia lourteigiana (Plowman 2262, US)
Miconia lugonis (Ramos 2538, US)
Miconia luteynii (Luteyn 7512, US)
Miconia macrantha (Rivera 726, US)
Miconia macrothyrsa (Cuatrecasas 4160, US)
Miconia macrotis (Callejas 4091, US)
Miconia magdalena (Callejas 5266, US)
Miconia majalis (André 2768, photo in US)
Miconia manicata (Lehmann 6688, US)
Miconia marginata (Schultes 17082, US)
Miconia maroana (Barbosa 8384, COL)
Miconia matthaei (Callejas 9282, US)
Miconia mazanana (Pipoly 15423, US)
Miconia megalantha (Denslow 2325, US)
Miconia mesmeana (Croat 56463, US)
Miconia micayana (Killip 1184, US)
Miconia mimica (Killip 7936, US)
Miconia minuta (Cuatrecasas 8704, US)
Miconia minutiflora (Betancurt 1348, US)
Miconia miocarpa (Uribe-Uribe 3007, US)
Miconia mirabilis (Tillet 737-228, US)
Miconia mituana (Sastre 2210, US)
Miconia mocquerysii (Uribe-Uribe 3958, US)
Miconia mollicula (Uribe-Uribe 6433, US)
Miconia mulleola (Cuatrecasas 12935, US)
Miconia multiplinervia (Cuatrecasas 8410, US)
Miconia multispicata (Uribe-Uribe 6341, US)
Miconia mutisiana (Cuatrecasas 2506, MA)
Miconia myriantha (Zarucchi 1709, US)
Miconia myrtillifolia (Cuatrecasas 28669, US)
Miconia napoana (Schwabe 73/171, US)
Miconia nervosa (González 2443, US)
Miconia neurotricha (Triana s.n., photo in US)
Miconia nigripes (Lehmann 5474, US)
Miconia nodosa (Schultes 3240, US)
Miconia notabilis (Devia 997, US)
Miconia nutans (Callejas 2703, US)
Miconia obovata (Bro. Tomas 23701, US)
Miconia ochracea (Camp E-342, US)
Miconia oinochrophylla (Callejas 2343, US)
Miconia oraria (Forero 1507, US)
Miconia orcheotoma (Espinol 3341, US)
Miconia oreogena (Barclay 6684, US)
Miconia orescia (Uribe-Uribe 5560, US)
Miconia pachydonta (Philipson 1482, COL)
Miconia paleacea (King 6093, US)
Miconia pandurata (Cited in Cogniaux 1891)
Miconia parvifolia (Fosberg 19063, US)
Miconia paspaloides (Woronow 6436, cited in description)
Miconia pastoensis (Jameson 415, photo in US)

- Miconia pedicellata* (Cuatrecasas 13570, US)
Miconia penicillata (Pennell 5808, US)
Miconia pergamentacea (Fosberg 21104, US)
Miconia perijensis (Cuatrecasas 25175, US)
Miconia phaeochaeta (Schwabe 73/163, US)
Miconia phaeophylla (Little 7307, US)
Miconia phanerostila (Moreno 887, NY)
Miconia pichinchensis (Cited by Wurdack 1980)
Miconia piperifolia (Cited in description)
Miconia platypoda (Cuatrecasas 7185, NY)
Miconia plena (Lawrance 152, US)
Miconia plethorica (Uribe-Uribe 4777, US)
Miconia poecilantha (Barclay 3602, US)
Miconia poeppigii (Schultes 24931, US)
Miconia polyneura (Cuatrecasas 27687, US)
Miconia popayanensis (Lehmann 5473, US)
Miconia porphyrotricha (Gentry 55205, MO)
Miconia pozensis (Sandeman 6114, US)
Miconia prasina (Sastre 3032, US)
Miconia prasinifolia (Killip 9011, US)
Miconia pseudoradula (Sneidern 1447, US)
Miconia psychrophila (Díaz 2164, US)
Miconia pterocaulon (Denslow 2346, US)
Miconia puberula (Gasche 102, US)
Miconia pubipetala (Cuatrecasas 6991, US)
Miconia pulvinata (Smith 1841, US)
Miconia punctata (Cuatrecasas 15741, US)
Miconia punctibullata (Morales-P. 1884, COL)
Miconia puracensis (Cuatrecasas 23449, US)
Miconia pustulata (Cuatrecasas 20804, US)
Miconia quintuplinervia (Luteyn 12425, US)
Miconia racemosa (Killip 3860, US)
Miconia radulaefolia (Schultes 19622, US)
Miconia rava (Cuatrecasas 21705, US)
Miconia reclinata (Barr 17704, US)
Miconia reducens (Forero 4772, US)
Miconia resima (Cuatrecasas 8669, US)
Miconia reticulata (Triana 4008, US)
Miconia rhodantha (Bro. Daniel 624, US)
Miconia rigens (Jaramillo 843, US)
Miconia rivetii (Mendoza 883, FMB)
Miconia rubens (Lehmann 5951, US)
Miconia rubiginosa (Cuatrecasas 18710, US)
Miconia rubricans (Silverstone-Sopkin 4456A, US)
Miconia rufescens (Cuatrecasas 4761, US)
Miconia ruficalyx (Lozano 5583, US)
Miconia rugosa (Schultes 15606, US)
Miconia salicifolia (Cuatrecasas 9325, US)
Miconia sandemanii (Sandeman 5837, US)
Miconia schlimii (Smith 13, US)
Miconia sciurea (Gutierrez 2524, US)
Miconia semisterilis (Cuatrecasas 11304, US)
Miconia seposita (Luteyn 12580, US)

Miconia serrulata (Zarucchi 1967, US)
Miconia setosa (Agredo 548, NY)
Miconia shattuckii (Haught 4927, US)
Miconia silverstonei (Silverstone-Sopkin 3849, US)
Miconia simplex (Denslow 2294, US)
Miconia smaragdina (Forero 2941, US)
Miconia smithii (Kirkbride 1952, US)
Miconia sneidernii (Sneidern 5525, photo in US)
Miconia sordida (Triana 4031, US)
Miconia spatellophora (Cuatrecasas 8794-B, US)
Miconia spennerostachya (Rudas 2875, US)
Miconia spicellata (Grubb 46, US)
Miconia spichigeri (Brand 1356, US)
Miconia spinulidentata (Barclay 5938, US)
Miconia spinulosa (Kirkbride 2019, US)
Miconia spireifolia (Cuatrecasas 8520, US)
Miconia splendens (Cuatrecasas 10819, US)
Miconia sprucei (Pinto 1002, COL)
Miconia squamulosa (Garcia-Barriga 20248, US)
Miconia stenostachya (Jaramillo 717, US)
Miconia stenourea (López-Figueiras 8530, US)
Miconia stephananthera (Uribe-Uribe 3602, US)
Miconia stevensiana (Almeda 10402, COL)
Miconia stipitata (Cuatrecasas 16916, US)
Miconia stipularis (Cuatrecasas 18091, US)
Miconia subalpina (Pennell 10520, US)
Miconia submacrophylla (Cuatrecasas 9025, US)
Miconia subsessilifolia (Escobar 3334, US)
Miconia summa (Garcia-Barriga 17690, US)
Miconia superposita (Cuatrecasas 23324, US)
Miconia symlocoidaea (Cited in description)
Miconia tamana (Mendoza 7209, FMB)
Miconia tenuis (Cited by Cogniaux 1891)
Miconia ternatifolia (Garcia-Barriga 20466, US)
Miconia tetrasperma (Schultes 3737, US)
Miconia theizans (Luteyn 13204, US)
Miconia tinifolia (Devia 645, US)
Miconia titanophylla (Garcia-Barriga 18200, US)
Miconia tomentosa (Maguire 44070, US)
Miconia toroi (Toro 278, US)
Miconia tovarensis (Orozco 2497, US)
Miconia traillii (Zarucchi 1762, US)
Miconia transversa (Cuatrecasas 17459, US)
Miconia triangularis (Schultes 2037, US)
Miconia tricaudata (Barclay 6579, US)
Miconia trichophora (Killip 14848, US)
Miconia trinervia (Cuatrecasas 16680, US)
Miconia triplinervis (Thomas 5495, US)
Miconia truncata (Cuatrecasas 6829, US)
Miconia tuberculata (Callejas 2297, US)
Miconia tubulosa (Lawrance 484, US)
Miconia turgida (Uribe-Uribe 6174, US)
Miconia ulmariooides (Grant 10229, NY)

- Miconia umbriensis* (Klug 1926, US)
Miconia uribei (Uribe-Urube 6580, US)
Miconia urticoides (Forero 2456, US)
Miconia vallensis (Cuatrecasas 21667, US)
Miconia velutina (Schlim 545, US)
Miconia verrucosa (Cuatrecasas 27866, US)
Miconia versicolor (Uribe-Urube 6245, US)
Miconia violacea (Lehmann 3842, US)
Miconia voronovii (Cuatrecasas 13164, US)
Miconia wittii (Stergios 4113, US)
Miconia wurdackii (Uribe-Urube 5683, US)
Miconia zarucchii (Zarucchi 1632, US)
- Venezuela (212 spp)**
- Miconia abbreviata* (Liesner 18990, US)
Miconia abysmophila (Maguire 42206, US)
Miconia acinodendron (Bunting 11010, US)
Miconia acutifolia (Steyermark 109698, US)
Miconia aeruginosa (Tillett 744-247, US)
Miconia affinis (Bunting 11599, US)
Miconia aguitensis (Huber 12594, US)
Miconia alata (Fernández 803, US)
Miconia albicans (Steyermark 62359, US)
Miconia alborufescens (Cowan 31490-A, US)
Miconia aliquantula (Valero 33, US)
Miconia alternans (Steyermark 115105, US)
Miconia amacurensis (Rosales 241, US)
Miconia amblyandra (Cited by Michelangeli & Cotton 2008)
Miconia amilcariana (Stergios 17582, US)
Miconia ampla (Zimmermann 104, US)
Miconia apostachya (Liesner 24363, US)
Miconia appendiculata (Tillett 737-150, US)
Miconia araguensis (Wurdack 2688, US)
Miconia arbutifolia (Steyermark 105031, US)
Miconia argyrophylla (Stergios 11556, US)
Miconia aristata (Tate 897, US)
Miconia astroplocama (Cited by Almeda 2009)
Miconia avia (Wurdack 2766, US)
Miconia aymardii (Aymard 4412, US)
Miconia barbinervis (Tillett 737-83, US)
Miconia bernardii (Bernardi 3209, US)
Miconia berryi (Berry 132, US)
Miconia biglandulosa (Aymard 7168, US)
Miconia biglomerata (Cited by Cogniaux 1891)
Miconia bilopezii (López-Figueiras 8688, US)
Miconia borjensis (Huber 486, US)
Miconia boxii (Box 3745, US)
Miconia brachybotrya (Liesner 17481, US)
Miconia brachygyna (Dorr 7699, US)
Miconia bracteata (Liesner 23603, US)
Miconia breteleri (Wurdack 2765, US)
Miconia brevipes (Liesner 17930, US)

- Miconia bubalina* (Bunting 12801, US)
Miconia buxifolia (Steyermark 101147A, US)
Miconia cacumina (Cited by Berry 2001)
Miconia campestris (Huber 10361, US)
Miconia canaguensis (Wurdack 2744, US)
Miconia carassana (Williams 14832, US)
Miconia cataractae (Steyermark 98326, US)
Miconia cautis (Maguire 37498, US)
Miconia centrodesma (Liesner 18196, US)
Miconia ceramicarpa (Steyermark 86513, US)
Miconia cernua (Funck 1557, photo in US)
Miconia chaetodon (Croat 54425, US)
Miconia chamissois (Cited by Michelangeli & Cotton 2008)
Miconia chapensis (Cited in description)
Miconia chionophila (Ruiz-Terán 7991, US)
Miconia chrysophylla (Steyermark 101771, US)
Miconia ciliata (Steyermark 106134, US)
Miconia cinchonifolia (Fernández 7180, US)
Miconia cionotricha (Steyermark 101461, US)
Miconia crassinervia (Bernardi 6630, US)
Miconia curta (Steyermark 93879, US)
Miconia decurrens (Huber 8422, US)
Miconia dioica (Steyermark 1226, US)
Miconia dispar (Liesner 4123, US)
Miconia dodecandra (Davidse 4668, US)
Miconia dolichopoda (Ruiz-Terán 1267, US)
Miconia dolichorrhyncha (Boom 6015, US)
Miconia donaeana (Wurdack 2743, US)
Miconia dunstervillei (Steyermark 100904, US)
Miconia egensis (Fernández 4365, US)
Miconia elaeoides (Wurdack 2785, US)
Miconia elata (Steyermark 123426, US)
Miconia elviraе (Luteyn 9635, US)
Miconia eugenioides (Wurdack 42673, US)
Miconia fallax (Wurdack 34463, US)
Miconia fissa (Cited by Michelangeli & Cotton 2008)
Miconia floribunda (Wurdack 2747, US)
Miconia fragilis (Aymard 4170, US)
Miconia fragrans (Liesner 21538, US)
Miconia funckii (Wurdack 2693, US)
Miconia globuliflora (Cited by Michelangeli & Cotton 2008)
Miconia goudotii (Ruiz-Terán 3460, US)
Miconia gracilis (Liesner 10899, US)
Miconia gratissima (Fernández 5890, US)
Miconia grossidentata (Steyermark 113891, US)
Miconia guaiquinimiae (Maguire 33024, US)
Miconia holosericea (Steyermark 96689, US)
Miconia huberi (Huber 9519, US)
Miconia hypoleuca (Steyermark 96079, US)
Miconia ibaguensis (Sobrevila 79178, US)
Miconia iluensis (Steyermark 98089, US)

- Miconia impetiolaris* (Stergios 15113, US)
Miconia inaequalifolia (Zimmermann 91, US)
Miconia jahnii (Ruiz-Terán 3273, US)
Miconia kappleri (Cited by Michelangeli & Cotton 2008)
Miconia kavanayensis (Lasser 1879, US)
Miconia lacera (Bunting 10991, US)
Miconia laevigata (Maier 2172, US)
Miconia laevipilis (Cardozo 1499, US)
Miconia lamprophylla (Liesner 10764, US)
Miconia larensis (Steyermark 55275, US)
Miconia lasseri (Delgado 604, US)
Miconia lateriflora (Morillo 9197, US)
Miconia latifolia (Alston 6832, US)
Miconia lehmannii (Tillett 746-586, US)
Miconia leiotricha (Berry 1153, US)
Miconia lepidota (Bunting 3779, US)
Miconia liesneri (Liesner 10837, US)
Miconia ligulata (Delascio 3197, US)
Miconia limitaris (Tillett 747-1171, US)
Miconia lonchophylla (Ruiz-Terán 917, US)
Miconia longifolia (Wurdack 2812, US)
Miconia longispicata (Wurdack 43494, US)
Miconia lourteigiana (Steyermark 101449, US)
Miconia lucida (Van der Werff 7630, US)
Miconia macrodon (Delascio 3326, US)
Miconia macrothyrsa (Wurdack 257, US)
Miconia macrotis (Tamayo 2941, US)
Miconia magdalenae (Breteler 4596, US)
Miconia manicata (Cited by Michelangeli & Cotton 2008)
Miconia marginata (Holst 2647, US)
Miconia mariae (Steyermark 1369, US)
Miconia maroana (Williams 14431, US)
Miconia matthaei (Steyermark 107198, US)
Miconia meridensis (Stergios 4736, US)
Miconia mesmeana (Hahn 5002, US)
Miconia metallica (Steyermark 103696, US)
Miconia micrantha (Cited by Michelangeli & Cotton 2008)
Miconia minutiflora (Tillett 737-109, US)
Miconia mirabilis (Liesner 9796, US)
Miconia mituana (Thomas 3314, US)
Miconia mocquerysii (Mocqueris 968, US)
Miconia mollicula (Wurdack 2730, US)
Miconia mulleola (Wurdack 2795, US)
Miconia multiflora (Cited by Michelangeli & Cotton 2008)
Miconia multispicata (Steyermark 127222, US)
Miconia myriantha (Gines 5136, US)
Miconia myrtillifolia (Ruiz-Terán 1185, US)
Miconia nervosa (Steyermark 120532, US)
Miconia nitidissima (Wurdack 2739, US)
Miconia novemnervia (Ruiz-Terán 2000, US)
Miconia paupercula (Funck 122, US)
Miconia perijensis (Wood 92, US)
Miconia perobscura (Croat 59421, US)
Miconia perturbata (Maguire 42474, US)
Miconia phaeophylla (Ramirez 4226, US)
Miconia pisinna (Wurdack 2794, US)
Miconia pluknetii (Liesner 19005, US)
Miconia poeppigii (Bruijn 1213, US)
Miconia polita (Liesner 20946, US)
Miconia prasina (Bunting 11298, US)
Miconia pseudoaplostachya (Maguire 34715, US)
Miconia pseudocapsularis (Fernández 5458, US)
Miconia pterocaule (Liesner 9586, US)
Miconia pubipetala (Morillo 3527, US)
Miconia pulvinata (Wurdack 2726, US)
Miconia punctata (Colella 1558, US)
Miconia pyrifolia (Steyermark 94169, US)
Miconia racemosa (Steyermark 98985, US)
Miconia radulaefolia (Maguire 44075, US)
Miconia reducens (Davidse 18728, US)
Miconia resimoides (Pittier 12985, US)
Miconia rhytidophylla (Liesner 6374, US)
Miconia roraimensis (Maguire 42137, US)
Miconia rubens (Fendler 1833, US)
Miconia rubiginosa (Bunting 9355, US)
Miconia rufescens (Steyermark 108276, US)
Miconia ruficalyx (Steyermark 87292, US)
Miconia rugosa (Liesner 19542, US)
Miconia ruizteranii (Ruiz-Terán 3204, US)
Miconia rupestris (Steyermark 93971, US)
Miconia rupticalyx (Maguire 42013, US)
Miconia salebrosa (Wurdack 2745, US)
Miconia salicifolia (Cited by Wurdack 1973)
Miconia sanctiphilippi (Steyermark 95858, US)
Miconia schlimii (Steyermark 105793, US)
Miconia serialis (Stergios 5364, US)
Miconia serrulata (Bunting 9695, US)
Miconia silicicola (Steyermark 130111, US)
Miconia simplex (Bunting 6800, US)
Miconia smaragdina (Ruiz-Terán 8219, US)
Miconia spicellata (Delascio 3040, US)
Miconia spinulosa (Wurdack 2690, US)
Miconia splendens (Steyermark 120575, US)
Miconia sprucei (Liesner 3688, US)
Miconia squamulosa (Ruiz-Terán 1356, US)
Miconia stenostachya (Pittier 115, US)
Miconia stephananthera (Huber 26960, US)
Miconia steyermarkii (Fernández 8051, US)
Miconia strigosa (Liesner 17000, US)
Miconia suaveolens (Stergios 4708, US)
Miconia superba (Chavie 400, US)
Miconia sylvatica (López-Palacios 1334, US)
Miconia tabayensis (Gehringer 595, US)
Miconia tachirensis (Steyermark 101038, US)
Miconia tamana (Steyermark 100878, US)

Miconia tetraspermoides (Lasser 1874, US)
Miconia theizans (Liesner 12759, US)
Miconia tillettii (Hernández 452, US)
Miconia tinifolia (Ruiz-Terán 6328, US)
Miconia tomentosa (Plowman 2277, US)
Miconia tovarensis (Wurdack 2716, US)
Miconia traillii (Liesner 25740, MO)
Miconia trinervia (Boom 6021, US)
Miconia triplinervis (Breteler 4593, US)
Miconia trujillensis (Steyermark 104813, US)
Miconia truncata (Liesner 9004, US)
Miconia tuberculata (Wurdack 2673, US)
Miconia ulmariooides (Wurdack 2675, US)
Miconia undata (Liesner 6708, US)
Miconia velutina (Cuatrecasas 28352, US)
Miconia versicolor (Maguire 44218, US)
Miconia wittii (Cited in description)
Miconia yatuensis (Maguire 36695, US)

Trinidad-Tobago (42 spp)

Miconia acinodendron (Purseglove 6515, US)
Miconia affinis (Snow 10, NY)
Miconia albicans (Broadway s.n., US)
Miconia ampla (Broadway 5855, NY)
Miconia argyrophylla (Broadway 6055, US)
Miconia barbinervis (Cited by Cogniaux 1891)
Miconia bracteata (Broadway 6034, US)
Miconia bupalina (Cited by Cogniaux 1891)
Miconia centrodesma (Cowan 1480, US)
Miconia chrysophylla (Britton 1193, NY)
Miconia ciliata (Cited by Williams 1934)
Miconia dodecandra (Broadway 6151, NY)
Miconia dolichorrhyncha (Broadway 6136, K)
Miconia fragrans (Snow 13, US)
Miconia holosericea (Snow 2, US)
Miconia hypoleuca (Britton 2401, US)
Miconia ibaguensis (Cited by Williams 1934)
Miconia kappleri (Schultes 4862, NY)
Miconia lacera (Purseglove 6499, US)
Miconia laevigata (Cowan 1377, US)
Miconia lanata (Snow 11, US)
Miconia lateriflora (Cited by Williams 1934)
Miconia longifolia (Adams 14376, NY)
Miconia macrothyrsa (Hekking 1394, US)
Miconia macrotis (Broadway 6625, US)
Miconia matthaei (Britton 1807, NY)
Miconia minutiflora (Broadway 6630, NY)
Miconia mirabilis (Smith 10002, US)
Miconia myriantha (Eggers 1359, US)
Miconia nervosa (Smith 10075, US)
Miconia plukenetii (Cowan 1483, US)
Miconia prasina (Crosby 58, US)
Miconia pubipetala (Eggers 1150, US)

Miconia punctata (Snow s.n., US)
Miconia racemosa (Britton 2134, NY)
Miconia rubiginosa (Cited by Williams 1934)
Miconia ruficalyx (Snow 6, US)
Miconia savannarum (Britton 2468, NY)
Miconia serrulata (Britton 2893, NY)
Miconia splendens (Broadway 7565, NY)
Miconia stenostachya (Britton 696, US)
Miconia tomentosa (Britton 1017, NY)

Guyana (88 spp)

Miconia abbreviata (Clarke 8597, US)
Miconia acinodendron (Jenman 4235, US)
Miconia acutifolia (Hahn 5463, US)
Miconia aeruginosa (Henkel 945, US)
Miconia affinis (Jansen-Jacobs 4260, US)
Miconia alata (Tillett 45382, US)
Miconia albicans (Jansen-Jacobs 3614, US)
Miconia alborufescens (Pipoly 7926, US)
Miconia alternans (Irwin 178, US)
Miconia amacurensis (Maas 3947, US)
Miconia ampla (Tillett 5859, US)
Miconia apostachya (Boom 7744, US)
Miconia argyrophylla (De la Cruz 2052, US)
Miconia brachybotrya (Clarke 8595, CAS)
Miconia bracteata (Cowan 39250, US)
Miconia brevipes (Cowan 39258, US)
Miconia bupalina (Pipoly 8285, US)
Miconia cacumina (Cited in description)
Miconia campestris (Cowan 2082, US)
Miconia carassana (Henkel 4873, US)
Miconia centrodesma (Henkel 1202, US)
Miconia ceramicarpa (Smith 2555, US)
Miconia chrysophylla (Smith 3607, US)
Miconia ciliata (Cowan 1784, US)
Miconia diaphanea (Clarke 7341, US)
Miconia dodecandra (Cowan 1886, US)
Miconia dolichorrhyncha (Clarke 7153, US)
Miconia egensis (De la Cruz 3737, US)
Miconia fallax (Cowan 39260, US)
Miconia fanshawei (Cited in description)
Miconia fragilis (Maguire 40501, US)
Miconia fragrans (Hahn 4327, US)
Miconia francavillana (Henkel 6868, US)
Miconia gratissima (Maguire 23037, US)
Miconia holosericea (Fanshawe 2499, US)
Miconia hypoleuca (Tillett 45879, US)
Miconia ibaguensis (Smith 2440, US)
Miconia impetiolaris (Clarke 7915, CAS)
Miconia lappacea (Clarke 5414, US)
Miconia lasseri (Maguire 32591, US)
Miconia lateriflora (Maguire 22813, US)

- Miconia lepidota* (De la Cruz 4385, US)
Miconia longifolia (Jenman s.n., NY)
Miconia longispicata (Clarke 7196, US)
Miconia loreyoides (Clarke 10382, US)
Miconia macrothyrsa (Pipoly 9487, US)
Miconia maguirei (Boom 7439, US)
Miconia marginata (Maguire 23537, US)
Miconia mariae (Hahn 4273, US)
Miconia matthaei (Pipoly 8173, US)
Miconia minutiflora (Hollowell 724, CAS)
Miconia mirabilis (Gleason 610, US)
Miconia myriantha (Pinkus 217, US)
Miconia nervosa (Stoffers 268, US)
Miconia phaeophylla (Tillett 45801, US)
Miconia platypoda (Clarke 2862, CAS)
Miconia pluknetii (Maguire 22945, US)
Miconia poeppigii (Tiwari 641, US)
Miconia polita (Pipoly 10914, US)
Miconia prasina (Cowan 1761, US)
Miconia pseudoaplostachya (Henkel 5100, US)
Miconia pubipetala (De la Cruz 1038, US)
Miconia punctata (Acevedo-Rodriguez 6086, US)
Miconia pyrifolia (Boom 8127, US)
Miconia racemosa (Pipoly 11728, US)
Miconia radulaefolia (Hoffmann 2283, US)
Miconia regelii (Smith 2721, US)
Miconia roraimensis (Ule 8692, photo in US)
Miconia rubiginosa (Henkel 938, US)
Miconia rufescens (Pipoly 9442, US)
Miconia ruficalyx (De la Cruz 4542, US)
Miconia rugosa (Maguire 32442, US)
Miconia rupestris (Liesner 23227, US)
Miconia rupticalyx (Boom 8972, US)
Miconia sagotiana (Wilson-Browne 315, NY)
Miconia serialis (Jansen-Jacobs 3822, US)
Miconia serrulata (Pipoly 8091, US)
Miconia silicicola (Tate 212, NY)
Miconia splendens (Smith 2814, US)
Miconia stenostachya (Hoffmann 3718, US)
Miconia stephananthera (McDowell 4103, US)
Miconia superba (Tillett 45556, US)
Miconia tetraspermoides (Tillett 45511, US)
Miconia theizans (Clarke 9517, CAS)
Miconia tillettii (Boom 7481, US)
Miconia tinifolia (Liesner 23229, US)
Miconia tomentosa (Pipoly 11542, US)
Miconia traillii (Clarke 4439, CAS)
- Suriname (64 spp)**
- Miconia acinodendron* (Maguire 23745, US)
Miconia acuminata (Hostman 1265, NY)
Miconia affinis (Cited by Wurdack et al. 1993)
Miconia alata (Kramer 2721, US)
- Miconia albicans* (Donselaar 135, US)
Miconia aliquantula (Sauvain 511, US)
Miconia alternans (Hekking 1074, US)
Miconia ampla (Maguire 24087, US)
Miconia apostachya (Maas 3334, US)
Miconia argyrophylla (Maguire 39110, US)
Miconia bracteata (Donselaar 2409, US)
Miconia brevipes (Oldenburger 1130, US)
Miconia cacatia (Mori 8425, US)
Miconia campestris (Teunissen LBB 15843, US)
Miconia ceramicarpa (Lindeman 532, US)
Miconia chrysophylla (Stoffers 477, US)
Miconia ciliata (Maguire 24287, US)
Miconia diaphanea (Irwin 55090, US)
Miconia dodecandra (Maguire 24415, US)
Miconia egensis (Lindeman 6474, US)
Miconia fallax (Oldenburger 752, NY)
Miconia francavillana (Lindeman 486, US)
Miconia gratissima (Donselaar 1496, US)
Miconia holosericea (Kramer 2720, US)
Miconia hypoleuca (B.W. 630, US)
Miconia ibaguensis (Kramer 2064, US)
Miconia kappleri (Cited by Wurdack et al. 1993)
Miconia lappacea (Cited by Wurdack et al. 1993)
Miconia lasseri (Maas LBB 11066, US)
Miconia lateriflora (Maguire 24109, US)
Miconia lepidota (B.W. 5848, US)
Miconia longifolia (Maguire 23844, US)
Miconia macrothyrsa (Oldenburger 571, US)
Miconia matthaei (Maguire 24906, US)
Miconia minutiflora (Lindeman 2512, NY)
Miconia mirabilis (Croat 53855, US)
Miconia myriantha (B.W. 3826, US)
Miconia nervosa (Lindeman LBB 12112, US)
Miconia phaeophylla (Kramer 2960, US)
Miconia platypoda (Boer 1557, NY)
Miconia pluknetii (Lindeman 7022, US)
Miconia poeppigii (Irwin 55450, US)
Miconia prasina (Cowan 39106, US)
Miconia pubipetala (Irwin 55479, US)
Miconia punctata (Cited by Wurdack et al. 1993)
Miconia pyrifolia (Lanjouw 2864, NY)
Miconia racemosa (Maguire 25007, US)
Miconia rubiginosa (Oldenburger 1149, US)
Miconia rufescens (Oldenburger 684, US)
Miconia ruficalyx (LBB 13374, US)
Miconia rugosa (Donselaar 1663, US)
Miconia sagotiana (Maguire 23872, US)
Miconia sarmentosa (Cited by Wurdack et al. 1993)
Miconia serialis (Lanjouw 1278, US)
Miconia serrulata (Maguire 23641, US)
Miconia splendens (Lanjouw 1296, US)
Miconia stenostachya (Rombouts 376, NY)

- Miconia stephananthera* (Cited by Wurdack et al. 1993)
Miconia tetraspermooides (Lindeman 5254, US)
Miconia tillettii (Maguire 24533, US)
Miconia tomentosa (Jongkind 17038, US)
Miconia traillii (Maguire 24689, US)
Miconia trimera (Irwin 54673A, US)
Miconia tschudyoides (Donselaar 1655, US)
- French Guiana (67 spp)**
- Miconia acinodendron* (Halleé 606, US)
Miconia acuminata (De Granville 5352, US)
Miconia affinis (Mori 8697, US)
Miconia alata (Irwin 55279, US)
Miconia albicans (Oldeman 2261, US)
Miconia aliquantula (Mori 18226, US)
Miconia alternans (De Granville 10133, US)
Miconia amacurensis (Cited by Wurdack et al. 1993)
Miconia ampla (Grenand 1302, US)
Miconia apostachya (Cited by Wurdack et al. 1993)
Miconia argyrophylla (De Granville 12492, US)
Miconia bracteata (De Granville 4546, US)
Miconia cacatin (De Granville B-5058, US)
Miconia centrodesma (De Granville 1017, US)
Miconia ceramicarpa (Cremers 7924, US)
Miconia chrysophylla (Cowan 38758, US)
Miconia ciliata (Hoff 5149, US)
Miconia diaphanea (Cremers 6751, US)
Miconia dispar (De Granville B-4757, US)
Miconia egensis (Oldeman 1908, NY)
Miconia elata (Oldeman 3060, US)
Miconia eriodonta (Cremers 5738, US)
Miconia fragilis (Prevost 1086, US)
Miconia francavillana (Cowan 38842, US)
Miconia gratissima (Wachenheim 121, US)
Miconia holosericea (De Granville 6451, US)
Miconia hypoleuca (Lescure 943, US)
Miconia kappleri (Oldeman B-2889, US)
Miconia lappacea (Cremers 9863, US)
Miconia lasseri (Oldeman 1404, US)
Miconia lateriflora (De Granville 6327, US)
Miconia lepidota (Melinon 35, US)
Miconia longifolia (Aubreville 352, US)
Miconia longispicata (Feuillet 10221, US)
Miconia macrothyrsa (Cited by Wurdack et al. 1993)
Miconia matthaei (De Granville 4545, US)
Miconia melinonis (Sastre 6389, US)
Miconia minutiflora (Mori 15622, US)
Miconia mirabilis (Cremers 5739, US)
Miconia mutabilis (Cited by Wurdack et al. 1993)
Miconia myriantha (Cremers 10623, US)
Miconia nervosa (De Granville 5281, US)
Miconia oldemanii (Cremers 8711, US)
Miconia phaeophylla (Wachenheim s.n., US)
- Miconia plukenetii* (De Granville 3498, US)
Miconia poeppigii (Prevost 1005, US)
Miconia prasina (Oldeman B-771, US)
Miconia pseudoaplostachya (Cited by Wurdack et al. 1993)
Miconia pubipetala (Cremers 7474, US)
Miconia punctata (Prevost 2863, US)
Miconia pyrifolia (Cited by Wurdack et al. 1993)
Miconia racemosa (Cremers 5737, US)
Miconia rufescens (Veyret 4394, US)
Miconia ruficalyx (Oldeman B-2344, US)
Miconia sagotiana (Prevost 1118, US)
Miconia sarmentosa (Prevost 1707, US)
Miconia sastrei (Sastre 1365, US)
Miconia serialis (Rech s.n., US)
Miconia serrulata (Mori 8946, US)
Miconia splendens (Halleé 659, US)
Miconia tetraspermooides (Oldeman B-2777, US)
Miconia tillettii (De Granville 4492, US)
Miconia tomentosa (Cremers 6662, US)
Miconia traillii (Oldeman T-893, US)
Miconia trimera (Cremers 6510, US)
Miconia trinervia (Grenand 738, US)
Miconia tschudyoides (Cremers 8378, US)
- Ecuador (254 spp)**
- Miconia abbreviata* (Neill 6730, US)
Miconia aequatorialis (Little 677, US)
Miconia aeruginosa (Harling 11526, US)
Miconia affinis (Lugo 2338, US)
Miconia aggregata (Boeke 384, US)
Miconia albicans (Rose 23379, US)
Miconia alypifolia (Maguire 44312, US)
Miconia amazonica (Renner 69318, US)
Miconia amnicola (Hurtado 3051, US)
Miconia ampla (Palacios 7819, US)
Miconia appendiculata (Jaramillo 2945, US)
Miconia arboricola (Tipaz 1757, US)
Miconia ascendens (Drew E-359, US)
Miconia asclepiadea (Grubb 1413, US)
Miconia aspergillaris (Asplund 17709, US)
Miconia asperrima (Madison 3550, US)
Miconia asplundii (Cited in description)
Miconia aspratilis (Camp E-704, US)
Miconia astroplocama (Madison 3489, US)
Miconia aulocalyx (Harling 7419, US)
Miconia aurea (Camp E-1133, US)
Miconia aureoides (Cazalet 7709, US)
Miconia auritinoda (Luteyn 11007, US)
Miconia barbeyana (Harling 11890, US)
Miconia barbinervis (Lugo 1747, US)
Miconia barbipilis (Steyermark 54308, US)
Miconia barclayana (Cazalet 5444, US)

- Miconia beneolens* (King 7941, US)
Miconia benoistii (Cited in description)
Miconia benthamiana (Mexia 8420, US)
Miconia biglandulosa (Baker 6268, US)
Miconia bipatrialis (Harling 8241, US)
Miconia bolivarensis (Dodson 14316, US)
Miconia bordoncilloana (Neill 13478, CAS)
Miconia brachybotrya (Zaruma 196, US)
Miconia brachycalyx (Palacios 5178, US)
Miconia bracteolata (Maguire 44273, US)
Miconia brevitheca (Gentry 72562, US)
Miconia bubalina (Camp E-1544, US)
Miconia bullata (Jaramillo 7333, US)
Miconia caelata (Jameson s.n., US)
Miconia caesariata (Anderson 12702, US)
Miconia cajanumana (Espinosa E-1993, US)
Miconia calignosa (Cited in description)
Miconia calvescens (Harling 9705, US)
Miconia campii (Camp E-1614, US)
Miconia cannabina (Brandbyge 30523, US)
Miconia capitellata (Harling 13684, US)
Miconia carassana (Jaramillo 30935, US)
Miconia castillensis (Camp E-5162, US)
Miconia castrensis (Camp E-4728, US)
Miconia cazaletii (Harling 6990, US)
Miconia centrodesma (Lugo 4565, US)
Miconia centrodesmoides (Hurtado 184, US)
Miconia centronioides (Little 6406, US)
Miconia centrophora (Lehmann 6232, photo in US)
Miconia cercophora (Lugo 1573, US)
Miconia chionophila (Barclay 8354, US)
Miconia chlorocarpa (Palacios 3953, US)
Miconia chrysophylla (Brandbyge 35363, US)
Miconia ciliata (Játiva 756, US)
Miconia cladonia (Steyermark 53604, US)
Miconia clathrantha (Harling 26739, US)
Miconia collayensis (Camp E-4890, US)
Miconia conformis (Neill 6696, US)
Miconia corazonica (Camp E-1664, US)
Miconia corymbiformis (Moran 47, US)
Miconia cosangensis (Kirkbride 4160, US)
Miconia crebribullata (Dobson 1080, US)
Miconia crinita (Boeve 552, US)
Miconia crocea (Lugo 487, US)
Miconia cuneata (Cited by Renner 1999)
Miconia cuprea (Camp E-4736, US)
Miconia cutucuensis (Camp E-1297, US)
Miconia dapsiliflora (Cited in description)
Miconia decurrens (Harling 13912, US)
Miconia denticulata (Cited by Wurdack 1980)
Miconia dielsii (Maas 2972, US)
Miconia difficilis (Tate 554, US)
Miconia dissimulans (Maguire 44366, US)
Miconia divergens (Rubio 921, US)
Miconia dodecandra (Asplund 18866, US)
Miconia dodsonii (Maguire 44365, US)
Miconia dolichorrhyncha (Brandbyge 30333, US)
Miconia dorsiloba (Camp E-1150, US)
Miconia dudleyi (Harling 9987, US)
Miconia egensis (Cited by Ulloa & Neill 2005)
Miconia elata (Neill 7133, US)
Miconia erioclada (Harling 12925, US)
Miconia espinosae (Harling 20594, US)
Miconia explicita (Clarck 2553, US)
Miconia filamentosa (Lawesson 43343, US)
Miconia floribunda (Palacios 6404, US)
Miconia fosbergii (Kessler 2620, US)
Miconia fosteri (Foster 3617, US)
Miconia fuliginosa (Prieto P-314, US)
Miconia gibba (Palacios 6211, US)
Miconia glandulistyla (Kirkbride 4293, US)
Miconia glaucescens (Palacios 6586, US)
Miconia glyptophylla (Camp E-4565, US)
Miconia gonioclada (Camp E-225, US)
Miconia goniostigma (Mexia 6687, US)
Miconia gracilis (Játiva 791, US)
Miconia grandifoliata (Little 727, US)
Miconia grayana (Steyermark 52931, US)
Miconia guayaquilensis (Dodson 6400, US)
Miconia hadrophylla (Hoover 2205, US)
Miconia harlingii (Palacios 3954, US)
Miconia heterochaeta (Lugo 5431, US)
Miconia hexamera (Harling 23011, US)
Miconia hirsutivena (Cited in description)
Miconia hookeriana (Lugo 3411, US)
Miconia huigrensis (Camp E-4015, US)
Miconia hylophila (Brandbyge 33885, US)
Miconia hymenanthera (Davis 291, US)
Miconia ibaguensis (Harling 25146, US)
Miconia idiogena (Palacios 7178, US)
Miconia imitans (Ollgaard 35593, US)
Miconia impetiolaris (Dodson 8760, US)
Miconia inanis (Lehmann 5658, photo in US)
Miconia innata (Camp E-728, US)
Miconia insularis (Harling 9923, US)
Miconia intricata (Gentry 70098, US)
Miconia jahnii (Palacios 3926, US)
Miconia jorgensenii (Jorgensen JP-58, US)
Miconia kraenzlinii (Palacios 6794, US)
Miconia lacera (Játiva 597, US)
Miconia laevigata (Játiva 6, US)
Miconia lamprophylla (Palacios 8727, US)
Miconia lasiocalyx (Little 6547, US)
Miconia lateriflora (Romoleroux 2140, US)
Miconia latifolia (Luteyn 14503, US)
Miconia laxa (Cited in description)

- Miconia leandroides* (Cited in description)
Miconia ledifolia (Maguire 44319, US)
Miconia lennartanderssonii (Harling 19235, US)
Miconia ligustrina (Camp E-5143, US)
Miconia littlei (Gentry 12176, US)
Miconia livida (Madison 3343, US)
Miconia longifolia (Játiva 11, US)
Miconia longisetosa (Sodiro 490, US)
Miconia loreyoides (Palacios 6046, US)
Miconia lourteigiana (Ellenberg 3268, US)
Miconia loxensis (Cited by Wurdack 1980)
Miconia lugonis (Lugo 1711, US)
Miconia lutescens (Maguire 44344, US)
Miconia macbrydeana (MacBryde 983, US)
Miconia machinazana (Neill 16909, NY)
Miconia macrotis (Hitchock 21315, US)
Miconia manicata (Dorr 6362, US)
Miconia matthaei (Cerón 3094, US)
Miconia mazanana (Harling 7494, US)
Miconia media (Camp E-5237A, US)
Miconia mediocris (Camp E-705, US)
Miconia medusa (Espinosa 1559, US)
Miconia micropetala (Maguire 44367, US)
Miconia minuta (Baker 5552, US)
Miconia minutiflora (Bennett 3785, US)
Miconia multiplinervia (Tipaz 1042, US)
Miconia multispicata (Palacios 6491, US)
Miconia namandensis (Espinosa E-1077, US)
Miconia napoana (Lugo 2686, US)
Miconia nasella (Camp E-1324, US)
Miconia nervosa (Grubb 1440, US)
Miconia nodosa (Luteyn 8419, US)
Miconia notabilis (Cited by Wurdack 1980)
Miconia nutans (Neill 6945, US)
Miconia obscura (Maguire 44335, US)
Miconia ochracea (Asplund 17104, US)
Miconia oellgaardii (Cited in description)
Miconia oligantha (Lugo 22, photo in US)
Miconia ombrophila (Little 632, US)
Miconia onaensis (Prieto P-211, US)
Miconia onychocalyx (Cited in description)
Miconia oraria (Little 6306, US)
Miconia orcheotoma (Palacios 10552, US)
Miconia oreescia (Palacios 7272, US)
Miconia pailasana (Camp E-4902, US)
Miconia paleacea (Palacios 10307, US)
Miconia paludigena (Luteyn 11207, NY)
Miconia papillosa (Maguire 44239, US)
Miconia pastazana (Lugo 1465, US)
Miconia pausana (Camp E-1603, US)
Miconia penningtonii (Cazalet 5602, US)
Miconia pernettifolia (Steyermark 54344, US)
Miconia phaeochaeta (Hurtado 2391, US)
- Miconia phlebodes* (Baker 5472, US)
Miconia pichinchensis (Asplund 9791, US)
Miconia pilaloensis (Holm-Nielsen 24407, US)
Miconia pisinniflora (Asplund 18591, US)
Miconia poeppigii (Palacios 7724, US)
Miconia poortmannii (Hitchcock 21339, US)
Miconia porphyrotricha (Asplund 10163, US)
Miconia prasina (Bennett 3662, US)
Miconia prietoi (Maguire 44287, US)
Miconia procumbens (Lugo 3245, US)
Miconia prominens (Camp E-791, US)
Miconia protuberans (Harling 16433, US)
Miconia pseudocentrophora (Lugo 511, US)
Miconia pterocaulon (Jaramillo 31302, US)
Miconia pujana (Lugo 418, US)
Miconia punctata (Palacios 6612, US)
Miconia pustulata (Mexia 7707, US)
Miconia quadripora (Little 205, US)
Miconia radula (Dodson 780, US)
Miconia radulaefolia (Brandbyge 36031, US)
Miconia reburrosa (Camp E-1604, US)
Miconia reducens (Játiva 2111, US)
Miconia renneri (Cited in description)
Miconia rhonhofiae (Cited in description)
Miconia rimbachii (Rimbach 786, US)
Miconia rivalis (Little 455, US)
Miconia rivetii (Maas 2859, US)
Miconia robinsoniana (Bentley 144, US)
Miconia rotundifolia (Wurdack 1199, US)
Miconia ruizii (Madsen 75872, US)
Miconia rupticalyx (Cited by Wurdack 1980)
Miconia russea (Harling 12812, US)
Miconia salicifolia (Cazalet 5355, US)
Miconia scabra (Cited by Cogniaux 1891)
Miconia schunkei (Holm-Nielsen 4386, US)
Miconia sciurea (Palacios 639, US)
Miconia scutata (Alvarado 138, US)
Miconia seposita (Lugo 5903, US)
Miconia serrulata (Mexia 7067, US)
Miconia seticaulis (Harling 17633, US)
Miconia smaragdina (Cited in Wurdack 1980)
Miconia sodiroi (Cited in description)
Miconia sparrei (Sparre 19290, US)
Miconia spennerostachya (Dodson 15083, US)
Miconia spinulidentata (Drew E-322, US)
Miconia splendens (Harling 7379, US)
Miconia stelligera (Cited by Renner 1999)
Miconia stenophylla (Prieto P-312, US)
Miconia stipitata (Cited by Wurdack 1980)
Miconia suborbicularis (Maguire 44364, US)
Miconia subspicata (Brandbyge 32215, US)
Miconia tenensis (Harling 12858, US)
Miconia tephrodes (Camp E-4872, US)

- Miconia ternatifolia* (Harling 14049, US)
Miconia theizans (Dodson 1335, US)
Miconia tinifolia (Camp E-325, US)
Miconia tomentosa (Brandbyge 30616, US)
Miconia transversa (Van der Werff 11878, US)
Miconia triangularis (Asplund 9190, US)
Miconia trinervia (Lugo 3064, US)
Miconia triplinervis (Ollgaard 34915, US)
Miconia vaccinioides (Ollgaard 58226, US)
Miconia venulosa (Asplund 18898, US)
Miconia versicolor (Palacios 9646, US)
Miconia vesca (Steyermark 53582, US)
Miconia villonacensis (Harling 13845, US)
Miconia zamorensis (Cited in description)
Miconia zubenetana (Gudiño 1682, US)
- Peru (302 spp)**
- Miconia abbreviata* (Smith 990, NY)
Miconia acinodendron (Smith 1427, US)
Miconia acuminata (Klug 3342, US)
Miconia adinantha (Dudley 10384, US)
Miconia adrieni (Young 659, NY)
Miconia affinis (Schunke 4987, US)
Miconia aggregata (Dillon 6425, US)
Miconia albicans (Killip 24777, US)
Miconia aligera (Dudley 10612, US)
Miconia alpina (Cano 3838, US)
Miconia alternans (McDaniel 16824, NY)
Miconia alypifolia (Ochoa 11675, NY)
Miconia amazonica (Smith 3898, US)
Miconia amnicola (Killip 27530, US)
Miconia ampla (Schunke 5703, US)
Miconia amplexicaulis (Mexia 8288, US)
Miconia appendiculata (Klug 2365, US)
Miconia aprica (Killip 24246, US)
Miconia argyrophylla (Foster 10874, US)
Miconia asclepiadea (Rodríguez 2776, MO)
Miconia aspergillaris (Diaz S. 2160, NY)
Miconia asperrima (Barbour 3960, US)
Miconia astroplocama (McDaniel 10780, US)
Miconia atrofusca (Macbride 5188, NY)
Miconia aulocalyx (Rimachi 1830, NY)
Miconia aurea (Schunke 2902, US)
Miconia aureoides (Schunke 13752, US)
Miconia axinaeoides (Madison 10237-70, US)
Miconia ayacuchensis (Dudley 11992, US)
Miconia bailloniana (Schunke 6034, US)
Miconia bangii (McCarrol 74, US)
Miconia barbeyana (Foster 7715, US)
Miconia barbinervis (Klug 2619, US)
Miconia beneolens (Dillon 6178, US)
Miconia benthamiana (Dillon s.n., US)
Miconia biglandulosa (Wurdack 1839, US)
- Miconia bipatrialis* (Wurdack 914, US)
Miconia brachyanthera (Wurdack 1058, US)
Miconia brachybotrya (Croat 19545, US)
Miconia bracteolata (Sagástegui 9530, US)
Miconia brevis (Emperaire 5637, US)
Miconia brevistylis (Weberbauer 3553, photo in US)
Miconia bubalina (Smith 1231, US)
Miconia bullata (Wurdack 1705, US)
Miconia caelata (Ochoa 14803, US)
Miconia caerulea (Williams 7757, US)
Miconia cajanumana (Cano 1507, US)
Miconia calvescens (Barbour 5018, US)
Miconia cannabina (Klug 1976, US)
Miconia carassana (Plowman 6960, US)
Miconia carpishana (Maguire 44436, US)
Miconia cauingia (Hutchison 6159, US)
Miconia cazaletii (Cited by Ulloa et al. 2004)
Miconia centrodesma (Bell 88-68, US)
Miconia centrodesmoides (Davidson 9267, US)
Miconia centrophora (Hutchison 5791, US)
Miconia chionophila (Sánchez 780, US)
Miconia choriophylla (Dudley 10405, US)
Miconia chrysophylla (Williams 4809, US)
Miconia ciliaris (Schunke 6001, US)
Miconia ciliata (Knapp 8271, US)
Miconia clathrantha (Barbour 2629, US)
Miconia clivorum (Wurdack 1794, US)
Miconia coelestis (Foster 7565, US)
Miconia comosa (Spruce 4438, photo in US)
Miconia condylata (Foster 10356, US)
Miconia cookii (Vargas 13979, US)
Miconia cordata (Metcalf 30602, US)
Miconia crassifolia (Wurdack 516, US)
Miconia crassipes (Wurdack 827, NY)
Miconia crassistigma (López 5393, US)
Miconia cremophylla (Smith 4467, US)
Miconia cretacea (Plowman 4881, US)
Miconia cuneata (Schunke 11661, NY)
Miconia cuspidata (Wallnöfer 110-20388, US)
Miconia cyanocarpa (Smith 7650, US)
Miconia dasyclada (Madison 10097-70, US)
Miconia decurrens (Schunke 5904, US)
Miconia demissifolia (Madison 10168, US)
Miconia densifolia (Hutchison 5064, US)
Miconia denticulata (Dillon 6268, US)
Miconia diaphanea (Gentry 19755, US)
Miconia dichrophylla (Asplund 13415, G)
Miconia dielsii (Smith 4610, US)
Miconia dipsacea (Killip 22411, US)
Miconia dispar (Schunke A121, US)
Miconia divisoriana (Ferreyra 1669, US)
Miconia dodecandra (Hutchison 5994, US)
Miconia dolichorrhyncha (Vásquez 3896, US)

- Miconia donaeana* (Smith 5437, US)
Miconia dorsiloba (McDaniel 29933, US)
Miconia duckei (Rimachi 7103, US)
Miconia dudleyi (Dudley 11489, US)
Miconia dumetosa (Wurdack 1433, NY)
Miconia egensis (Ayala 2710, US)
Miconia egregia (Díaz 2881, US)
Miconia elata (Belshaw 3472, US)
Miconia elongata (Boeke 3231, NY)
Miconia emendata (Killip 25974, US)
Miconia eriocalyx (Croat 17910, NY)
Miconia erioclada (Barbour 4398, US)
Miconia expansa (Klug 3201, US)
Miconia falcata (Weberbauer 4998, US)
Miconia fallax (Killip 24775, US)
Miconia ferreyrae (Allard 22073, US)
Miconia fictilis (Macbride 4431, NY)
Miconia filamentosa (Klug 938, US)
Miconia firma (López 7772, US)
Miconia flaccida (Killip 25625, US)
Miconia floccosa (Smith 2786, US)
Miconia floribunda (Wurdack 1057, US)
Miconia fosteri (Schunke 3921, US)
Miconia fruticulosa (Weberbauer 3383, photo in US)
Miconia galactantha (Smith 7869, US)
Miconia glaberrima (Macbride 5771, US)
Miconia glandulifera (Mathews 1729 or 680, photo in US)
Miconia glaucescens (Wachter 5, US)
Miconia glomerata (Killip 25934, US)
Miconia glutinosa (Weberbauer 1325, photo in US)
Miconia grandifoliata (McDaniel 16259, US)
Miconia griffisii (Macbride 4382, photo in US)
Miconia grisea (Weberbauer 3346, photo in US)
Miconia hamata (Gentry 23006, US)
Miconia herrerae (Nuñez 8557, US)
Miconia heterochaeta (Kukijat 267, US)
Miconia heteromera (Foster 9319, US)
Miconia hexapetala (Smith 375, US)
Miconia histothrix (Schunke 7553, US)
Miconia holosericea (Killip 24766, US)
Miconia hookeriana (Wurdack 2019, US)
Miconia hospitalis (Dudley 11853, US)
Miconia huanucensis (Asplund 13173, photo in US)
Miconia hutchisonii (Sagástegui 8581, US)
Miconia hygrophila (Killip, US)
Miconia ibaguensis (Macbride 5488, US)
Miconia icosandra (Killip 25781, US)
Miconia idroboi (Ancuash 8, US)
Miconia igniaria (Cano 1505, US)
Miconia impetiolaris (Cited by Almeda 2009)
Miconia inamoena (Ule 6735, US)
Miconia innata (Foster 7772, US)
Miconia juruensis (McDaniel 16319, US)
Miconia klugii (Klug 1141, US)
Miconia koepckeana (Wolfe 12413, US)
Miconia lacera (Klug 2230, US)
Miconia lachnoclada (Madison 10378, US)
Miconia laciniata (Sagástegui 12994, US)
Miconia laevigata (Simpson 558, US)
Miconia lambayequensis (Quiroz 1976, US)
Miconia lamprophylla (Schunke 3441, US)
Miconia lanata (Gentry 57512, US)
Miconia lasiocalyx (Davison 9226, US)
Miconia lasiostyla (Evinger 470, US)
Miconia lateriflora (Ferreyra 17260, US)
Miconia latifolia (Friedberg 236, US)
Miconia laurina (Pavón s.n., photo in US)
Miconia lechleri (Lechler 2565, photo in US)
Miconia lepidota (Cited by Cogniaux 1891)
Miconia lilacina (Pearce s.n., photo in US)
Miconia livida (Schunke 11894, US)
Miconia longifolia (Killip 29777, US)
Miconia longiracemosa (Williams 4249, US)
Miconia longisepala (Klug 3233, US)
Miconia loretensis (Rimachi 4096, US)
Miconia loreyoides (Vargas 11020, US)
Miconia lourteigiana (Dudley 11751, US)
Miconia loxensis (López 7715, US)
Miconia lugonis (Ruokoleinen 3012, US)
Miconia lugubris (Asplund s.n., US)
Miconia lutescens (Dillon 6284, US)
Miconia macrothyrsa (Chrostowski 70-126, US)
Miconia macrotis (Barrier 588, US)
Miconia madisonii (Madison 10295, US)
Miconia malatestae (Macbride 5176, US)
Miconia mapirensis (Wallnöfer 11-29788, US)
Miconia martiniana (Williams 7212, US)
Miconia matthaei (Smith 1062, US)
Miconia mazanana (Vásquez 7223, US)
Miconia media (Hutchison 5728, US)
Miconia membranacea (Foster 9192, US)
Miconia micropetala (Smith 2763, US)
Miconia miles-morganii (Boeke 2032, US)
Miconia minutiflora (Klug 936, US)
Miconia mirabilis (Cited by Wurdack et al. 1993)
Miconia modica (Weberbauer 7552, photo in US)
Miconia molinopampana (Wurdack 1428, US)
Miconia mollis (Gentry 45564, US)
Miconia monzoniensis (Young 580, US)
Miconia multispicata (Gentry 29484, US)
Miconia muricata (Schunke 9747, US)
Miconia myriantha (Killip 27346, US)
Miconia myrtiformis (Smith 8715, US)
Miconia napoana (Vásquez 853, NY)
Miconia nerifolia (Hutchison 5529, US)

- Miconia nervosa* (Knapp 8260, US)
Miconia nigricans (Wurdack 1261, US)
Miconia nitida (Pavón s.n., photo in US)
Miconia nobilis (Killip 26006, US)
Miconia nutans (Schunke 7798, US)
Miconia obscura (Poeppig 34, US)
Miconia opacifolia (Rimachi 4925, US)
Miconia ottikeri (Macbride 4430, US)
Miconia pachydonta (Klug 2143, US)
Miconia paeminoza (Wallnöfer 12-24388, US)
Miconia paleacea (Smith 3994, US)
Miconia paludigena (Wurdack 1218, US)
Miconia pandurata (Spruce s.n., photo in US)
Miconia paradisica (Sagástegui 5962, US)
Miconia pavoniana (Asplund s.n., US)
Miconia pedicellata (Wojtkowski 8231, US)
Miconia persicariifolia (Plowman s.n., US)
Miconia peruviana (Vargas 6872, US)
Miconia phaeophylla (Vásquez 7701, US)
Miconia phanerostila (Vásquez 9667, US)
Miconia phlebodes (Wurdack 1965, US)
Miconia pisinniflora (Cited by Ulloa et al. 2004)
Miconia plumifera (Pearce s.n., photo in US)
Miconia poeppigii (Huashikat 1185, US)
Miconia polychaeta (Madison 10112-70, US)
Miconia polyneura (Cited by Macbride 1941)
Miconia polytopica (Dudley 10533, US)
Miconia prancei (Spichiger 3002, US)
Miconia prasina (Croat 19780, US)
Miconia procumbens (Nuñez 6125, US)
Miconia pseudocentrophora (Rauh 2188, NY)
Miconia pterocaule (Schunke 10088, US)
Miconia puberula (Schunke 5890, US)
Miconia pubipetala (Cited by Macbride 1941)
Miconia pujana (Berlin 240, US)
Miconia pulgari (Macbride 4145, photo in US)
Miconia pulverulenta (Young 1539, US)
Miconia punctata (Smith 6152, US)
Miconia punicea (Dudley 10663, US)
Miconia pyrifolia (Vásquez 251, US)
Miconia quadrifolia (Mathews 1724, US)
Miconia quadripora (Tillett 673-165, US)
Miconia radula (Smith 7782, US)
Miconia radulaefolia (Revilla 75, US)
Miconia reflexipila (Killip 25908, US)
Miconia retusa (Ule 6366, US)
Miconia rimachii (Vásquez 1032, US)
Miconia riparia (Wasshausen 1061, US)
Miconia rivalis (Foster 7861, US)
Miconia rivetii (Cited by Ulloa et al. 2004)
Miconia rotundifolia (Wurdack 1199, US)
Miconia rubiginosa (Wojtkowski 7239, US)
Miconia rufescens (Weberbauer 6259, US)
Miconia rufiramea (Dudley 11143, US)
Miconia rugifolia (Dudley 10705, US)
Miconia rugosa (Gentry 31625, US)
Miconia ruizii (Soukoup 2410, US)
Miconia russea (Smith 5989, US)
Miconia salicifolia (Vargas 4299, US)
Miconia saltuensis (Hutchison 5935, US)
Miconia sanguinea (Vargas 9813, US)
Miconia saxatilis (Macbride 4481, photo in US)
Miconia schunkei (Schunke 5756, US)
Miconia secundifolia (Wurdack 1419, US)
Miconia semisterilis (Huashikat 409, US)
Miconia serrulata (Foster 7866, US)
Miconia setulosa (Peyton 1336, US)
Miconia sparrei (Morawetz 18-22888, US)
Miconia spennerostachya (Dudley 10107, US)
Miconia spichigeri (Spichiger 2000, US)
Miconia splendens (Foster 9335, US)
Miconia stelligera (Plowman 5855, US)
Miconia stenostachya (Aguilar 1036, CAS)
Miconia subandicola (Belshaw 3396, US)
Miconia subglabra (Spruce 4842, photo in US)
Miconia subspicata (Huashikat 571, US)
Miconia subulipetala (Rimachi 3421, US)
Miconia sulcata (Schunke 423, US)
Miconia tenensis (Knapp 8052, US)
Miconia terborghii (Dudley 10794, US)
Miconia terera (Foster 9159, US)
Miconia ternatifolia (Wojtkowski 415, US)
Miconia tetragona (Gentry 20753, US)
Miconia tetrasperma (Gentry 21202, US)
Miconia thaminantha (Dudley 10531, US)
Miconia theizans (Madison 10232-70, US)
Miconia thyrsiflora (Dudley 11157, US)
Miconia thysanophylla (Kayap 1414, US)
Miconia tiliifolia (Scott 882, US)
Miconia tiri (Wojtkowski 7998, US)
Miconia tomentosa (Plowman 5091, US)
Miconia traillii (Rimachi 5088, US)
Miconia triangularis (Gentry 45130, US)
Miconia trichogona (Macbride 5178, US)
Miconia trinervia (Ferreyra 44267, US)
Miconia triplinervis (Schunke 1336, US)
Miconia urbaniana (Weberbauer 5012, photo in US)
Miconia uvida (Wurdack 2153, US)
Miconia vaccinoides (Weberbauer 3998, US)
Miconia vargasii (Metcalf 30774, US)
Miconia venulosa (Schunke 12020, US)
Miconia vitiflora (Macbride 5165, US)
Miconia vittata (Klug 2873, US)
Miconia wagneri (Schunke 5478, US)
Miconia weberbaueri (Wolfe 12475, US)
Miconia wolfei (Wallnöfer 11-20688, US)

Miconia woytkowskii (Foster 9313, US)
Miconia zubenetana (Klug 2208, US)

Bolivia (141 spp)

Miconia abbreviata (Smith 13026, US)
Miconia acuminata (Beck 8054, US)
Miconia aeruginosa (Buchtien 1709, US)
Miconia affinis (Izawa 1, US)
Miconia albicans (Coimbra 754, US)
Miconia alborufescens (Rodriguez 569, US)
Miconia alternans (Anderson 12046, US)
Miconia amabilis (Buchtien 1084, US)
Miconia amissa (Solomon 9554, US)
Miconia ampla (Daly 6376, US)
Miconia argyrophylla (Killeen 5788, US)
Miconia aureoides (Wasshausen 541, US)
Miconia axinaeoides (Boeke 1470, US)
Miconia bangii (Bang 828, US)
Miconia barbeyana (Acevedo-Rodriguez 6603, US)
Miconia barbinervis (Cárdenas 703, US)
Miconia biacuta (Lewis 40839, US)
Miconia biappendiculata (Cited by Cogniaux 1891)
Miconia biformis (Nee 42036, US)
Miconia boliviensis (Lewis 881105, US)
Miconia boomii (Boom 4478, US)
Miconia brittonii (Luteyn 13704, US)
Miconia bubalina (Vargas 2499, US)
Miconia calvescens (Solomon 7909, US)
Miconia centrodesma (Buchtien 1721, US)
Miconia chamissoi (Nee 40469, US)
Miconia chionophila (Solomon 18947, US)
Miconia chrysophylla (Nee 41415, US)
Miconia ciliaris (Smith 14005, US)
Miconia ciliata (Killeen 6503, US)
Miconia coelestis (Cited by Cogniaux 1891)
Miconia collatata (Seidel 5759, US)
Miconia cookii (Buchtien 983, US)
Miconia cordata (Beck 2182, US)
Miconia cremophylla (Cited by Cogniaux 1891)
Miconia cretacea (Seidel 4851, US)
Miconia cuneata (Altamirano 1884, MO)
Miconia cyanocarpa (Solomon 12236, US)
Miconia decurrens (Krukoff 10725, US)
Miconia dichrophylla (Bang 2650, C)
Miconia dispar (Vargas 635, US)
Miconia dodecandra (Solomon 7552, US)
Miconia dolichorrhyncha (Daly 6661, US)
Miconia duckei (Cárdenas 7342, US)
Miconia elata (Nee 39432, US)
Miconia elongata (Solomon 12501, US)
Miconia eriocalyx (Croat 51277, US)
Miconia erioclada (Smith 13807, US)

Miconia fallax (Beck 13649, US)
Miconia ferruginata (Daly 6312, NY)
Miconia flavesrens (Lewis 37321, US)
Miconia fosteri (Vargas 963, US)
Miconia galactantha (Lewis 38933, US)
Miconia glomerulifera (Bang 2856, US)
Miconia gratissima (Anderson 11962, US)
Miconia heliotropoides (Hopkins 244, NY)
Miconia herzogii (Killeen 4417, US)
Miconia hirta (Seidel 1258, US)
Miconia holosericea (Killeen 4477, US)
Miconia hygrophila (Solomon 18515, US)
Miconia ibaguensis (Beck 12281, US)
Miconia incachacana (Steinbach 9459, US)
Miconia ioneura (Steinbach 8337, NY)
Miconia juruensis (Solomon 4627, US)
Miconia klugii (Boom 4294, US)
Miconia kuntzei (Kunze s.n., photo in US)
Miconia lamprophylla (Cited by Almeda 2009)
Miconia lanata (Rusby 2303, US)
Miconia lasiocalyx (Beck 3091, US)
Miconia lateriflora (Vargas 605, US)
Miconia latifolia (Cited by Wurdack 1980)
Miconia latistigma (Cárdenas 4589, US)
Miconia lepidota (Cited by Cogniaux 1891)
Miconia livida (Rusby 2296, US)
Miconia longicuspida (Herzog 326, Z)
Miconia longifolia (Mostacedo 36, US)
Miconia longispicata (Guillén 1259, US)
Miconia loretensis (Beck 20656, US)
Miconia lourteigiana (Prance 7569, US)
Miconia macrothyrsa (Nee 41460, US)
Miconia mandonii (Solomon 7391, US)
Miconia mapirensis (Krukoff 11215, US)
Miconia matthaei (Nee 38699, US)
Miconia micrantha (Solomon 9328, US)
Miconia micropetala (Cited by Wurdack 1980)
Miconia minutiflora (Buchtien 1134, US)
Miconia molybdea (Solomon 14211, US)
Miconia multiflora (Kessler 5766, US)
Miconia multispicata (Solomon 14832, US)
Miconia muricata (Killeen 3731, US)
Miconia myriantha (Anderson 12009, US)
Miconia nervosa (Izawa 49, US)
Miconia obliqua (Beck 4905, US)
Miconia pentlandii (Feuerer 8551a, US)
Miconia persicariifolia (Bang 1410, US)
Miconia phanerostila (Prance 8367, US)
Miconia plumifera (Lewis 881163, US)
Miconia poeppigii (Gentry 44344, US)
Miconia polygama (Beck 3730, US)

- Miconia polytopica* (Buchtien 5498, US)
Miconia prasina (Nee 39636, US)
Miconia puberula (Williams 5376, US)
Miconia pubicalyx (Buchtien 999a, photo in US)
Miconia punctata (Beck 16377, US)
Miconia pyrifolia (Killeen 5834, US)
Miconia quadrialata (Beck 25988, US)
Miconia recondita (Lewis 40838, US)
Miconia resima (Bang 2626, US)
Miconia rosea (Cárdenas 762, US)
Miconia rubiginosa (Beck 21187, US)
Miconia rufescens (Gentry 75514, US)
Miconia ruficalyx (Pennington 60, US)
Miconia ruizii (Perry 1011, US)
Miconia rupticalyx (Solomon 9257, US)
Miconia rusbyana (Rusby 2259, US)
Miconia sanguinea (Beck 13635, US)
Miconia serrulata (Solomon 17695, US)
Miconia sessilifolia (Feuerer 8885b, US)
Miconia setulosa (Beck 7586, US)
Miconia spennerostachya (Solomon 10770, US)
Miconia splendens (Killeen 4474, US)
Miconia staphidioides (Solomon 14839, US)
Miconia steinbachii (Steinbach 367, US)
Miconia stelligera (Williams 621, NY)
Miconia stellulata (Beck 16331, US)
Miconia stenocardia (Cited by Cogniaux 1913)
Miconia stenostachya (Nee 39702, US)
Miconia stephananthera (Helme 723, US)
Miconia subandicola (Beck 8210, US)
Miconia ternatifolia (Solomon 12209, US)
Miconia tetrasperma (Krukoff 10450, US)
Miconia theizans (Solomon 17464, US)
Miconia tiliifolia (Helme 757, US)
Miconia tomentosa (Beck 13696, US)
Miconia trinervia (Nee 43252, US)
Miconia triplinervis (Seidel 2240, US)
Miconia undata (Krukoff 11206, US)
Miconia uvifera (Weddell 4284, photo in US)
Miconia valida (Bang 2630, photo in US)
Miconia vilhenensis (Thomas 5609, US)
Miconia wagneri (Cárdenas 7342, NY)
- Brazil North (162 spp)**
- Miconia abbreviata* (Prance 9714, US)
Miconia acinodendron (Renner 420, US)
Miconia acuminata (Lowrie 311, US)
Miconia acutifolia (Steyermak 104000, US)
Miconia affinis (Lowrie 275, US)
Miconia alata (Mori 17270, US)
Miconia albicans (Rosa 456, US)
Miconia alborufescens (Prance 9599, US)
Miconia aliquantula (Kuhlmann 1970, US)
- Miconia alternans* (Archer 7989, US)
Miconia amacurensis (Daly 4039, US)
Miconia amazonica (Krukoff 8056, US)
Miconia amnicola (Mori 9153, US)
Miconia ampla (Pires 11505, US)
Miconia apostachya (Madison 6317, US)
Miconia argyrophylla (Ratter 6321, US)
Miconia astrotricha (Martius s.n., M)
Miconia aulocalyx (Renner 504, US)
Miconia aurea (Maas P12832, US)
Miconia aureoides (Prance 12166, US)
Miconia barbinervis (Krukoff 5120, US)
Miconia biglandulosa (Maas P12778, US)
Miconia bracteata (Irwin 47512, US)
Miconia brevipes (Sette-Silva 724, US)
Miconia bubalina (Vieira 857, US)
Miconia burchellii (Souza 24454, UPCB)
Miconia calvescens (Sperling 6023, US)
Miconia campestris (Pires 3949, US)
Miconia cannabina (Prance 23944, US)
Miconia carassana (Rodrigues 1599, US)
Miconia cautis (Fróes 28127, US)
Miconia centrodesma (Prance 9597, US)
Miconia ceramicarpa (Martinelli 6754, US)
Miconia chrysophylla (Daly 1934, US)
Miconia ciliata (Silva 305, US)
Miconia cinchonifolia (Amaral 522, US)
Miconia collatata (Cid Ferreira 4537, US)
Miconia comptifolia (Prance 4986, US)
Miconia cowanii (Austin 7265, US)
Miconia crassinervia (Pennington P22648, US)
Miconia cretacea (Campbell 8906, US)
Miconia cuspidata (Renner 560, US)
Miconia decurrens (Silveira 452, US)
Miconia diaphanea (Cid Ferreira 1417, US)
Miconia dichrophylla (Rodrigues 8054, INPA)
Miconia dispar (Cid Ferreira 4152, US)
Miconia dodecandra (Pipoly 6643, US)
Miconia dolichorrhyncha (Cid Ferreira 8680, US)
Miconia duckei (Lleras P17326, US)
Miconia egensis (Gentry 13033, US)
Miconia elata (Pires 14590, US)
Miconia elegans (Prance 9714, US)
Miconia emendata (Prance 8160, US)
Miconia eriocalyx (Daly 6932, US)
Miconia erioclada (Goldenberg 1294, UPCB)
Miconia eriodonta (Black 52, US)
Miconia eugeniooides (Alencar 462, US)
Miconia fallax (Vieira 655, US)
Miconia ferruginata (Fróes 30085, US)
Miconia fissa (Cid Ferreira 7155, US)
Miconia fosteri (Prance 12570, US)
Miconia fragrans (Silva 60642, US)

- Miconia grandifoliata* (Duarte 6899, US)
Miconia grattissima (Prance 20467, US)
Miconia heliotropoides (Ribeiro 1416, US)
Miconia holosericea (Prance 29492, US)
Miconia hypoleuca (Rodrigues 8603, US)
Miconia ibaguensis (Prance 7960, US)
Miconia impetiolaris (Prance 10252, US)
Miconia inaequalifolia (Krukoff 8936, US)
Miconia juruensis (Ule 5827, photo in US)
Miconia kavanayensis (Mori 17242, US)
Miconia lamprophylla (Prance 7318, NY)
Miconia lanata (Nelson 835, US)
Miconia lappacea (Silva 1993, US)
Miconia lasseri (Cavalcante 861, US)
Miconia lateriflora (Archer 8105, US)
Miconia lepidota (Milliken 659, US)
Miconia longifolia (Egler 46015, US)
Miconia longispicata (Ducke 164, US)
Miconia lourteigiana (Prance 2461, US)
Miconia lugonis (Goldenberg 961, UPCB)
Miconia macrothyrsa (Prance 9280, US)
Miconia macrotis (Kirkbride 2748, INPA)
Miconia manauara (Cid Ferreira 6723, INPA)
Miconia marginata (Rodrigues 166, US)
Miconia maroana (Nelson 1725, US)
Miconia matthaei (Prance 9752, US)
Miconia mattogrossensis (Nee 34714, US)
Miconia mazanana (Fróes 23901, US)
Miconia melinonis (Mexia 5961, US)
Miconia minutiflora (Nelson 1399, US)
Miconia mirabilis (Mori 17519, US)
Miconia multispicata (Thomas 4943, US)
Miconia myriantha (Silva 254, US)
Miconia nambyquarae (Cid Ferreira 4491, US)
Miconia navioensis (Miller 7716, US)
Miconia neblinensis (Maguire 60484, US)
Miconia nervosa (Prance 26540, US)
Miconia pachydonta (Anderson 12169, US)
Miconia paleacea (Goldenberg 954, UPCB)
Miconia phaeophylla (Mori 17678, US)
Miconia phanerostila (Prance 14249, US)
Miconia platypoda (Nee 42859, US)
Miconia poeppigii (Cid Ferreira 8088, US)
Miconia prancei (Prance 17973, US)
Miconia prasina (Pipoly 6882, US)
Miconia procumbens (Daly 10152, NY)
Miconia pseudocapsularis (Prance 21605, US)
Miconia pterocaule (Goldenberg 971, NY)
Miconia puberula (Prance 13814, US)
Miconia pubipetala (Ducke 1231, US)
Miconia punctata (Pipoly 6732, US)
Miconia pyrifolia (Rodrigues 8294, US)
Miconia racemosa (Sobel 4934, US)
Miconia radulaefolia (Cid Ferreira 8654, US)
Miconia regelii (Daly 7104, US)
Miconia rhytidophylla (Prance 3678, US)
Miconia rimachii (Prance 5748, US)
Miconia rubiginosa (Prance 29495, US)
Miconia rufescens (Cavalcante 810, US)
Miconia ruficalyx (Daly 4145, US)
Miconia rugosa (Prance 2245, US)
Miconia sagotiana (Pires 1486, US)
Miconia santaremensis (Ginzberger s.n., photo in US)
Miconia sarmentosa (Cowan 38536, US)
Miconia sastrei (De Granville 12436, US)
Miconia schunkei (Goldenberg 989, UPCB)
Miconia schwackei (Ducke 201, US)
Miconia secundiflora (Silva 2204, US)
Miconia serialis (Lobo 26, US)
Miconia serrulata (Steward P13253, US)
Miconia silicicola (Vicentini 1911, UPCB)
Miconia spichigeri (Ongley P21736, US)
Miconia splendens (Cid Ferreira 4153, US)
Miconia sprucei (Kawasaki 114, US)
Miconia stelligera (Prance 12385, US)
Miconia stellulata (Carreira 529, US)
Miconia stenostachya (Rabello 3748, US)
Miconia stephananthera (Ducke 1381, US)
Miconia strigosa (Prance 21685, US)
Miconia subandicola (Goldenberg 1387, UPCB)
Miconia subsimplex (Cid Ferreira 65, US)
Miconia subspicata (Michelangeli 1390, UPCB)
Miconia ternatifolia (Ramos 269, INPA)
Miconia tetragona (Cid Ferreira 8408, US)
Miconia tetrasperma (Prance 3508, US)
Miconia tetraspermoides (Lleras P19603, US)
Miconia tiliifolia (Prance 13782, US)
Miconia tillettii (Prance 10171, US)
Miconia tomentosa (Lowrie 633, US)
Miconia traillii (Rabello 3099, US)
Miconia trimera (Egler 46690, US)
Miconia trinervia (Irwin 48049, US)
Miconia triplinervis (Lima 528, US)
Miconia truncata (Cid Ferreira 3766, US)
Miconia umbrosa (Riedel 217, NY)
Miconia undata (Thomas 4922, US)
Miconia vilhenensis (Vieira 611, US)
Miconia wagneri (Prance 12362, US)
Miconia wittii (Fróes 27892, US)
Miconia woytkowskii (Teixeira 693, US)
- Brazil Central-West (68 spp)**
- Miconia acuminata* (Cordeiro 1168, US)
Miconia affinis (Berg P18522, US)
Miconia alata (Amaral 850, NY)
Miconia albicans (Irwin 7383, US)

Miconia alborufescens (Windisch 2160, US)
Miconia ampla (Philcox 3589, US)
Miconia apostachya (Harley 10702, US)
Miconia argyrophylla (Cid Ferreira 6230, US)
Miconia burchellii (Ratter 4337, US)
Miconia calvescens (Anderson 9607, US)
Miconia chamissois (Hatschbach 38727, US)
Miconia chartacea (Irwin 24915, US)
Miconia chrysophylla (Berg 18509, US)
Miconia ciliata (Amaral 839, US)
Miconia collatata (Prance 59642, US)
Miconia cubatanensis (Irwin 8956, US)
Miconia cuspidata (Ratter 1138, US)
Miconia cyathanthera (Ratter 4769, US)
Miconia dichrophylla (Silva 4748, INPA)
Miconia dispar (Cid Ferreira 6199, US)
Miconia dodecandra (Silva 277, US)
Miconia dolichorrhyncha (Hatschbach 37650, US)
Miconia egensis (Thomas 3940, US)
Miconia elegans (Thomas 3853, US)
Miconia fallax (Cid Ferreira 6102, US)
Miconia ferruginata (Irwin 24937, US)
Miconia gratissima (Harley 11170, US)
Miconia heliotropoides (Harley 10606, US)
Miconia hirtella (Irwin 13814, US)
Miconia holosericea (Dubbs 1284, US)
Miconia ibaguensis (Hatschbach 63856, US)
Miconia irwinii (Harley 11386, US)
Miconia lanata (Kirkbride 3496, US)
Miconia lepidota (Ratter 1378, US)
Miconia leucocarpa (Pott 9044, UPCB)
Miconia ligustroides (Maguire 44539, US)
Miconia macrothyrsa (Anderson 7292, US)
Miconia macrotis (Harley 10207, US)
Miconia matthaei (Harley 10650, US)
Miconia mattogrossensis (Philcox 3907, US)
Miconia minutiflora (Anderson 9349, US)
Miconia nambyquarae (Irwin 6566, US)
Miconia nervosa (Cordeiro 1169, US)
Miconia paucidens (Hatschbach 58671, UPCB)
Miconia pepericarpa (Irwin 19278, US)
Miconia prasina (Hatschbach 36094, US)
Miconia punctata (Harley 10812, US)
Miconia pyrifolia (Thomas 4181, US)
Miconia regelii (Thomas 3953, US)
Miconia rimachii (Thomas 3866, US)
Miconia rubiginosa (Irwin 18567, US)
Miconia rufescens (Cordeiro 1118, US)
Miconia sagotiana (Hatschbach 38186, US)
Miconia sclerophylla (Irwin 26633, US)
Miconia sellowiana (Silva 15, US)
Miconia serrulata (Harley 10740, NY)
Miconia spennerostachya (Malme 2228, US)

Miconia splendens (Cid Ferreira 6187, US)
Miconia staminea (Ratter 3280, US)
Miconia stenostachya (Hatschbach 38801, US)
Miconia tetraspermoides (Harley 11171, US)
Miconia theizans (Hatschbach 36359, US)
Miconia tiliifolia (Hatschbach 63827, US)
Miconia tomentosa (Santos 1680, US)
Miconia traillii (Hatschbach 62638, US)
Miconia vilhenensis (Goldenberg 419, US)
Miconia weddellii (Weddell 3001, photo in US)
Miconia woytkowskii (Ivanauskas 4317, UPCB)

Brazil Northeast (71 spp)

Miconia affinis (Daly 453, US)
Miconia alata (Fróes 11564, US)
Miconia albicans (Irwin 32564, US)
Miconia alborufescens (Harley 16848, US)
Miconia amacurensis (Lima 11, US)
Miconia amoena (Eupunino 311, US)
Miconia apostachya (Harley 10702, US)
Miconia buddlejoides (Goldenberg 389, CEPEC)
Miconia calvescens (Hage 1039, US)
Miconia carvalhoana (Melo et al. 1692, K)
Miconia caudigera (Mori 12806, US)
Miconia centrodesma (Mori 9297, US)
Miconia ceramicarpa (Baleé 773a, US)
Miconia chamissois (Fonseca 492, US)
Miconia chartacea (Harley 22569, US)
Miconia ciliata (Carvalho 985, US)
Miconia cinnamomifolia (Hatschbach 48781, US)
Miconia compressa (Amorim 880, US)
Miconia cubatanensis (Carvalho 3753, US)
Miconia cuspidata (Araujo 74, UPCB)
Miconia cyathanthera (Harley 26380, US)
Miconia dodecandra (Belém 910, US)
Miconia dorsaliporosa (Paixão 851, CEPEC)
Miconia elegans (Hatschbach 53408, US)
Miconia fallax (Ribeiro 44, RB)
Miconia fasciculata (Goldenberg 870, CEPEC)
Miconia ferruginata (Harley 21193, US)
Miconia francavillana (Mori 13127, US)
Miconia herpetica (Fonseca 472, US)
Miconia hirtella (Noblick 371, US)
Miconia holosericea (Taylor E1272, US)
Miconia hypoleuca (Noblick 2384, US)
Miconia ibaguensis (Harley 21276, US)
Miconia irwinii (Giulietti CFCR 1310, US)
Miconia johnwurdackiana (Hatschbach 47984, CAS)
Miconia jucunda (Jesus 576, US)
Miconia kappleri (Baleé 105, US)
Miconia latecrenata (Santos 975, US)
Miconia lateriflora (Jangoux 418, US)
Miconia lepidota (Lima 5, US)

- Miconia leucocarpa* (Magalhães 160, UPCB)
Miconia ligustroides (Harley 21198, US)
Miconia longifolia (Jangoux 866, US)
Miconia lurida (Eupunino 316, US)
Miconia macrothyrsa (Rodrigues 133, UPCB)
Miconia minutiflora (Pinto 102/86, US)
Miconia mirabilis (Eupunino 473, US)
Miconia mourae (Amorim 5907, NY)
Miconia nervosa (Harley 19476, US)
Miconia octopetala (Santos 1073, US)
Miconia paniculata (Santos 176, US)
Miconia pepericarpa (Harley 15457, US)
Miconia poeppigii (Krukoff 1973, US)
Miconia prasina (Moraes 1043, US)
Miconia pyrifolia (Mori 10929, US)
Miconia racemosa (Ribeiro 1208, US)
Miconia rimalis (Harley 25819, US)
Miconia rubiginosa (Thomas 9666, US)
Miconia ruficalyx (Santos 840, US)
Miconia rufipila (Blanchet 3952, photo in US)
Miconia sclerophylla (Hatschbach 47413, US)
Miconia serialis (Eugênio 928, US)
Miconia serrulata (Santos 185, US)
Miconia speciosa (Amorim 4914, CEPEC)
Miconia splendens (Silva 702, US)
Miconia stenostachya (Belém 700, US)
Miconia subcordata (Blanchet 1011, photo in US)
Miconia theizans (Carvalho 1719, US)
Miconia tomentosa (Daly D210, US)
Miconia trianae (Glaziou 16931, BR)
Miconia tristis (Santos 898, US)

Brazil Southeast (129 spp)

- Miconia affinis* (Araújo s.n., US)
Miconia albicans (Trinta 1059, US)
Miconia alborufescens (Pabst 7113, US)
Miconia altissima (Glaziou 16969, US)
Miconia amoena (Weinberg 604, US)
Miconia angelana (Romero & Nakajima 3773, HUFU)
Miconia argyraea (Glaziou 16905, R)
Miconia augusti (Glaziou 16895, R)
Miconia baumgratziana (Caddah 730, UPCB)
Miconia brasiliensis (Ribeiro 865, US)
Miconia brunnea (Goldenberg 455, UPCB)
Miconia buddlejoides (Hatschbach 49931, CEPEC)
Miconia burchellii (Irwin 22956, US)
Miconia cabucu (Hoehne 28306, US)
Miconia calvescens (Gomes 1147, US)
Miconia capixaba (Thomaz 819, UEC)
Miconia castaneiflora (Caddah 447, UPCB)
Miconia caudigera (Arbo 4499, US)
Miconia centrodesma (Shepherd 10973, UEC)
Miconia chamissois (Eiten 1919, US)
- Miconia chartacea* (Renner 2113, US)
Miconia ciliata (Ribeiro 36, UPCB)
Miconia cinerascens (Emmerich 1616, US)
Miconia cinerea (Glaziou 11974, R)
Miconia cinnamomifolia (Felippe 41, US)
Miconia cipoensis (Pirani CFSC 12699, US)
Miconia collatata (Ratter 8761, UEC)
Miconia corallina (Irwin 29464, US)
Miconia cubatanensis (Romero 2071, US)
Miconia cyathanthera (Nakajima 1086, US)
Miconia dichroa (Glaziou 18223, K)
Miconia discolor (Brade 17134, US)
Miconia divaricata (Gardner 391, US)
Miconia dodecandra (Araújo 6162, US)
Miconia dorsaliporosa (Goldenberg 915, UPCB)
Miconia dura (Sello s.n., cited by Triana 1871)
Miconia eichleri (Glaziou 8679, US)
Miconia elaeodendron (Baumgratz 797, RB)
Miconia elegans (Romero 1146, US)
Miconia fallax (Harley 24955, US)
Miconia fasciculata (Gardner 183, US)
Miconia ferruginata (Eiten 3223, US)
Miconia flammea (Casaretto 2710, TO)
Miconia fluminensis (Ule 4229, photo in US)
Miconia formosa (Glaziou 17061, R)
Miconia francavillana (Pirani 3470, US)
Miconia gigantea (Herb. Urban 17548, P)
Miconia gilva (Glaziou 16897, R)
Miconia glazioviana (Glaziou 16943, US)
Miconia herpetica (Glaziou 13836, US)
Miconia hirtella (Romero 547, US)
Miconia holosericea (Fontella 95, US)
Miconia hyemalis (Hoehne s.n., SPF 13806)
Miconia hypoleuca (Sobral 4747, US)
Miconia ibaguensis (Nakajima 1404, US)
Miconia inaequidens (Herb. Deless. s.n., G-DEL)
Miconia inconspicua (Brade 109462, US)
Miconia irwinii (Hatschbach 31533, US)
Miconia jucunda (Brade 6073, US)
Miconia klotzschii (Sello B1126, US)
Miconia kollmannii (Kollmann 5064, MBML)
Miconia kriegeriana (Chiavegatto 87, RB)
Miconia labiakiana (Kollmann 7438, UPCB)
Miconia laeta (Glaziou s.n., US)
Miconia latecrenata (Davis 60572, UEC)
Miconia lepidota (Hatschbach 46290, US)
Miconia leucocarpa ((Eiten 3271, US)
Miconia ligustroides (Irwin 19578, US)
Miconia longicuspis (Lima VII.990, US)
Miconia lurida (Claussen 48, US)
Miconia lymanii (Maguire 44559, US)
Miconia macrothyrsa (Irwin 2342, US)
Miconia maximowicziana (Glaziou 17543, R)

- Miconia mellina* (St. Hilaire D62, US)
Miconia mendoncae (Mexia 5244, US)
Miconia michelangeliana (Kollmann 10743, MBML)
Miconia minutiflora (Schiavini 105, US)
Miconia mirabilis (Martinelli 9717, US)
Miconia molesta (Glaziou 17542, US)
Miconia mourae (Moura 132, P)
Miconia multinervia (Langsdorff s.n., P)
Miconia nervosa (Martinelli 9759, US)
Miconia oblongifolia (Glaziou 13437, R)
Miconia octopetala (Hatschbach 49984, US)
Miconia organensis (Glaziou 17946, US)
Miconia ovalifolia (Glaziou 16908, US)
Miconia ovata (Cited by Cogniaux 1891)
Miconia paniculata (Pabst 5208, US)
Miconia paradoxa (Hatschbach 28986, US)
Miconia paucidens (Goldenberg 28482, UEC)
Miconia penduliflora (Goldenberg 1536, UPCB)
Miconia pennipilis (Sello 1060, US)
Miconia pepericarpa (Eiten 3389, US)
Miconia petroniana (Glaziou 9463, US)
Miconia petropolitana (Glaziou 9463, US)
Miconia picinguabensis (Pedroni 628, UEC)
Miconia polyandra (Hoehne 23835, US)
Miconia prasina (Araújo 1542, US)
Miconia pseudoeichleri (Glaziou 16935, US)
Miconia pulchra (Glaziou 17541, P)
Miconia pusilliflora (Kuhlmann 659, US)
Miconia pyrifolia (Botta 2934, UEC)
Miconia rabenii (Raben s.n., photo in US)
Miconia racemifera (Brade 16770, US)
Miconia rimalis (Hatschbach 29942, US)
Miconia robusta (Silveira RB 111547, US)
Miconia rubiginosa (Romero 522, US)
Miconia rufipila (Lombardi 5376, UPCB)
Miconia ruschiana (Kollmann 3424, UPCB)
Miconia saldanhae (Peixoto 3500, US)
Miconia sclerophylla (Maguire 44766, US)
Miconia sellowiana (Handro 77285, US)
Miconia serrulata (Kirizawa 1673, SP)
Miconia setosociliata (Glaziou 16934, US)
Miconia shepherdii (Goldenberg 71, UEC)
Miconia speciosa (St. Hilaire 340, P)
Miconia splendens (Folli 324, US)
Miconia staminea (Eiten 7838, US)
Miconia stenostachya (Araújo 272, US)
Miconia subvernica (Glaziou 590, R)
Miconia tentaculifera (Arbo 3946, US)
Miconia theizans (Romero 1939, US)
Miconia trianae (Williams 1456, US)
Miconia tristis (Silva 139, UEC)
Miconia urophylla (Mimura 564, US)
Miconia valtheri (Goldenberg 76, UPCB)
- Miconia vismioides* (Collector unknown, BM)
Miconia warmingiana (Irwin 30284, US)
Miconia willdenowii (Lemos 7529, US)
- Brazil South (40 spp)**
- Miconia albicans* (Hatschbach 27101, US)
Miconia brasiliensis (Klein 107, US)
Miconia brunnea (Goldenberg 736, UPCB)
Miconia buddlejoides (Hatschbach 19920, MBM)
Miconia cabucu (Klein 9677, US)
Miconia calvescens (Reitz 18102, US)
Miconia chamissois (Hatschbach 13354, US)
Miconia chartacea (Reitz 3186, US)
Miconia cinerascens (Reitz 9384, US)
Miconia cinnamomifolia (Klein 1013, US)
Miconia collatata (Hatschbach 14294, US)
Miconia cubatanensis (Klein 8289, US)
Miconia discolor (Reitz 2108, US)
Miconia dodecandra (Hatschbach 14396, US)
Miconia eichleri (Hatschbach 31030, US)
Miconia fasciculata (Reitz 3213, US)
Miconia flammea (Smith 7991, US)
Miconia hyemalis (Reitz 4749, US)
Miconia ibaguensis (Hatschbach 40200, US)
Miconia inconspicua (Reitz 5213, US)
Miconia jucunda (Klein 876, US)
Miconia lagunensis (Reitz 61, US)
Miconia latecrenata (Smith 5663, US)
Miconia ligustroides (Hatschbach 17628, US)
Miconia longicuspis (Camargo 101, UPCB)
Miconia lymanii (Smith 7878, US)
Miconia paniculata (Klein 2339, US)
Miconia paucidens (Hatschbach 39284, MBM)
Miconia petropolitana (Ferreira, G.F. 37, US)
Miconia prasina (Hatschbach 19135, US)
Miconia pusilliflora (Klein 1309, US)
Miconia racemifera (Meyer 362, UPCB)
Miconia ramboi (Wasum 2118, US)
Miconia sellowiana (Wasum 8689, US)
Miconia splendens (Lindeman 1676, US)
Miconia stenostachya (Hatschbach 20029, US)
Miconia theizans (Hatschbach 9568, US)
Miconia tristis (Reitz 6523, US)
Miconia valtheri (Goldenberg 718, UPCB)
Miconia willdenowii (Goldenberg 808, UPCB)
- Paraguay (17 spp)**
- Miconia albicans* (Hahn 1744, US)
Miconia amambayensis (Hassler 11381, photo in US)
Miconia calvescens (Zardini 12696, US)
Miconia chamissois (Zardini 28015, US)
Miconia cinerascens (Hahn 2135, US)
Miconia collatata (Hassler 10566, NY)

Miconia discolor (Brunner 896, US)
Miconia fallax (Cited by Cogniaux 1891)
Miconia ibaguensis (Schinini 29379, US)
Miconia paraguayensis (Hassler 26127, photo in US)
Miconia paucidens (Schinini 20229, US)
Miconia prasina (Cited by Wurdack et al. 1993)
Miconia pusilliflora (Zardini 32517, US)
Miconia staminea (Zardini 16863, US)
Miconia stenostachya (Marin 792, G)
Miconia theizans (Schinini 9139, US)
Miconia tristis (Marin 793, G)

Argentina (10 spp)

Miconia calvescens (Krapovickas 26666, US)
Miconia chamissois (Pedersen 5429, US)
Miconia cinerascens (Krapovickas 23370, US)
Miconia collatata (Curran 27, US)
Miconia discolor (Vanni 3066, US)
Miconia ioneura (Cabrera 10581, US)
Miconia molybdea (Giusti 12110, US)

Miconia petropolitana (Legname 5556, US)
Miconia pusilliflora (Vanni 826, US)
Miconia staminea (Krapovickas 23776, US)

Uruguay (1 sp)

Miconia hyemalis (Cited by Wurdack 1962)

Sri Lanka (1 sp)

Miconia calvescens (Fosberg 58147, NY; invasive)

Hawaiian Is. (1 sp)

Miconia calvescens (Staples 770, NY; invasive)

Society Is. (1 sp)

Miconia calvescens (Cited by Meyer 1996; invasive)

Australia (1 sp)

Miconia calvescens (Cited by Csurhes 2008; invasive)

LIST 4. *Miconia* names synonymized under other genera.

Miconia africana Jacq.-Fél., Bull. Mus. Hist. Nat. Paris 10: 642 (1938).
= *Leandra quinquedentata* (DC.) Cogn.

Miconia agrestis (Aubl.) Baill., Hist. Pl. 7: 35 (1880).
Basionym: *Melastoma agrestis* Aubl.
= *Leandra agrestis* (Aubl.) Raddi.

Miconia angulata (Griseb.) M.Gómez, Anales Hist. Nat. 23: 68 (1894).
= *Calycogonium angulatum* Griseb.

Miconia angustiflora Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 246 (1850).
= *Tetrazygia bicolor* (Mill.) Cogn.

Miconia angustifolia (Sw.) Griseb., Goett. Abh. 7: 69 (1857).
Basionym: *Melastoma angustifolia* Sw.
= *Tetrazygia angustifolia* (Sw.) DC.

Miconia atrata (Spring) Wawra, Oesterr. Bot. Z. 13: 89 (1883).
= *Clidemia atrata* Spring.

Miconia aschenborniana S.Schauer ex Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 246 (1850).
= *Leandra cornoides* (Schltdl. & Cham.) Cogn.

Miconia baillonii M.Gómez, Anales Hist. Nat. 23: 69 (1894).
= *Conostegia clidemiooides* C.Wright ex Griseb.

Miconia baracoana M.Gómez, Anales Hist. Nat. 23: 68 (1894).
= *Calycogonium clidemiooides* Griseb.

Miconia barbicaulis Gleason, Bull. Torrey Bot. Club 68: 248 (1941).
= *Tococa symphyandra* (Triana) Cogn.

Miconia bergii Beurl., Kongl. Vetensk. Acad. Handl. 40: 129 (1854).
= *Conostegia xalapensis* (Bonpl.) D.Don ex DC.

Miconia blancheana Urb., Repert. Spec. Nov. Regni Veg. 17: 405 (1921).
= *Pachyanthus blancheanus* (Urb.) Urb. & Ekman..

Miconia brachycentra (Griseb.) M.Gómez, Anales Hist. Nat. 23: 69 (1894).
Basionym: *Graffenrieda brachycentra* Griseb.
= *Tetrazygia brachycentra* (Griseb.) C. Wright.

Miconia calcarata Gleason, Bull. Torrey Bot. Club 72 (5): 474 (1945).
= *Tessmannianthus calcaratus* (Gleason) Wurdack.

Miconia capillaris (Sw.) M.Gómez, Anales Hist. Nat. 23: 68 (1894).
Basionym: *Melastoma capillaris* Sw.
= *Sagraea capillaris* (Sw.) DC.

Miconia caryophyllaea (DC.) Triana, Trans. Linn. Soc. London 28 (1): 106 (1871).
Basionym: *Truncaria caryophyllaea* DC.
= *Tococa caryophyllaea* (DC.) S.S.Renner.

Miconia catalpifolia Kraenzl., Vierteljahrsschr. Naturf. Ges. Zürich 76: 153 (1931).
= *Graffenrieda latifolia* (Naudin) Triana.

Miconia christophoriana (Ham.) DC., Prodr. 3: 200 (1828).
Basionym: *Melastoma christophorianum* Ham.
= *Tetrazygia discolor* (L.) DC.

Miconia chrysandra Griseb., Cat. Pl. Cub. 99 (1866).
= *Graffenrieda chrysandra* (Griseb.) Triana.

Miconia cinnamomea Beurl., Kongl. Vetensk. Acad. Handl. 40: 131 (1854).
= *Conostegia cinnamomea* (Beurl.) Wurdack.

Miconia coccinea (Rich.) Judd & Skean, Bull. Florida State Mus., Biol. Sci. 36: 61 (1991).
Basionym: *Melastoma coccineum* Rich.
= *Charianthus alpinus* (Sw.) R.A.Howard.

Miconia consimilis Pilg., Verh. Bot. Vereins Prov. Brandenburg 47: 170 (1905).
= *Graffenrieda gracilis* (Triana) L.O.Williams.

Miconia corymbosa (Rich.) Judd & Skean, Bull. Florida State Mus., Biol. Sci. 36: 61 (1991).
Basionym: *Melastoma corymbosum* Rich.
= *Charianthus corymbosus* (Rich.) Cogn.

Miconia cuabae Urb., Repert. Spec. Nov. Regni Veg. 19: 305 (1924).
= *Henriettea cuabae* (Urb.) Borhidi.

Miconia emarginata Ruiz & Pav., Syst. Veg. Fl. Peruv. Chil. 1: 106 (1798).
= *Graffenrieda emarginata* (Ruiz & Pav.) Triana.

Miconia erioneura Cogn., Mem. Torrey Bot. Club 6: 38 (1896).
= *Tococa erioneura* (Cogn.) Wurdack.

Miconia erythrophylla Ule, Notizbl. Königl. Bot. Gart. Berlin 6: 357 (1915).
= *Tococa erythrophylla* (Ule) Wurdack.

Miconia fadyenii (Hook.) Judd & Skean, Bull. Florida State Mus., Biol. Sci. 36 (2): 61 (1991).
= *Tetrazygia fadyenii* Hook.

Miconia glandulosa (Sw.) Naudin, Ann. Sci. Nat., Bot., Ser. 3 16: 244 (1850).
Basionym: *Melastoma glandulosa* Sw.
= *Tetrazygia hispida* (Sw.) Macfad.

Miconia heterophylla (Naudin) M.Gómez, Anales Hist. Nat. 23: 68 (1894).
= *Calycogonium heterophyllum* Naudin.

Miconia incarum J.F.Macbr., Trop. Woods 17: 12 (1929).
= *Meriania radula* (Benth.) Triana.

Miconia leblondii Judd & Skean, Bull. Florida State Mus., Biol. Sci. 36: 61 (1991).
= *Charianthus nodosus* (Desr.) Triana.

Miconia lima (Desr.) M.Gómez, Anales Hist. Nat. 23: 69 (1894).
Basionym: *Melastoma lima* Desr.
= *Leandra lima* (Desr.) Judd & Skean.

Miconia lindeniana (Naudin) M.Gómez, Anales Hist. Nat. 23: 68 (1894).

= ***Calycogonium lindenianum*** Naudin.

Miconia lundelliana L.O.Williams, Fieldiana, Bot. 29 (10): 573 (1963).

= ***Pachyanthus lundellianus*** (L.O.Williams) Judd & Skean.

Miconia microphylla (C.Wright) M.Gómez, Anales Hist. Nat. 23: 69 (1894).

= ***Calycogonium microphyllum*** C.Wright.

Miconia monocephala Urb., Symb. Antill. 9: 117 (1923).

= ***Pachyanthus monocephalus*** (Urb.) Borhidi.

Miconia neomicrantha Judd & Skean, Bull. Florida State Mus., Biol. Sci. 36 (2): 62 (1991).

= ***Ossaea micrantha*** (Sw.) Macfad. ex Cogn.

Miconia nitens Benth., J. Bot. 2: 313 (1840).

= ***Tococa nitens*** (Benth.) Triana.

Miconia oleifolia (Griseb.) M.Gómez, Anales Hist. Nat. 23: 69 (1894).

= ***Pachyanthus oleifolius*** Griseb.

Miconia ossaeiformis Naudin, Ann. Sci. Nat., Bot., Ser. 3 17: 212 (1850).

= ***Mecranium purpurascens*** (DC.) Triana.

Miconia ottoschulzii Urb. & Ekman, Ark. Bot. 22A (17): 39 (1929).

= ***Graffenrieda ottoschulzii*** (Urb. & Ekman) Urb. & Ekman.

Miconia pallens (Spreng.) Triana, Trans. Linn. Soc. London 28 (1): 103 (1871).

Basionym: *Melastoma pallens* Spreng.

= ***Tetrazygia pallens*** (Spreng.) Cogn.

Miconia penninervis (Griseb.) M.Gómez, Anales Hist. Nat. 23: 68 (1894).

= ***Clidemia penninervis*** Griseb.

Miconia platyphylla (Benth.) L.O.Williams, Fieldiana, Bot. 29 (10): 575 (1963).

= ***Tococa platyphylla*** Benth.

Miconia plicata (Griseb.) M.Gómez, Anales Hist. Nat. 23: 69 (1894).

= ***Calycogonium plicatum*** Griseb.

Miconia poiretii (Griseb.) M.Gómez, Anales Hist. Nat. 23: 69 (1894).

= ***Pachyanthus poiretii*** Griseb.

Miconia pterosepala Urb., Symb. Antill. 9: 116 (1923).

= ***Clidemia pterosepala*** (Urb.) Alain.

Miconia purpurascens (Aubl.) DC., Prodr. 3: 179 (1828).

Basionym: *Melastoma purpurascens* Aubl.

= ***Aciotis purpurascens*** (Aubl.) Triana.

Miconia purpurea (D.Don) Judd & Skean, Bull. Florida State Mus., Biol. Sci. 36: 62 (1991).

= ***Charianthus purpureus*** D.Don.

Miconia rangeliana Wright ex Griseb., Cat. Pl. Cub. 99 (1866).

= ***Tetrazygia laxiflora*** Naudin.

Miconia rosmarinifolia (Griseb.) M.Gómez, Anales Hist. Nat. 23: 68 (1894).
= ***Calycogonium rosmarinifolium*** Griseb.

Miconia rostrata (Bertol.) Triana, Trans. Linn. Soc. London 28 (1): 131 (1871).
Basionym: *Melastoma rostratum* Bertol.
= ***Conostegia xalapensis*** (Bonpl.) D.Don ex DC.

Miconia rupicola Gleason, Bull. Torrey Bot. Club 52: 383 (1925).
= ***Conostegia polyandra*** Benth.

Miconia scandens Ruiz & Pav., Fl. Peruv. 4: 395 (1802).
= ***Adelobotrys adscendens*** (Sw.) Triana.

Miconia secundo-angustifolia M.Gómez, Anales Hist. Nat. 23: 68 (1894).
= ***Pachyanthus angustifolius*** Griseb.

Miconia secundo-lanceolata M.Gómez, Anales Hist. Nat. 23: 69 (1894).
= ***Calycogonium lanceolatum*** Griseb.

Miconia setimarginata Pittier, Bol. Soc. Venez. Ci. Nat. 11: 27 (1947).
= ***Tococa oligantha*** Gleason.

Miconia stahlii (Cogn.) M.Gómez, Dicc. Bot. Nom. Vulg. Cub. Puerto-Riquenos 37 (1889).
= ***Tetrazygia stahlii*** Cogn.

Miconia stenopetala Griseb., Fl. Brit. W. Indian Is. 711 (1864).
= ***Platycentrum clidemoides*** Naudin.

Miconia stylosa Cogn. in Mart., Fl. bras. 14 (4): 257 (1887).
= ***Tococa subciliata*** (DC.) Triana.

Miconia subciliata DC., Prodr. 3: 187 (1828).
= ***Tococa subciliata*** (DC.) Triana.

Miconia subcrustulata Beurl., Kongl. Vetensk. Acad. Handl. 40: 130 (1854).
= ***Conostegia subcrustulata*** (Beurl.) Triana.

Miconia subhirsuta (DC.) M. Gomez, Anales Hist. Nat. 23: 69 (1894).
Basionym: *Conostegia subhirsuta* DC.
= ***Conostegia icosandra*** (Sw. ex Wikstr.) Urb.

Miconia symphyandra Triana, Trans. Linn. Soc. London 28 (1): 126 (1871).
= ***Tococa symphyandra*** Cogn.

Miconia umbilicata (Bertol.) Triana, Trans. Linn. Soc. London 28 (1): 131 (1871).
Basionym: *Melastoma umbilicatum* Bertol.
= ***Conostegia xalapensis*** (Bonpl.) D.Don ex DC.

Miconia wrightii (Griseb.) Triana, Trans. Linn. Soc. London 28 (1): 103 (1871).
= ***Pachyanthus wrightii*** Griseb.

Miconia xalapensis (Bonpl.) M. Gomez, Anales Hist. Nat. 23: 69 (1894).
Basionym: *Melastoma xalapense* Bonpl.
= ***Conostegia xalapensis*** (Bonpl.) D.Don ex DC.