



Two new combinations in the genus *Bornmuellera* (Brassicaceae)

IVANA REŠETNIK¹, GERALD M. SCHNEEWEISS² & ZLATKO LIBER¹

¹ University of Zagreb, Faculty of Science, Department of Botany, Marulićev trg 9a, HR-10000 Zagreb, Croatia.

E-mail: ivana.resetnik@biol.pmf.hr

² University of Vienna, Department of Systematic and Evolutionary Botany, Rennweg 14, A-1030 Vienna, Austria

The genus *Bornmuellera* Haussknecht (1897: 71) encompasses seven species distributed from the south-western Balkan Peninsula to Anatolia (Warwick *et al.* 2006). The original description of the genus encompassed only one species, *B. tymphaea* (Hausskn.) Haussknecht (1897: 72) which he previously described within *Vesicaria* Adanson (1763: 420). Haussknecht characterized the genus with non-saccate deciduous sepals, white flowers with obtuse petals, filaments with appendages at the base, sessile subglobose glabrous fruits and two seeds in each loculi. The other taxa known at the time, *B. cappadocica* (Willdenow 1800: 452) Cullen & T.R. Dudley in Dudley & Cullen (1965: 228), *B. glabrescens* (Boiss. & Balansa in Boissier 1856: 32) Cullen & T.R. Dudley in Dudley & Cullen (1965: 228) and *B. baldaccii* (Degen 1896: 413) Heywood (1964: 61), were left classified in either *Ptilotrichum* Meyer (1831: 64, e.g., Degen 1896, Boissier 1867) or *Vesicaria* (Boissier 1856). Later, Heywood (1964) and subsequently Dudley & Cullen (1965) taxonomically re-defined the genus, whose new circumscription was followed by succeeding authors (Hartvig 2002, Warwick *et al.* 2006).

Recent molecular phylogenetic analyses of the tribe Alysseae (Rešetnik *et al.* 2013) revealed that the monotypic genera *Leptoplax* Schulz (1933: 92) and *Physocardamum* Hedge (1968: 293) phylogenetically nest within *Bornmuellera* and thus should be included in it. *Leptoplax emarginata* (Boissier 1842: 160) Schulz (1933: 92), endemic to Greece, was previously considered closely related to and even included in *Peltaria* Jacquin (1762: 260) of tribe Thlaspidaceae (Dudley & Cullen 1965, Ball 1993, Appel & Al-Shehbaz 2003, Warwick *et al.* 2008) because of shared compressed indehiscent silicules compared to terete, thin leathery fruits in *Bornmuellera*. As is often the case in Brassicaceae (Koch *et al.* 2003 and references therein, Alexander *et al.* 2010, Moazzeni *et al.* 2010), the similarity in fruit characters is superficial and the compressed siliculæ in *Peltaria* and *Leptoplax* are of independent origin. Diagnostic morphological characters shared between *Leptoplax* and *Bornmuellera* include malpighiaceous hairs, non-saccate sepals, white flowers with obtuse petals and filaments with appendages. Both genera share a base chromosome number of $x=8$ (Constantinidis *et al.* 2002; compared to $x=7$ in *Peltaria*: Warwick & Al-Shehbaz 2006) and an ecological specialization to serpentine soils and the capability of nickel hyperaccumulation. Furthermore, *L. emarginata* hybridises easily and extensively with sympatric *Bornmuellera* species, *B. tymphaea* and *B. baldaccii* (Ball 1993, Hartvig 1986, 2002). A molecular phylogenetic study based on nuclear ITS data (Cecchi *et al.* 2010) inferred *Leptoplax* as sister to the European members of *Bornmuellera*. When considering Anatolian *Bornmuellera* taxa, *L. emarginata* is inferred by both nuclear and plastid sequence data as embedded within *Bornmuellera* (Rešetnik *et al.* 2013).

Previously, *Physocardamum* was tentatively placed in Lepidieae (Hedge 1968, Al-Shehbaz 1986) or not assigned to any tribe (Al-Shehbaz *et al.* 2006). The tribal placement of *Physocardamum* in Alysseae was only recently clarified by molecular data (Warwick *et al.* 2010, Rešetnik *et al.* 2013). This is supported by morphological data, because most features of *P. davisii* Hedge (1968: 293) are shared with *Bornmuellera*, including perennial life form, presence of malpighiaceous trichomes, non-saccate sepals, white petals, inflated and glabrous siliculæ, and biovulate loculi; the only difference is that *Physocardamum* possesses edentate filaments (Hedge 1968). Molecular data identified *P. davisii* as sister to Anatolian *B. cappadocica* and resolved this Turkish clade as sister to a clade including the European species, i.e., including *Leptoplax* (Rešetnik *et al.* 2013).

Therefore, in the light of available molecular data (Rešetnik *et al.* 2013) and a re-evaluation of morphological characters, the monotypic genera *Leptoplax* and *Physocardamum* should be merged with *Bornmuellera*, requiring the following new combinations.

New combinations in *Bornmuellera* Hausskn.

Bornmuellera emarginata (Boiss.) Rešetnik, *comb. nov.*

basionym: *Ptilotrichum emarginatum* Boissier (1842: 160), synonyms: *Koniga emarginata* (Boiss.) Nyman (1855: 200); *Peltaria emarginata* (Boiss.) Haussknecht (1893: 111); *Leptoplax emarginata* (Boiss.) O.E. Schulz (1933: 92)
Type: [Greece, W Aegean] “Hab. in monte Delphi Euboea”, ?1837, Aucher-Éloy no. 228 (G-BOISS photo!)

Bornmuellera davisii (Hedge) Rešetnik, *comb. nov.*

basionym: *Physocardamum davisii* Hedge (1968: 293)

Type: “Turkey, B9 Ağrı: 2 km SW of Hamur (Murat valley). 1670 m. Sloping meadows; perennial; fruits inflated; pale green; flowers white.” 2 June 1966, Davis 44017 (<http://data.rbge.org.uk/herb/E00386088>) (E photo!)

References

- Adanson, M. (1763) *Families des Plantes* 2. Vincent, Paris. 640 pp.
- Alexander, P.J., Windham, M.D., Govindarajulu, R., Al-Shehbaz, I.A. & Bailey, C.D. (2010) Molecular phylogenetics and taxonomy of the genus *Thysanocarpus* (Brassicaceae). *Systematic Botany* 35: 559–577.
<http://dx.doi.org/10.1600/036364410792495926>
- Al-Shehbaz, I.A. (1986) The genera of *Lepidieae* (Cruciferae; Brassicaceae) in the southeastern United States. *Journal of the Arnold Arboretum* 67: 265–311.
- Al-Shehbaz, I.A., Beilstein, M.A. & Kellogg, E.A. (2006) Systematics and phylogeny of the Brassicaceae (Cruciferae): an overview. *Plant Systematics and Evolution* 259: 89–120.
<http://dx.doi.org/10.1007/s00606-006-0415-z>
- Appel, O. & Al-Shehbaz, I.A. (2003) Cruciferae. In: Kubitzki, K. & Bayer, C. (Eds.) *The families and Genera of Vascular Plants V. Flowering Plants, Dicotyledons: Malvales, Capparales, and non-betalain Caryophyllales*. Springer, Berlin, pp. 75–174.
http://dx.doi.org/10.1007/978-3-662-07255-4_17
- Ball, P.W. (1993) *Peltaria* Jacq. In: Tutin, T.G., Burges, N.A., Chater, A.O., Edmondson, J.R., Heywood, V.H., Moore, D.M., Valentine, D.H., Walters, S.M. & Webb, D.A. (Eds.) *Flora Europaea* 1, 2nd ed. Cambridge University Press, Cambridge, pp. 358.
- Boissier, P.E. (1842) *Plantae aucherianae orientalis enumeratae, cum novarum specierum descriptione*. *Annales des Sciences Naturelles Botanique* II 17: 150–162.
- Boissier, P.E. (1856) *Diagnoses Plantarum Orientalium Novarum*. Series secunda, no. 5. B. Herrmann, Lipsia. 118 pp.
- Boissier, P.E. (1867) *Flora Orientalis* 1. H. Georg, Basel, Switzerland. 1017 pp.
- Cecchi, L., Gabrielli, R., Arnetoli, M., Gonnelli, C., Hasko, A. & Selvi, F. (2010) Evolutionary lineages of nickel hyperaccumulation and systematics in European *Alysseae* (Brassicaceae): evidence from nrDNA sequence data. *Annals of Botany* 106: 751–767.
- Constantinidis, T., Bareka, E-P. & Kamari, G. (2002) Karyotaxonomy of Greek serpentine angiosperms. *Botanical Journal of the Linnean Society* 139: 109–124.
<http://dx.doi.org/10.1046/j.1095-8339.2002.00044.x>
- Degen, A. (1896) Bemerkungen über einige orientalische Pflanzenarten. XXII–XXVII. *Österreichische Botanische Zeitschrift* 46: 411–418.
<http://dx.doi.org/10.1007/bf01794719>
- Dudley, T.R. & Cullen, J. (1965) Studies in the Old World *Alysseae* Hayek. *Feddes Repertorium* 71: 218–228.
<http://dx.doi.org/10.1002/fedr.19650710105>
- Hartvig, P. (1986) *Leptoplax* O.E. Schulz. In: Strid, A. (Ed.) *Mountain Flora of Greece* 1. Cambridge University Press, Cambridge, pp. 275–276.
- Hartvig, P. (2002) *Bornmuellera* Hausskn. In: Strid, A. & Tan, K. (Eds.) *Flora Hellenica* 2. A. R. G. Gantner, Ruggell, pp. 233–234.
- Haussknecht, C. (1893) Symbolae ad floram graecam. Aufzählung der im Sommer 1885 in Griechenland gesammelten Pflanzen. *Mitteilungen des Thüringischen Botanischen Vereins* 3–4: 96–116.
- Haussknecht, C. (1897) Drei neue Cruciferen-Gattungen der orientalischen Flora. *Mitteilungen des Thüringischen Botanischen Vereins* 11: 68–76.
- Hedge, I.C. (1968) *Physocardamum*: a new genus of Cruciferae from Turkey. *Notes from the Royal Botanic Garden, Edinburgh* 28: 293–296.
- Heywood, V.H. (1964) *Bornmuellera* Hausskn. In: Tutin, T.G., Heywood, V.H., Burges, N.A., Valentine D.H., Walters, S.M. & Webb, D.A. (eds.) *Flora Europaea* 1. Cambridge University Press, Cambridge, pp. 306.
- Jacquín, N. J. (1762) *Enumeratio stirpium plerarumque, quae sponte crescunt in agro Vindobonensi, montibusque confinibus*. Joannis Pauli Kraus, Vindobona, 315 pp.

- Koch, M.A., Al-Shehbaz, I.A. & Mummenhoff, K. (2003) Molecular systematics, evolution, and population biology in the mustard family (Brassicaceae). *Annals of the Missouri Botanical Garden* 90: 151–171.
<http://dx.doi.org/10.2307/3298580>
- Meyer, C.A. (1831) *Ptilotrichum* C.A.Mey. In: Ledebour, C.F. (ed.) *Flora Altaica* 3. Berolini, pp. 64–67.
- Moazzeni, H., Zarre, S., Al-Shehbaz, I.A. & Mummenhoff, K. (2010) Phylogeny of *Isatis* (Brassicaceae) and allied genera based on ITS sequences of nuclear ribosomal DNA and morphological characters. *Flora* 205: 337–343.
<http://dx.doi.org/10.1016/j.flora.2009.12.028>
- Nyman, C.F. (1855) *Sylloge Florae Europaeae*. Oerebroa. 442 pp.
<http://dx.doi.org/10.5962/bhl.title.11052>
- Rešetnik, I., Satovic, Z., Schneeweiss, G.M. & Liber, Z. (2013) Phylogenetic relationships in Brassicaceae tribe *Alysseae* inferred from nuclear ribosomal and chloroplast DNA sequence data. *Molecular Phylogenetics and Evolution* 69: 772–786.
<http://dx.doi.org/10.1016/j.ympev.2013.06.026>
- Schulz, O.E. (1933) *Leptoplax* O.E. Schulz. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 66: 92.
- Warwick, S.I. & Al-Shehbaz, I.A. (2006) Brassicaceae: Chromosome number index and database on CD-Rom. *Plant Systematics and Evolution* 259: 237–248.
<http://dx.doi.org/10.1007/s00606-006-0421-1>
- Warwick, S.I., Francis, A. & Al-Shehbaz, I.A. (2006) Brassicaceae: Species checklist and database on CD-Rom. *Plant Systematics and Evolution* 259: 249–258.
<http://dx.doi.org/10.1007/s00606-006-0422-0>
- Warwick, S.I., Mummenhoff, K., Sauder, C.A., Koch, M.A. & Al-Shehbaz, I.A. (2010) Closing the gaps: phylogenetic relationships in the Brassicaceae based on DNA sequence data of nuclear ribosomal ITS region. *Plant Systematics and Evolution* 285: 209–232.
<http://dx.doi.org/10.1007/s00606-010-0271-8>
- Warwick, S.I., Sauder, C.A. & Al-Shehbaz, I.A. (2008). Phylogenetic relationships in the tribe *Alysseae* (Brassicaceae) based on nuclear ribosomal ITS DNA sequences. *Canadian Journal of Botany* 86: 315–336.
<http://dx.doi.org/10.1139/b08-013>
- Willdenow, C.L. (1800) *Species Plantarum, editio quarta, tomus III*. Nauk, Berolini, 847 pp.