



Nomenclatural observations on the genus “*Bryopsidella*” (Ulvophyceae, Bryopsidaceae)

GIOVANNI FURNARI* & MARIO CORMACI

Department of Biological, Geological and Environmental Sciences, University of Catania, Section Vegetal Biology, Via A Longo, 19, 95125 Catania, Italy

* Corresponding author (furnari.giovanni41@gmail.com)

Feldmann (1970: 249), quoting the results of Hustede (1960), who had shown that *Bryopsis halymeniae* Berthold (1882: 497) was the gametophyte of a *Derbesia* sp.¹, pointed out that this species could not belong to the genus *Bryopsis* differing from it in chloroplasts with no pyrenoids and gametes without a stigma. On that basis, as well as because of its peculiar life cycle, he stated that *Bryopsis halymeniae* should be considered the type of a new genus: *Bryopsidella*. Even though Feldmann (1970) did not explicitly propose the new combination *Bryopsidella halymeniae*, such a binomial was reported by Gallardo *et al.* (1993: 412, note 60) and by Boudouresque & Perret-Boudouresque (1987: 83).

The genus *Bryopsidella*, when proposed by Feldmann (1970), was invalid because no diagnosis was provided as required by Art. 38.1 of Melbourne Code (McNeill *et al.* 2012). Moreover, it was invalid because the type species is not a validly published name as required by Art. 10.3 of Melbourne Code (McNeill *et al.* 2012). In fact, *Bryopsis halymeniae* is invalid because it is accompanied only by habitat and locality information and not by a description or diagnosis, as required by Art. 38.1 of Melbourne Code (McNeill *et al.* 2012) (Table 1). Moreover, Art. 38.10 cannot be applied to that species in that no figures were provided by Berthold (1882). The fact that the species was invalid was not recognized in the Index Nominum Algarum [<http://ucjeps.berkeley.edu/INA.html>] nor in AlgaeBase (Guiry & Guiry, 2014). Conversely, the invalid status of that species name was pointed out by Wynne (2005: 252), who however, attributed *Bryopsis halymeniae* to “Berthold ex Funk (1927: 330, fig. 17f)”. However, Funk gave no diagnosis of that species either in 1927 or in 1955 (Funk 1955: 21 and Pl. II figs 1–3) (Table 1), and figures provided by him are insufficient to validate the species because it was published after 1 January 1908 [see Arts 38.8 and 38.10 of Melbourne Code (McNeill *et al.* 2012)]. Therefore, *B. halymeniae* cannot be considered as having been validated by Funk, but remains an invalid name.

The genus *Bryopsidella* was validated by Rietema (1975: 1, footnote) who published the Latin diagnosis of the genus and definitely included its type species, referring to the validly published species *Derbesia neglecta* Berthold (1880: 77) [see Art. 10.3 of Melbourne Code (McNeill *et al.* 2012)]. It should be noted that *Bryopsidella* is generally attributed to Feldmann in Rietema [see Index Nominum Algarum and AlgaeBase (Guiry & Guiry 2014)], while Feldmann gave only the permission to Rietema to publish the Latin diagnosis of the genus. Therefore, the author of the genus is only Rietema (or Feldmann ex Rietema) and not Feldmann in Rietema.

Finally, the combination *Bryopsidella neglecta* (Berthold) Rietema (Rietema 1975: 1, footnote) is invalid because it lacks a full and direct reference to the basionym that includes the page on which the basionym was published [see Art. 41.5 of Melbourne Code (McNeill *et al.* 2012)]. Therefore, the following new combination is here proposed:

Bryopsidella neglecta* (Berthold) G. Furnari et Cormaci, *comb. nov.

Basionym: *Derbesia neglecta* Berthold (Mitt. Zool. Stat., Neapel, 2: 77, 1880).

Acknowledgements

Dr. William J. Woelkerling is acknowledged for suggestions given during the preparation of this note. Both anonymous referees are acknowledged for their comments.

1. A more complete paper showing that *Derbesia neglecta* and *Bryopsis halymeniae* are phases of one species, was later published by Hustede (1964).

TABLE 1. “Descriptions” of *Bryopsis halymeniae* given by Berthold (1882), Funk (1927) and Funk (1955), and respective English translations.

Berthold (1882: 497)
Im sommer auf <i>Halymenia floresia</i> zarte grüne Rasen bildend. Auf der Rhede von Neapel vor der Santa Lucia in 7–12 meter tiefe.
In summer on <i>Halymenia floresia</i> tender green-grass forming. On the roads of Naples in front of Santa Lucia at a 7–12 meter depth.
Funk (1927: 330)
Bildet auf <i>Halymenia floresia</i> und <i>ulvoidea</i> im Hochsommer und Herbst zarte grüne Räschen bildend. Nach Berthold bei Santa Lucia in 7–12 m Tiefe. Ich erhielt sie einmal auf den genannten Halymenien im September 1924 von der Secca di San Giovanni (15–20 m), fruktifizierend.
It forms on <i>Halymenia floresia</i> and <i>ulvoidea</i> in high summer and autumn delicate green grass. After Berthold at Santa Lucia in depth 7–12 m. I obtained it once on the mentioned Halymenien in September 1924 from the Secca di San Giovanni (15–20 m), fertile
Funk (1955: 21)
Entwickelt sich regelmässig im Sommer auf <i>Halymenia floresia</i> , seltener auf <i>H. ulvoidea</i> , auf denen sie mehr oder weniger dichte Rasen in Höhe von 1–1,5 cm bildet. Sie ist bei Santa Lucia in Tiefen von 10–20 m häufig. Fruktifizierend habe ich sie früher im September gefunden.
Developed on a regular basis in the summer on <i>Halymenia floresia</i> , rarely on <i>H. ulvoidea</i> on which it is more or less dense turf in height of 1–1.5 cm. It is common in Santa Lucia at depths of 10–20 m. I have found them fertile earlier in September

References

- Berthold, G. (1880) Zur Kenntnis der Siphoneen und Bangiaceen. *Mittheilungen aus der Zoologischen Station zu Neapel* 2: 72–82.
- Berthold, G. (1882) Über die Verteilung der Algen im Golf von Neapel nebst einem Verzeichnis der bisher daselbst beobachteten Arten. *Mittheilungen aus der Zoologischen Station zu Neapel* 3: 393–536.
- Boudouresque, C.-F. & Perret-Boudouresque, M. (1987) *A checklist of the benthic marine algae of Corsica*. GIS Posidonie publications, Marseille, 121 pp.
- Feldmann, J. (1970) Reproduction and life-cycle of siphonous marine Chlorophyceae. *XI International Botanical Congress. Seattle, Washington. Abstracts*: 249.
- Funk, G. (1927) Die Algenvegetation des Golfs von Neapel. *Pubblicazioni della Stazione Zoologica di Napoli* 7 (Supplemento): 1–507.
- Funk, G. (1955) Beiträge zur Kenntnis der Meeresalgen von Neapel: Zugleich mikrophotographischer Atlas. *Pubblicazioni della Stazione Zoologica di Napoli*. 25 (Supplemento): i–x, 1–178.
- Gallardo, T., Gómez Garreta, A., Ribera, M.A., Cormaci, M., Furnari, G., Giaccone, G. & Boudouresque, C.-F. (1993) Checklist of Mediterranean Seaweeds, II. Chlorophyceae Wille *s.l.* *Botanica Marina* 36: 399–421.
<http://dx.doi.org/10.1515/botm.1993.36.5.399>
- Guiry, M.D. & Guiry, G.M. (2014) *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. Available at <http://www.algaebase.org> (accessed 21 January 2014).
- Hustede, H. (1960) Über den Generationswechsel zwischen *Derbesia neglecta* Berth. und *Bryopsis halymeniae* Berth. *Naturwissenschaften* 47: 19.
<http://dx.doi.org/10.1007/bf00628462>
- Hustede, H. (1964) Entwicklungsphysiologische Untersuchungen über den Generationswechsel zwischen *Derbesia neglecta* Berth. und *Bryopsis halymeniae* Berth. *Botanica Marina* 6: 134–142.
<http://dx.doi.org/10.1515/botm.1964.6.1-2.134>
- McNeill, J., Barrie, F.R., Buck, W.R., Demoulin, V., Greuter, W., Hawksworth, D.L., Herendeen, P.S., Knapp, S., Prado, J., Prud'homme van Reine, W.F., Smith, G.F., Wiersema, J.H. & Turland, N.J. (2012) International Code of Nomenclature for algae, fungi and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011. *Regnum Vegetabile* 154: 1–208.
- Rietema, H. (1975) *Comparative investigations on the life-histories and reproduction of some species in the siphonous green algal genera Bryopsis and Derbesia*. University of Groningen, Groningen, 130 pp.
- Wynne, M.J. (2005) Two new species of *Bryopsis* (Ulvophyceae, Chlorophyta) from the Sultanate of Oman, with a census of currently recognized species in the genus. *Contributions from the University of Michigan Herbarium* 24: 229–256.