



## The genus *Melobesia* (Corallinales, Rhodophyta) from the subtropical South Atlantic, with the addition of *M. rosanoffii* (Foslie) Lemoine

VANESSA POLETTO BORGES<sup>1</sup>, EDUARDO BASTOS<sup>1</sup>, MANUELA BERNARDES BATISTA<sup>1</sup>, ZENILDA BOUZON<sup>2</sup>, CINTIA LHULLIER<sup>1</sup>, EDER CARLOS SCHMIDT<sup>2</sup>, MARINA NASRI SISSINI<sup>1</sup> & PAULO ANTUNES HORTA<sup>1-3</sup>

<sup>1</sup> Federal University of Santa Catarina, Botany Department, 88010-970, Florianópolis, Brazil.

<sup>2</sup> Program in Cell Biology and Development, Department of Cell Biology, Embryology and Genetics, Federal University of Santa Catarina 88049-900, CP 476, Florianópolis, SC, Brazil.

<sup>3</sup> Corresponding author. E-mail: [pahorta@ccb.ufsc.br](mailto:pahorta@ccb.ufsc.br)

### Abstract

The calcareous crusted epiphytic algae are a group of extremely delicate, fragile and poorly studied algae. The subfamily Melobesioideae (Corallinophycidae, Rhodophyta) includes the genus *Melobesia*, which has no record of molecular analysis. However, thanks to measurement data, it was possible to find enough similarity for taxonomic identification of two species: *Melobesia rosanoffii* (Foslie) Lemoine, described for the first time in South Atlantic Ocean, and *Melobesia membranacea* (Esper) Lamouroux, first described in southern Brazil. The group has undergone several changes of classification from animals to plants. Today crusty coralline algae have great importance due to the possibility of easy spore dispersal between oceans.

**Key words:** Coralline red algae, Epiphyte crustose coralline, Hapalidiaceae, Brazil

### Introduction

The subfamily Melobesioideae (Corallinophycidae, Rhodophyta) involve crustose to fruticose plants, with cells of continuous vegetative filaments linked by cell fusions; genicula absent; multiporate plate composed of cells at maturity (Taylor 1960; Harvey 2003). The algae also present tetrasporangial and bisporangial conceptacles. This subfamily includes nine genera, and among these are the delicate epiphytes as representatives of the genus *Melobesia* J.V. Lamouroux 1812:186 (Woelkerling 1988).

Lamouroux (1812) described the genus *Melobesia* for the species *M. membranacea*, and defined as calcified plants, largely or entirely pseudoparenchymatous, growing completely attached to the surface of various substrates (most commonly other algae), thallus construction dimerous throughout, upright filaments each normally terminating at the thallus surface in an epithelial cell, outermost epithelial cell walls rounded or flattened but not flared at the corners.

Gametangia and carposporangia develop in uniporate conceptacles. Spermatangia (male gametangia) and carpogonia (female gametangia) are produced in different conceptacles; male and female conceptacles are occasionally formed on the same plant but much more commonly formed on different plants. Spermatangia are formed on unbranched filaments arising from the conceptacle chamber floor and roof. Carpogonial branches 1–2 are celled with basal auxiliary cells, arising from the conceptacle chamber floor. Even after fertilization senescent carpogonial branches and the respective trycogines persist in the center of the conceptacle floor. Carposporophytes develop in these female conceptacles, after fertilization.

Tetrasporangia/bisporangia formed in conceptacles on separate plants from gametangia and carposporangia. Roofs of tetrasporangial/bisporangial conceptacles multiporate. Each pore is blocked by apical plug that prevents sporangia release, which should just occur after the complete maturation of these structures. Bisporangia each contain two spores but otherwise are similar to tetrasporangia (Woelkerling 1988; Wilks & Woelkerling 1991; Woelkerling *et al.* 1993).

Despite being understudied, epiphyte coralline diversity can present key information to the understanding of biogeographical and ecological aspects of the marine biodiversity. On the one hand, the epiphytic group distribution can help in the description of the main routes of marine diversity dispersion (van den Hoek 1987), improving hypothesis about environments connectivity. On the other hand, they play important role in reducing herbivory of flesh seaweeds

and seagrasses, a possible understudied coevolutionary process (Martin et al. 2008), that is threatened by the observed and predicted ocean acidification.

Biogeographically, *Melobesia* appears to be widespread (Pacific Islands; North, Central and South America, South Africa, Africa, Ireland, Europe, Atlantic Islands, Caribbean Islands, Western Atlantic, Indian Ocean, Islands South-west Asia, Asia, South-east Asia, Australia and New Zealand, even Antarctic and the sub Antarctic islands), but many records require confirmation, due the high complexity of performing a taxonomic study of this genus, several cases of **synonyms** and lack of molecular analysis.

Adapting methodologies of coralline red algae anatomy conventionally utilized for taxonomical studies in this group, the present study report new distribution records and evaluate the significance of characters that have been used to help delimit the specie based on morphological and reproductive features

## Material and Methods

Data were collected on the coast of Santa Catarina state, part of the warm temperate province on the Brazilian Coast (Horta et al. 2001), in five cities: Governador Celso Ramos—Praia Grande (27°21'26.26"S, 48°32'10.75"W) on May, 2012; Desert Island (27°16'13"S 48°19'56"W), August, 2012; Farol of Santa Marta (28°36'21.69"S 48°48'57.25"W) on January, 2013; city of Florianópolis – Sambaqui (27°29'24.61"S, 48°32'14.78"W) on February, 2013; Moleques Island (27°50'44.80"S, 48°25'46.77"W) on April 2013. All on the rocky shore, in the intertidal at low tide (<0.3 m). The studied specimens were deposited at the Phycological Herbarium of the Federal University of Santa Catarina (FLOR).

### *Light microscopy (LM)*

The collections were placed in plastic bags, transported to the laboratory with seawater, and latter preserved in 4% formalin-seawater, and transferred to 70% alcohol-water solution with 10% added glycerin after 24 hours. For light microscopy, decalcification was made with 10% hydrochloric acid for 20–40 minutes, dehydrated in increasing series of aqueous ethanol solutions and infiltrated with Historesin (Leica Historesin, Heidelberg, Germany), according to the instructions supplied by the manufacturer. Sectioning (6–10 µm thick) was made with a microtome, stained with acidified aqueous Toluidine blue (TB-O) (Moura *et al.* 1997, Horta 2002).

### *Scanning electron microscope (SEM)*

For the analysis in a scanning electron microscope, stubs were prepared following Horta (2002) with some modifications due the fragility of the algae were made. Conceptacle measurements were obtained from transversal fractures, perpendiculars to the thallus surface, regarding conceptacles presence, utilizing a razor blade to promote the perpendicular fracture. Conceptacle height is the distance from the conceptacle floor to the top of the chamber, and the diameter is the distance between the inner walls as described in Chamberlain (1983).

## Results

During taxonomic analysis of coralline epiphytes observed in seaweeds from Brazilian coast two species belonging to the family Hapalidiaceae and subfamily Melobesioideae. Due the presence of dimerous construction, cells of adjacent filaments joined by fusion cells and multiporated tetrasporangial conceptacles these taxa was identified as being representatives of the genus *Melobesia*. Among the evaluated specimens two dietetics morpho/anatomical aspects was observed, indicating the presence of two species: one known as *M. membranacea*, and a second as *M. rosanoffii*, a new record to South Atlantic.

*Melobesia membranacea* (Esper) Lamouroux 1812

**Basionym:** *Corallina membranacea* Esper

**Homotypic Synonyms:** *Corallina membranacea* Esper 1796; *Epilithon membranaceum* (Esper) Heydrich 1897; *Lithothamnion membranaceum* (Esper) Foslie 1898

**Type locality:** West coast of France (Dawson 1960: 8)

**Distribution in the World:** distributed worldwide.

**References to Brazil:** as *Melobesia membranacea* (Esper) J.V.Lamouroux (Taylor 1960, Oliveira Filho 1977), and as *Epilithon membranaceum* (Esper) Heydrich (Taylor, 1930).

**Specimens studied** Brazil, Santa Catarina, Desert Island; Farol of Santa Marta Sambaqui; Moleques Island.

## Description


### *Vegetative structure*

Calcareous plants reddish, purple or white (Figure 1A); Thallus encrusting, epigenous and completely affixes by cell adhesion. Structure pseudomarenchymatous with dorsiventral organization; Construction dimerous (Figure 1C), filaments composed of palisade or non-palisade cells 3–5  $\mu\text{m}$  long and 3–6  $\mu\text{m}$  high; distal wall of epithallial cells of adjacent filaments joined by cell-fusion (Figure 1D); secondary pit-connections and tricocytetes not observed.

### *Reproductive features*


Gametangia and carposporangia produced in uniporate conceptacles (Figure 1B). Tetrasporangia produced in multiporate conceptacles.

Gametangial thalli monoecious or dioecious; carpogonia and spermatangia produced in separate conceptacles. Carpogonia terminating 2–3 celled filaments arising from the female conceptacle chamber floor. Mature female carposporangial conceptacle roofs protruding above or flush with surrounding thallus surface, 18–25  $\mu\text{m}$  thick, composed of 5–8 layers of short, squat cells, conceptacle chambers 90–135  $\mu\text{m}$  in diameter and 55–97  $\mu\text{m}$  high. Carposporophytes with 12–20  $\mu\text{m}$  in diameter and 13–38  $\mu\text{m}$  high; when mature apparently consisting fusion cells bearing several gonimoblast filaments.




**FIGURE 1** Epiphytic coralline algae *Melobesia membranacea*. Image 1B. From scanning electron microscope; Images 1C and 1D from light microscopy. Fig. 1A. Plants growing on *Pterocladia capillacea* (S.G.Gmelin) Santelices & Hommersand. Fig. 1B. Uniporate conceptacle (arrow), note the detail of epithallial concavities raised above the thallus. Fig. 1C. Section through older portion of thallus showing dimerous organization. Fig. 1D. Detail of a hypothallus cell, CF indicates cell fusion.





**FIGURE 2** Reproductive features of *Melobesia membranacea*. Images from light microscopy. Fig. 2A. Carposporophytic conceptacle with mature carpospores (C). Fig. 2B. Image of a carposporophytic conceptacle with senescent carpogonial branches in the central region and carpospores (C) in the periphery.



**FIGURE 3** Reproductive features of *Melobesia membranacea*. Images from light microscopy. Fig. 3A. Detail of a male conceptacle with mature, unbranched spermatangial filaments arising from the chamber floor and roof (arrows) showing central filaments that resemble mature carpogonial branches. Fig. 3B. Section of a mature tetrasporangial conceptacle with mature tetrasporangium sporangium (T).

Spermatangial filaments arising from the floor and roof (Figure 3A), mature male conceptacles protruding above or flush with surrounding surface, 15–21  $\mu\text{m}$  thick, composed of 6–12 layers of more or less squat cells, conceptacle chambers 50–75  $\mu\text{m}$  in diameter and 47–55  $\mu\text{m}$  high.

Tetrasporangial conceptacle roofs protruding above or flush with surrounding surface, commonly but not always differentiated into a central sunken pore-plate; pore-plate composed of cells that are similar in size and shape to other roof cells, roof 5–6 cells thick above the chamber, with 10–14  $\mu\text{m}$  thick, conceptacle chambers 78–102  $\mu\text{m}$  in diameter and 60–102  $\mu\text{m}$  high; tetrasporangial initial situated one cell layer below the thallus surface, each mature sporangium with 14–21  $\mu\text{m}$  in diameter and 44–51  $\mu\text{m}$  long, containing zonately arranged tetraspores and possessing an apical plug that blocks a roof pores. Bisporangia with 35–38  $\mu\text{m}$  in diameter and 24–39  $\mu\text{m}$  long (Figure 3B).

*Melobesia rosanoffii* (Foslie) Lemoine 1912: LXI


**Basionym:** *Lithothamnion rosanoffii* Foslie 1908:5. Woelkerling 1993:191 *Epilithon rosanoffii* (Foslie) Foslie 1909:55

**Type locality:** Port Phillip Bay, Victoria, Australia (Womersley 1996: 173).

**Holotype:** TRH; includes slides 1342, 1641, and one unnumbered slide) (Womersley 1996: 173).

**Distribution in the World:** New Zealand (Nelson 2012), South Australia (Wilks & Woelkerling 1991, Womersley 1996), Tasmania (Wilks & Woelkerling 1991, Womersley 1996), and Victoria (Wilks & Woelkerling 1991, Womersley 1996, Morcom *et al.* 2005).

**Specimens studied:** Brazil, Santa Catarina, Governador Celso Ramos-Praia Grande.




**FIGURE 4.** Epiphytic coralline algae. Images 4B, 4C and 4D from scanning electron microscope. Fig. 4A. Plants growing on *Laurencia* sp. Arrows denote conceptacles. Fig 4B. Note the cell wall surface view of tetrasporangial conceptacles (TC). Fig 4C. Detail of epithallial concavities of a tetrasporangial conceptacle. Fig 4D. Note the pore plate of a young tetrasporangial multiporate conceptacle. Arrows denote pores.




## Description:

### *Vegetative structure*

Plants nongeniculate, entirely attached to the host alga by cell adhesion, encrusting (Figure 4A); construction dimerous (Figure 5); vegetative thallus 25–40  $\mu\text{m}$ , with epithelial cell 4–7  $\mu\text{m}$  in diameter and 4–8  $\mu\text{m}$  long and additional subtending cells 4–8  $\mu\text{m}$  in diameter and 5–7  $\mu\text{m}$  long; distal walls of the epithelial cells rounded or flattened (Figure 4, and figure 5A and 5B).



**FIGURE 5.** Vegetative features of *Melobesia rosanoffii* Fig 5A. Detail of margin of thallus observed by light microscopy. Primigenous cell (P) and epithelial cell (E). Fig 5B. Section through older portion of thallus, showing cell fusion (arrows). Dimerous organization.




**FIGURE 6.** Reproductive features of *Melobesia rosanoffii*. Both images from scanning electron microscope. Fig. 6A. Multiporate conceptacle, with evident two pores (arrows). Fig. 6B. Detail of tetrasporangia produced inside de conceptacle (T).

### *Reproductive features*

Gametangia and carposporangia produced in uniporate conceptacles (Figure 4B, 4C and 4D). Tetrasporangia produced in multiporate conceptacles (Figure 6). Carpogonia and spermatangia produced in separate conceptacles. Carpogonia terminating 2-celled filaments arising from the female conceptacle chamber floor. Mature female-carposporangial conceptacles roofs more or less flush with surrounding thallus surface, 17–21  $\mu\text{m}$  thick, composed of 5 layers of densely packed cells; conceptacle chambers 90–120  $\mu\text{m}$  in diameter and 60–74  $\mu\text{m}$  high. Carposporophytes when mature apparently lacking an evident fusion cell and consisting of several-celled gonimoblast filaments bearing terminal carposporangia 18–28  $\mu\text{m}$  in diameter and 16–32  $\mu\text{m}$  high. Spermatangial filaments unbranched, arising from the floor, walls and roofs of male conceptacle chambers, mature male conceptacle roofs more or less flush with surrounding thallus surface; conceptacle chambers 50–75  $\mu\text{m}$  in diameter and 34–38  $\mu\text{m}$  high (figure 7)

Tetrasporangial conceptacle roofs more or less flush with or slightly sunken below surrounding surface, central sunken pore-plate but pore-plate sharply differentiated anatomically from the remainder of roof and composed of cells that differ in size and shape from other roof cells, roof 12–18  $\mu\text{m}$  thick above the chamber; conceptacle chamber 51–83  $\mu\text{m}$  in diameter and 80–100  $\mu\text{m}$  high; tetrasporangial initials situated five or more cell layers below the thallus surface, scattered across the conceptacle chamber floor, each mature sporangium 23–35  $\mu\text{m}$  in diameter and 42–58  $\mu\text{m}$  long, containing zonately arranged tetraspores and possessing an apical plug that blocks a roof pore prior to spore release.



**FIGURE 7.** Reproductive features of *Melobesia rosanoffii* Fig 7A. Section of female-carpogonial conceptacle in vertical section with senescent carpogonial branches with persistent trychogina (GF). Fig 7B. Female conceptacle in vertical section, with carpogonial branches, arising from the floor with trychogine projected in the pore channel (arrow). Note the roof flush.

## Discussion

The coralline red algal genus *Melobesia* J.V.F. Lamouroux, 1812, Hapalidiaceae, Corallinales, Rhodophyta (see Harvey *et al.* 2003 for classification details), was established for three taxa: *M. membranacea*, *M. orbiculata*, and *M. verrucata*. Lamouroux considered *membranacea* to be an animal, but under the current ICBN, the name is treated as validly published as a plant in 1796 (Art. 45.4).

To the subfamily Melobesioideae consists now a total of 116 species (and infraspecific) names in the database, of which 17 are flagged as currently accepted taxonomically. At present, most species assigned to *Melobesia* are poorly or incompletely known (Guiry 2014). The record of *M. rosanoffii* is represented by a single collection from the northern tip of the North Island, growing as an epiphyte on *Lessonia* sp. and there were multi and uniporate conceptacles on adjacent plants. The identity of the specimen was confirmed by Adele Harvey based on sections. This is the first time that this specie is described from the Brazilian Coast.

Using diagnostic features employed in the taxonomy of non-geniculate coralline algae, the specimens were compared with the same taxa described from other parts of the world (Table 1), and identifications based on a combination of characteristics (Lamouroux 1812; Wilks and Woelkerling 1991; Womersley 1996).

Considering that epiphytic organisms may disperse more easily, we can start from the hypothesis that the diversity of the group would be higher than the presently described. Besides being cryptic, the diffusion of the equipment and the specific methodological approaches should improve the efforts investigating the diversity of this group. We can not forget that we are passing through a very dynamic moment where climatic changes, that increase the influence of the Indic in the South Atlantic Ocean (Beal *et al.* 2011), and the intensification of global marine transport (Whinam *et al.* 2005, Gregory 2009), can represent the necessary conditions to this group take more frequently hitchhiking from Indo/Pacific oceans to southwestern Atlantic coast. Thereby, the occurrence of *M. rosanoffii* probably results from the lack

of studies about epiphytic coralline algae on the Brazilian coast. However, we cannot ignore the possibility of recent arrival, as has been observed in the invasion of other taxa (Macreadie et al. 2011). These each time more connected oceans represent important challenge to the epiphytic coralline algae current taxonomists, once traditional approaches should be limited to provide robust answers requested nowadays.

Despite the modern analyses of the material provide good morphoanatomical information, the systematics should consider the possibility of the occurrence of cryptic species, as observed in other coralline red algae taxa (Bittner et al 2011). This particularity reinforces the need of additional molecular data for a better characterization of this group.

**TABLE 1.** Comparative table. A. Present study (*M. rosanoffii*); B. Present study (*M. membranacea*). C and D. Data from Wilks and Woelkerling (1991), about the species concerned. ND – no data.

Character	A	B	C	D
Thallus thickness	25–40	13–60	25–113	10–75
Epitelial cell length	3–5	3–5	2–3	2–8
Epitelial cell diameter	3–5	3–6	2–3	2–10
Postigenous cell length	5–7	4–10	2–15	2–15
Postigenous cell diameter	4–8	6–21	2–15	2–15
Tetrasporangial conceptacle chamber diameter	51–83	78–102	45–118	50–252
Tetrasporangial conceptacle chambre high	50–90	60–102	40–80	15–83
Tetrasporangial conceptacle roof thickness	12–18	10–14	12–25	ND
Number of cell layers in tetrasporangial conceptacle roof	4–6	5–6	2–7	2–5
Lengh of tetrasporangia	42–58	44–51	25–63	10–53
Diameter of tetrasporangia	23–35	14–21	10–25	22–65
Male conceptacle chamber diameter	50–75	50–75	100–138	57–130
Male conceptacle chamber high	34–38	47–55	30–75	25–40
Carposporangial conceptacle chamber diameter	90–120	90–135	100–193	50–250
Carposporangial conceptacle chamber high	60–74	55–97	30–55	25–80
Carposporangial conceptacle roof thickness	17–21	18–25	35–63	5–25
Number of cell layers in carposporangial conceptacle roof	5–8	5–8	5–14	2–8
Diameter of carposporangia	18–28	12–20	12–20	18–60
Length of carposporangia	16–32	13–38	12–25	ND

## Acknowledgments

We thank CAPES and CNPq (PROTAX) for research funds and scholarships provided, and also the authors would like to acknowledge the staff of the Central Laboratory of Electron Microscopy (LCME), Federal University of Santa Catarina, Florianopolis, Santa Catarina, Brazil, for the use of their transmission electron microscopy. Éder C. Schmidt holds a postdoctoral fellowship from CAPES. This study is part of the MSc dissertation of the first author.

## References

- Chamberlain, Y.M. (1985) The typification of *Melobesia membranacea* (Esper) Lamouroux (Rhodophyta, Corallinales). *Taxon* 34: 673–677. <http://dx.doi.org/10.2307/1222213>
- Chamberlain, Y.M. (1983) Studies in the Corallinales with special reference to Fosliella and Pnecophyllum in the British Isles. *Bulletin of British Museum* 11: 291–463.
- Dixon, P.S. (1970) The Rhodophyta: Some aspects of their biology ii. *Oceanography and Marine Biology: An Annual Reviews* 8: 307–325.
- Gray, J.E. (1864) *Handbook of British water-Weeds or algae*. Vol. 22. London: R. Hardwicke, pp. i–iv, 1–123.



- Guiry, M.D. & Guiry, G.M. (2014) *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. Available from: <http://www.algaebase.org> (accessed 30 January 2014).
- Harvey, A.S., Broadwater, S.T., Woelkerling, W.J. & Mitrovski, P.J. (2003) *Choreonema* (Corallinales, Rhodophyta): 18S rDNA phylogeny and resurrection of the Hapalidiaceae for the subfamilies Choreonematoideae, Austrolithoideae and Melobesioideae. *Journal of Phycology* 39: 988–998.  
<http://dx.doi.org/10.1046/j.1529-8817.2003.02158.x>
- Harvey, A.S., Woelkerling, W. J. & Millar, A.J.K. (2003) An account of the Hapalidiaceae (Corallinales, Rhodophyta) in south-eastern Australia. *Australian Systematic Botany* 16: 647–698.  
<http://dx.doi.org/10.1071/sb03008>
- Horta, P.A. (2002) Bases para a identificação das coralináceas não articuladas do litoral brasileiro – uma síntese do conhecimento. *Biotemas*, 15: 7–44.
- LamourouX, J.V.F. (1812) Sur la classification des Polypiers coralligènes non entièrement pierreux. *Nouveau Bulletin des Sciences par la Société Philomathique de Paris* 3: 181–188.
- Le Gall, L. & Saunders, G.W. (2007) A nuclear phylogeny of the Florideophyceae (Rhodophyta) inferred from combined EF2, small subunit and large subunit ribosomal DNA: establishing the Corallinophycidae *subclassis nov.* *Molecular Phylogenetics and Evolution* 43: 1118–1130.  
<http://dx.doi.org/10.1016/j.ympev.2006.11.012>
- McNeill, J., Barrie, F.R., Burdet, H.M., Demouline, V., Hawksworth, D.L., Marhold, K., Nicolson, D.H., Prado, J., Silva, P.C., Skog, J.E., Wiersema, J.H. & Turland, N.J. (2006) *International Code of Botanical Nomenclature (Vienna Code)* adopted by the Seventeenth International Botanical Congress Vienna, Austria, July 2005. pp. [i–iv], v–xviii + 1–568. Liechtenstein: A.R.G. Gantner Verlag.
- Morcom, N.F., Woelkerling, W.J. & Ward, S.A. (2005) Epiphytic nongeniculate corallines found on *Laurencia filiformis* f. *heteroclada* (Rhodomelaceae, Ceramiales). *Phycologia* 44: 202–211.  
[http://dx.doi.org/10.2216/0031-8884\(2005\)44\[202:encfol\]2.0.co;2](http://dx.doi.org/10.2216/0031-8884(2005)44[202:encfol]2.0.co;2)
- Moura, C.W. do N., Kraus, J.E. & Cordeiro-Marino, M. (1997) Metodologia para obtenção de cortes histológicos com historresina e coloração com azul de toluidina para algas coralináceas (Rhodophyta, Corellinales). *Hoemea* 24: 17–27.
- Nelson, W.A. (2012) Phylum Rhodophyta: red algae. In: *New Zealand inventory of biodiversity. Volume Three. Kingdoms Bacteria, Protozoa, Chromista, Plantae, Fungi*. (Gordon, D.P. Eds), pp. 327–346.
- Oliveira Filho, E.C. de (1977) *Algas marinhas bentônicas do Brasil. Tese de Livre Docencia*. São Paulo: Instituto de Biociências, Universidade de São Paulo. pp. 1–406.
- Silva, P.C., Basson, P.W. & Moe, R.L. (1996) Catalogue of the benthic marine algae of the Indian Ocean. *University of California Publications in Botany* 79: 1–1259.
- Taylor, W.R. (1930) A synopsis of the marine algae of Brazil. *Revue Algologique* 5: 1–35.
- Taylor, W.R. (1960) Marine algae of the eastern tropical and subtropical coasts of the Americas. Ann Arbor: The University of Michigan Press, pp. xi + 870, 14 figs, 80 plates.
- van den Hoek, C. (1987) The possible significance of long-range dispersal for the biogeography of seaweeds. *Helgoländer Meeresunters* 41: 261–272.  
<http://dx.doi.org/10.1007/bf02366191>
- Wilks, K.M. & Woelkerling, W.J. (1991) Southern Australian species of *Melobesia* (Corallinales, Rhodophyta). *Phycologia* 30: 507–533, 37 figs, 2 tables.  
<http://dx.doi.org/10.2216/i0031-8884-30-6-507.1>
- Woelkerling, W.J. & Chamberlain, Y.M. (2007) The genus *Melobesia* (Hapalidiaceae, Corallinales, Rhodophyta): typification update. *Phycologia* 46: 232–234.  
<http://dx.doi.org/10.2216/06-93.1>
- Woelkerling, W.J. & Mitrovski, P.J. (2003) *Choreonema* (Corallinales, Rhodophyta): 18S rDNA phylogeny and resurrection of the Hapalidiaceae for the subfamilies Choreonematoideae, Austrolithoideae and Melobesioideae. *Journal of Phycology* 39: 988–998.  
<http://dx.doi.org/10.1046/j.1529-8817.2003.02158.x>
- Woelkerling, W.J. (1988) The Coralline Red Algae: an analysis of the genera and subfamilies of nongeniculate Corallinales. pp. i–xi, 1–268, 259 figs, tables numbered by chapter. *London & Oxford: British Museum (Natural History) & Oxford University Press*.
- Woelkerling, W.J., Irvine, L.M. & Harvey, A.S. (1993) Growth-forms in non-geniculate coralline red algae (Corallinales, Rhodophyta). *Australian Systematic Botany* 6: 277–293.  
<http://dx.doi.org/10.1071/sb9930277>
- Woelkerling, W.J., Millar, A.J.K., Harvey, A. & Baba, M. (2008) Recognition of *Pachyarthron* and *Bossillaas* distinct genera in the Corallinales, subfamily Corallinoideae (Coarllinales, Rhodophyta). *Phycologia* 47: 265–293.  
<http://dx.doi.org/10.2216/ph07-85.1>
- Womersley, H.B.S. (1987) The marine benthic flora of southern Australia. *Part II*. pp. 481, 169 figs, 1 table, 8 plates, 4 maps. Adelaide: South Australian Government Printing Division.
- Womersley, H.B.S. (1996) The marine benthic flora of southern Australia - *Part IIIB - Gracilariales, Rhodymeniales, Corallinales and Bonnemaisoniales*. Vol. 5 pp. 1–392, 160 figs. Canberra & Adelaide: Australian Biological Resources Study & the State Herbarium of South Australia.
- Womersley, H.B.S. (1996) The marine benthic flora of southern Australia - *Part IIIB - Gracilariales, Rhodymeniales, Corallinales and Bonnemaisoniales*. Vol. 5 pp. 1–392, 160 figs. Canberra & Adelaide: Australian Biological Resources Study & the State Herbarium of South Australia.