



USING Next-Generation Sequencing (NGS) TO UNCOVER DIVERSITY OF WOOD-DECAYING FUNGI IN NEOTROPICAL ATLANTIC FORESTS

ALINE B.M. VAZ^{1,5}, PAULA L.C. FONSECA¹, LAURA R. LEITE³, FERNANDA BADOTTI⁴, ANNA C.M. SALIM³, FLAVIO M.G. ARAUJO³, SARA CUADROS-ORELLANA³, ÂNGELO A. DUARTE², CARLOS A. ROSA¹, GUILHERME OLIVEIRA⁶ & ARISTÓTELES GÓES-NETO^{1,7}

¹Department of Microbiology, Federal University of Minas Gerais (UFMG), Belo Horizonte, MG, 31270-901, Brazil.

²Department of Technology, State University of Feira de Santana (UEFS), Feira de Santana, BA, 44036900, Brazil.

³Oswaldo Cruz Foundation (FIOCRUZ-MG), Belo Horizonte, MG, 30.190-002, Brazil.

⁴Department of Chemistry, Federal Center of Technological Education of Minas Gerais (CEFET-MG), Belo Horizonte, MG, 30480-000, Brazil.

⁵Faculdade de Minas (FAMINAS), Belo Horizonte, MG, 31744-007, Brazil.

⁶Vale Institute of Technology - Sustainable Development Belém, PA, 66055-090, Brazil.

⁷Author for correspondence: arigoesneto@pq.cnpq.br / arigoesneto@gmail.com

Abstract

A targeted amplicon-based metagenomics approach (metabarcoding) provides detailed access to the diversity of the mycobiome in any substrate in distinct environments on Earth. Fungi are the main decomposers of lignocellulosic woody debris in terrestrial forested ecosystems, contributing significantly to the global carbon cycle. The main objectives of this study were to assess the fungal taxonomic diversity in fallen woody debris samples from two Neotropical forest fragments (rainforest and seasonal forest), to analyze the qualitative and quantitative components of the taxonomic diversity, and to investigate the functional diversity of the ecological groups detected. Our study comprised three main methodological steps: (i) sampling in the field; (ii) extraction of DNA, amplification of targeted segments and massively parallel sequencing; and (iii) data analysis and interpretation. A total of 110 molecular operational taxonomic units showing sequence similarity of 95% or more across the two collection sites using two DNA metabarcoding markers (*ITS1* and *ITS2*) were assigned to putative fungal genera in 59 families, 27 orders, and 3 phyla. The number of putative fungal genera and the relative abundance of reads for each genus are higher in the tropical rainforest site than in the tropical seasonal forest site. Most of the identified genera are ligninolytic and cellulolytic and/or hemicellulolytic Basidiomycota (Agaricomycetes) and Ascomycota (Sordariomycetes), but “sugar fungi” and fungi associated with plants and detritivorous insects were also detected. This is the first study using NGS as a rapid and large-scale useful strategy to uncover the diversity of wood-decaying fungi in tropical forests.

Key words: fungal metabarcoding, rainforest, seasonal tropical forest, lignocellulosic residues

Introduction

The plant litter in terrestrial forest ecosystems consists of dead plant parts, including both non-woody and woody plant residues, with the latter comprising twigs, stems, branches and trunks, which are collectively referred to as woody debris (WD) (Cannon & Sutton 2004). Woody material refers not only to proper wood but also to bark (outer and inner) and sap (from inner bark and sapwood) at any stage of decay. Saproxylic organisms include any species that depend, at least during some portion of their lifecycle, upon wounded or decayed woody material from living, weakened or dead trees (Stokland *et al.* 2012). Saproxylic biota degrade woody material by fragmentation or physical destruction (microfauna) and by mainly aerobic, but also anaerobic, enzymatic action (fungi and bacteria) (Hattaka 2001, Stokland *et al.* 2012). The predominant mycelial habit, along with the ability of fungi to produce extracellular lignocellulolytic enzymes that degrade dead lignified plant residues, make fungi the most important group associated with wood degradation (Zhou & Ingram 2000).

The initial decay process involves fungal communities of endophytes, pathogens and saprotrophs, that are present in plant tissues (Boddy 2001). The ecological functional groups of saproxylic fungi include the following: (a) “sugar

and staining fungi”, which degrade simple carbon compounds found in cell contents and saps in the initial stages of wood decomposition (mainly ascomycotan and basidiomycotan yeasts and a number of mycelial ascomycotan groups, but also Mucoromycotina); (b) structural wood-decaying fungi, which have full enzymatic ability to degrade cellulose, hemicelluloses and lignin (mainly Basidiomycota and certain groups of Ascomycota); (c) residual wood-decaying fungi, which use the products resulting from the decomposition of wood by structural wood-decaying fungi; (d) fungi associated with detritivorous or saprophagous animals (mainly insects of the orders Coleoptera, Isoptera and Diptera, as well as Acari), which can be sap feeders, inner bark consumers (phloemophagous) and wood consumers (xylophagous, xylemophagous); (e) animal-predatory (nematode-trapping) fungi; (f) mycoparasites (both biotrophic and necrotrophic) and fungicolous fungi; and (g) mycorrhizal fungi (Stokland *et al.* 2012).

Structural wood-decaying fungi are the main decomposers of wood and are usually categorized as white-, brown- and soft-rot fungi according to the mode of wood decay (Hattaka 2001). White-rot fungi degrade all of the structural components of wood (cellulose, hemicelluloses and lignin), either simultaneously or sequentially (selective delignification), whereas brown-rot fungi degrade cellulose and hemicelluloses, leaving the lignin fraction nearly unaltered (Morgenstern *et al.* 2008). All known white- and brown-rot wood-decaying fungi are basidiomycetes, and recent evidence based on phylogenomics suggests a continuum rather than a dichotomy between the white- and brown-rot modes of wood decay in Basidiomycota (Riley *et al.* 2014). Soft-rot fungi, which are restricted to Ascomycota, degrade the cellulose and hemicellulose fractions, but only in the central layer (S2) of the secondary cell walls, and they have limited capacity to modify lignin (Schwarze 2007).

The targeted amplicon-based metagenomics approach of massively parallel sequencing, a molecular, culture-independent approach, theoretically provides the most detailed access to the diversity of the mycobiome of any substrate. Nevertheless, possible methodological biases, limitations of the markers and bioinformatic analysis may lead to incorrect conclusions (Lindahl *et al.* 2013). In general, high fungal taxonomic diversity has been observed in various environmental compartments (air, soil, continental and oceanic sediments, and surface and ground water), as well as on the external and internal surfaces and microenvironments within macroorganisms. These studies typically consider molecular operational taxonomical units (MOTUs) to differentiate the environmental sequences at the species or genus level (Cuadros-Orellana *et al.* 2013). The internal transcribed spacer and 5.8S region of the nuclear ribosomal repeat unit have been used as the primary DNA fungal barcoding marker for species delimitation (Schoch *et al.* 2012). Both the ITS1 and ITS2 spacer regions can be used as DNA metabarcodes in fungal metagenomic studies (Blaalid *et al.* 2013). However, most of the NGS platforms commonly used (Roche, Life Technologies, Illumina) generate short reads (Goodwin *et al.* 2016), preventing the reliable identification at species level.

To date, studies specifically investigating the mycobiomes of decaying wood using targeted, amplicon-based metagenomics are few, and all of these studies were conducted strictly in palearctic boreal and temperate forests (Ovaskainen *et al.* 2010, Kubartova *et al.* 2012, Ovaskainen *et al.* 2013, Ottosson *et al.* 2015, Van der Wal *et al.* 2015, Hoppe *et al.* 2015, Jang *et al.* 2015, Runnel *et al.* 2015, Yamashita *et al.* 2015). Thus, as far as we know, there has not been a similar metabarcoding study in tropical, and more specifically, neotropical forests. Although there are many studies comprising collection, direct observation and/or culturing of fungi from decaying wood in the Atlantic Forest biome, (see Maia *et al.* 2015 for an extensive review of the literature), a metabarcoding study theoretically allows the most comprehensive access to the fungal diversity in any kind of substrate. Thus, the objective of our study is to characterize the fungal diversity in fallen woody debris from two climatically and vegetationally distinct, neotropical forest fragments using a targeted, amplicon-based (nrITS) metagenomics approach.

Material & Methods

Study areas

There are two main types of tropical broadleaf forests distributed across both the Neotropical and Paleotropical phytogeographical zones: (i) tropical rainforests, with high temperatures and high rainfall conditions throughout the year, and (ii) tropical seasonal forests, with year-round high temperatures but a clearly seasonal drought (Evert & Eichorn 2013). In this work two study areas were selected as representatives of each main type of tropical broadleaf forest: Rio Doce State Park (RDSP) and Serra do Conduru State Park (SCSP). The permissions for the field studies were obtained from the Instituto Estadual de Florestas (IRF): COL:045/12 for RDSP park and Instituto do Meio Ambiente e Recursos Hídricos (INEMA): 2013-009569/TEC/PESQ-0036 for SCSP park.

RDSP is situated in the state of Minas Gerais, Brazil (19°29’S–19°48’S and 42°28’W–42°38’W, in the interior

sub-region according to Ribeiro *et al.* 2009), with an altitude ranging from 230–515 m and a total area of 36,113 ha. The main vegetation is seasonally dry tropical forest, and the regional climate is classified as tropical wet and dry climate, Aw in the Köppen climate classification system, with a mean annual temperature of 23°C and a mean annual rainfall of 1300 mm. This area is characterized by a marked seasonality, with a humid season with intense precipitation in the summer and a dry season in the winter (Lopes *et al.* 2002). The surrounding areas near our sampling collection site are often visited by tourists leading to some anthropogenic impact.

SCSP is situated in the state of Bahia, Brazil (14°20'–14°30' S; 39°02'–39°08' W, in the Bahia sub-region according to Ribeiro *et al.* 2009), with an altitude ranging from 60–500 m and a total area of 9,275 ha. The main vegetation is tropical rainforest, and the regional climate is classified as tropical rainforest climate, Af in the Köppen climate classification system, with a mean annual temperature of 24°C and a mean annual rainfall of 2000 mm, which is well distributed throughout the year, i.e., without a dry season (Martini *et al.* 2007). In both study areas, the sample collection sites consisted of the main vegetation type: an open canopy environment near the lacunar system in RDSP and a dense canopy environment in SCSP.

Sample Collection

A total of 20 samples (10 per collection site) of fallen woody debris (FaWD) was randomly collected. The FaWD samples were collected in accordance to the following exclusion and inclusion criteria: (i) larger than 2 cm in diameter, (ii) including wood and bark, (iii) without visible epixylic lichens or bryophytes, (v) at least 5 m far apart from each other, and (vi) representing all stages of decay, according to Keller *et al.* (2004). Five subsamples were taken using a sterilized scalpel from different parts of each FaWD, and subsequently pooled forming one compound sample, which were placed in sterilized plastic bags, transported to the laboratory within 4 h, and stored at -20°C until processed.

DNA extraction, amplification, and massively parallel sequencing

Subsamples (1 g) from each wood sample were ground under liquid nitrogen, and metagenomic DNA was extracted from 100 mg samples according to Góes-Neto *et al.* (2005) with the following modifications: salt extraction buffer [(0.05 M Tris-HCl (pH 9), 0.005 M EDTA, 0.1 M NaCl, 1% SDS, 3% β-mercaptoethanol 3% polyvinyl-pyrrolidone (PVP)] and Proteinase K (50 µg/ml). The internal transcribed spacer (ITS) region of the rRNA gene was amplified using the primers ITS4 (5'-TCCTCCGCTTATTGATATGC-3') and ITS5 (5'-GGAAGTAAAAGTCGTAACAAGG-3') (Schoch *et al.* 2012). Each PCR reaction contained the following components: 1× PCR reaction buffer (100 mM Tris-HCl), 2.5 mM MgCl₂, 0.2 mM dNTP, 15 pmol of each oligonucleotide primer, 5 units Taq polymerase (Kappa, USA), 1 M betaine, 1 µg of BSA, 2% of DMSO and approximately 1 ng genomic DNA, and sterilized, deionized water. Cycle parameters were as follows: 1 initial denaturation cycle at 94°C for 2 min, followed by 35 denaturation cycles at 94°C for 1 min, and annealing at 60°C for 1 min, and extension at 72°C for 3 min, with a final extension cycle at 72°C for 5 min. At least three independent amplification reactions were performed from the same DNA extract. The PCR products were purified, quantified and combined until 100 ng of purified PCR product was obtained for each sample. Samples from the same collection site (RDSP or SCSP) were pooled in an equimolar proportion to produce a composite sample with a final content of 1 µg of purified PCR product, which was quantified using a DNA fluorescence assay.

An Ion Torrent adapter-ligated library was constructed following the Ion Fragment Library Kit (Life Technologies) protocol (Part #4467320 Rev. A). Briefly, fragmentation of 1 µg of each composite sample was performed using Adaptive Focused Acoustics™ (AFA; Covaris). After fragmentation, samples were end-repaired, and the Ion Torrent adapters P1 and A were ligated using DNA ligase. Following AMPure bead purification (Beckman Coulter, Brea, CA, USA), adapter-ligated products were nick-translated and PCR-amplified for a total of 10 cycles. The resulting library was also purified using AMPure beads, and the concentrations and sizes of fragmentation products were determined using an Agilent BioAnalyzer DNA 1000 Kit (Agilent Technologies). Sample emulsion PCR, emulsion breaking, and enrichment were performed using the Ion Xpress Template Kit (Part #4467389 Rev. B) according to the manufacturer's instructions. Briefly, an input concentration of one DNA template copy/Ion Sphere Particle (ISP) was added to the emulsion PCR master mix, and the emulsion was generated using an IKADT-20 mixer (Life Technologies). Next, the ISPs were recovered, and Dynabeads MyOne Streptavidin C1 beads (Life Technologies) were used to enrich for template-positive ISPs. ISP enrichment was confirmed using the Qubit 2.0 fluorometer (Life Technologies), and the sample was prepared for sequencing using the Ion Sequencing Kit protocol (Part #4467391 Rev. B). Each composite sample was loaded onto an Ion 316 chip and sequenced on the PGM (Personal Genome Machine) system for 110 cycles (Rothberg *et al.* 2011).

Data analyses

All of the generated sequences were deposited in (a) NCBI BioProject: PRJNA255944, BioSample: SAMN02934078, SAMN02934079 and (b) MG-RAST Locality 2 ITS1: 4576861.3 ITS2: 4576862.3, Locality 3 ITS1: 4576871.3 ITS2: 4576872.3. This study is part of the Brazilian Microbiome Project (BMP): <http://www.brmicrobiome.org/> (Pylro *et al.* 2013). The datasets were analyzed on a CentOS release 6.6 system. Sequences of short length (< 50 bp) and low quality (mean Phred score of $Q < 20$) with ambiguous character states (non-IUPAC) (maxambig = 0) and homopolymers longer than 8 bp (maxhomop = 8) were filtered out using PRINSEQ v0.15 (Schmieder & Edwards 2011). PCR amplification biases (overrepresented fragments) were also removed using USEARCH v.7.0.1090 (Edgar 2010). Subsequently, the FungalITSextractor was used for splitting ITS1 and ITS2 sequences (Nilsson *et al.* 2010), which were then clustered employing 97% stringency and filtered using a reference-based chimera filtering by using USEARCH v.7.0.1090 (Edgar 2010). Reads matching the internal transcribed spacers ITS1 or ITS2 were independently compared to the UNITE database (Kõljalg *et al.* 2013), using BLASTn v. 2.2.27 (E-value $\leq 1e-10$, minimum overlap length $\geq 90\%$, minimum identity $\geq 95\%$, Camacho *et al.* 2008), assigned to putative genera (Jumpponen & Jones 2009), and a fungal abundance table was generated. Non-informative names in the taxonomy dependent assignment were excluded prior to the diversity analyses (uncultured, fungal, fungi, fungus, isolate, *mycota, *mycetidae, leaf litter, unassignment, endophytic, endophyte, *mycetes, *mycete, *aceae, *ales, cf., symbiont, mycorrhiza, aff.) The number of fungal genera and the relative abundance of reads at the phylum, class and order levels were compared using barplots. All analyses were performed using R software (R Development Core Team 2016).

The ecological functional roles of the identified genera were investigated based on evidence at the gene and biochemical levels using the presence or absence of genes and/or enzymes related to lignocellulosic decomposition (degradation of cellulose, hemicelluloses and lignin), based on the CAZY (Carbohydrate-Active Enzymes: <http://www.cazy.org>), BRENDA (www.brenda-enzymes.org), NCBI (www.ncbi.nih.gov) databases, and the literature (Zhao *et al.* 2013, Riley *et al.* 2014). A majority consensus of these databases and literature was used to assign the ecological functional roles of putative fungal genera.

Results

The sequencing resulted in a total of 2.55 Mb, of which 687,662 and 1,861,279 reads were obtained for RDSP and SCSP, respectively (Table 1). Using both metabarcodes, a total of 110 genera was identified, 71 in RDSP and 85 in SCSP. In RDSP, 19.7% of genera were identified with ITS1, 43.7% with ITS2, and 36.6% with both markers. In SCSP, 18.8% of genera were identified with ITS1, 37.7% with ITS2, and 43.6% with both genomic regions.

TABLE 1. Summary of descriptive statistics.

	RDSP	SCSP	Total
Raw reads	687,662 (26.98%)	1,861,279 (73.02%)	2,548,941
Reads passing quality filter	628,970 (25.91%)	1,798,479 (74.09%)	2,427,449
Median read length (bp)	116	199	-
% G+C	43.48 \pm 8.61	47.06 \pm 5.58	-
Reads after 97% similarity clustering	259,138 (17.25%)	1,243,319 (82.75%)	1,502,547
Reads after FungalITSExtractor	114,696 (9.98%)	1,034,128 (90.02%)	1,148,824
ITS1	46,144 (10.76%)	382,758 (89.24)	428,902
ITS2	67,985 (10.38%)	587,037 (89.62%)	655,022
Reads assigned to any MOTUs			
ITS1	39,915 (30.32%)	91,720 (69.68%)	131,635
ITS2	59,999 (37.22%)	101,190 (62.78%)	161,189
Both markers	36,567 (30.3%)	84,109 (69.7%)	120,676
MOTUs (genus level) after all filters			

... Continued on next page

TABLE 1. (Continued)

	RDSP	SCSP	Total
ITS1	41	54	-
ITS2	58	70	-
Both markers	26	37	-

Note: Percentage values were calculated by line.

Regardless of the collection site or metabarcode used, a higher number of genera of Ascomycota than Basidiomycota were detected, and Zygomycota only in the SCSP (Fig. 1). The same pattern was observed when the relative abundance of reads was considered (Fig. 2).




FIGURE 1. Genera richness at the levels of phylum, class and order per study area and metabarcode using 95% sequence similarity with reference sequences in the UNITE database as surrogate for traditional taxonomic generic concepts.

Sordariomycetes (Ascomycota) and Agaricomycetes (Basidiomycota) exhibited the highest number of genera in the two collection sites. However, the relative abundances of reads in Saccharomycetes were considerably higher than all the other fungal classes.

At the ordinal level, Hypocreales (Ascomycota) showed the highest number of genera regardless of the collection site or metabarcode. Agaricales (Basidiomycota) showed the highest number of genera in the SCSP whereas Polyporales (Basidiomycota) was the highest in RDSP. However, when analyzing the total number of sequences, Saccharomycetales (Ascomycota) was the most abundant.

The genera that showed the highest number of sequences in RDSP were *Meyerozyma* (34%), *Scytalidium* (23%), *Resinicium* (16.7%), *Verticicola* (4.79%), *Pichia* (4.48%) and *Candida* (1.85%) whereas in SCSP were *Meyerozyma* (34.3%), *Pichia* (10.58%), *Asterostroma* (9.84%), *Resinicium* (8.35%), *Scytalidium* (7.48%), *Candida* (6.1%), and *Verticicola* (3.50%). Furthermore, when considering only ITS1, *Meyerozyma* and *Resinicium* were those with the highest number of reads, and for ITS2 were *Scytalidium* and *Verticicola*. All together, these seven genera (*Meyerozyma*, *Scytalidium*, *Resinicium*, *Pichia*, *Asterostroma*, *Candida*, *Verticicola*) represented 86.4% and 80.2% of the relative abundances of the reads for RDSP and SCSP, respectively (Fig. 2, and Table 2).




FIGURE 2. Relative abundances of reads of fungal genera per study area and metabarcode using 95% sequence similarity with reference sequences in the UNITE database as surrogate for traditional taxonomic generic concepts.

TABLE 2. Relative abundances of reads of fungal genera per study area and metabarcoding. Generic names were selected using a BLAST search of 95% sequence identity and names are based on the taxonomic concepts applied in the UNITE database.

Genus	RSDP		SCSP		Phylum	Class	Order	Family	Biochemical Activity ¹	Main Ecological Group		
	ITS1	ITS2	Total	ITS1							ITS2	Total
<i>Acanthostigma</i>	0	0	0	0	0.148	0.148	Ascomycota	Dothideomycetes	Pleosporales	Tubeufiaceae	HC	saprotroph
<i>Acremonium</i>	0.12	0.218	0.338	0.496	1.33	1.826	Ascomycota	Sordariomycetes	Hypocreales	no rank	HC	saprotroph
<i>Agaricus</i>	0	0	0	0.062	0	0.062	Basidiomycota	Agaricomycetes	Agaricales	Agaricales	LHC	saprotroph
<i>Anrodia</i>	0	0	0	0.062	0	0.062	Basidiomycota	Agaricomycetes	Polyporales	Fomitopsidaceae	HC	saprotroph
<i>Arthrobotrys</i>	0	0.073	0.073	0	0.074	0.074	Ascomycota	Orbiliomycetes	Orbiliales	Orbiliaceae	HC	saprotroph
<i>Arthrographis</i>	0.723	1.16	1.883	0.186	1.996	2.182	Ascomycota	Dothideomycetes	no rank	Eremomycetaceae	HC	saprotroph
<i>Aspergillus</i>	0	0	0	0	0.148	0.148	Ascomycota	Eurotiomycetes	Eurotiales	Trichocomaceae	HC	saprotroph
<i>Asterostroma</i>	3.133	0.145	3.278	12.392	6.8	19.192	Basidiomycota	Agaricomycetes	Russulales	Lachnoladiaceae	LHC	saprotroph
<i>Auerswaldia</i>	0	0	0	0	0.074	0.074	Ascomycota	Dothideomycetes	Dothideales	Dothideaceae	HC	saprotroph
<i>Aureobasidium</i>	0	0	0	0	0.148	0.148	Ascomycota	Dothideomycetes	Dothideales	Dothioraceae	HC	saprotroph
<i>Beauveria</i>	0	0	0	0	0.074	0.074	Ascomycota	Sordariomycetes	Hypocreales	Cordycipitaceae	HC	parasitic or mutualistic symbiont of animals
<i>Calocera</i>	0.12	1.668	1.788	0.124	1.183	1.307	Basidiomycota	Dacrymycetes	Dacrymycetales	Dacrymycetaceae	HC	saprotroph
<i>Candida</i>	2.651	1.378	4.029	7.373	4.582	11.955	Ascomycota	Saccharomycetes	Saccharomycetales	no rank	HC	parasitic or mutualistic symbiont of animals
<i>Chaetosphaeria</i>	0.723	0.725	1.448	0	0.148	0.148	Ascomycota	Sordariomycetes	Sordariales	Chaetosphaeriaceae	HC	saprotroph
<i>Chaunopycnis</i>	0	0	0	0.062	0.074	0.136	Ascomycota	Sordariomycetes	Hypocreales	Ophiocordycipitaceae	HC	parasitic or mutualistic symbiont of plants

...Continued on next page

TABLE 2. (Continued)

Genus	RSDP		SCSP		Phylum	Class	Order	Family	Biochemical Activity ¹	Main Ecological Group		
	ITS1	ITS2	Total	ITS1							ITS2	Total
<i>Chloridium</i>	0.361	0.218	0.579	0.991	1.33	2.321	Ascomycota	Sordariomycetes	Chaetosphaeriales	Chaetosphaeriaceae	HC	saprotroph
<i>Cladophialophora</i>	0	0	0	0	0.074	0.074	Ascomycota	Eurotiomycetes	Chaetothyriales	Herpotrichiellaceae	HC	parasitic or mutualistic symbiont of animals
<i>Cladosporium</i>	0.602	0.943	1.545	0.062	1.33	1.392	Ascomycota	Dothideomycetes	Capnodiales	Davidiellaceae	HC	parasitic or mutualistic symbiont of plants
<i>Clitopilus</i>	0	0	0	0.062	0	0.062	Basidiomycota	Agaricomycetes	Agaricales	Entolomataceae	LHC	saprotroph
<i>Clonostachys</i>	0	0.073	0.073	0	0	0	Ascomycota	Sordariomycetes	Hypocreales	Bionectriaceae	HC	parasitic or mutualistic symbiont of plants
<i>Collybia</i>	0	0	0	0	0.074	0.074	Basidiomycota	Agaricomycetes	Agaricales	Tricholomataceae	LHC	saprotroph
<i>Cordana</i>	0	0.073	0.073	0	0	0	Ascomycota	norank	norank	norank	HC	saprotroph
<i>Cordyceps</i>	0	0.073	0.073	0	0.074	0.074	Ascomycota	Sordariomycetes	Hypocreales	Cordycipitaceae	HC	parasitic or mutualistic symbiont of animals
<i>Cryptococcus</i>	0.12	1.015	1.135	0	0.739	0.739	Basidiomycota	Tremellomycetes	Tremellales	Tremellaceae	HC	parasitic or mutualistic symbiont of animals
<i>Curvularia</i>	0	0.073	0.073	0	0	0	Ascomycota	Dothideomycetes	Pleosporales	Pleosporaceae	HC	parasitic or mutualistic symbiont of animals
<i>Cyphellophora</i>	0.12	0.218	0.338	0.186	0	0.186	Ascomycota	Eurotiomycetes	Chaetothyriales	Chaetothyriaceae	HC	parasitic or mutualistic symbiont of animals

...Continued on next page

TABLE 2. (Continued)

Genus	RSDP		SCSP		Phylum	Class	Order	Family	Biochemical Activity ¹	Main Ecological Group		
	ITS1	ITS2	Total	ITS1							ITS2	Total
<i>Dactyliella</i>	0	0.145	0.145	0	0.443	0.443	Ascomycota	Orbiliiales	Orbiliaceae	HC	parasitic or mutualistic symbiont of animals	
<i>Debaryomyces</i>	0	0.218	0.218	0	0.074	0.074	Ascomycota	Saccharomycetes	Saccharomycetales	Saccharomycetaceae	HC	saprotroph
<i>Devriesta</i>	0	0	0	0	0.074	0.074	Ascomycota	Dothideomycetes	Capnodiales	Teratosphaeriaceae	HC	saprotroph
<i>Diplodia</i>	0	0	0	0	0.148	0.148	Ascomycota	Dothideomycetes	Botryosphaeriales	Botryosphaeriaceae	HC	parasitic or mutualistic symbiont of plants
<i>Epicoccum</i>	0.241	0	0.241	0	0.222	0.222	Ascomycota	Dothideomycetes	Pleosporales	Pleosporaceae	HC	parasitic or mutualistic symbiont of plants
<i>Exophiala</i>	0	0	0	0.248	0	0.248	Ascomycota	Eurotiomycetes	Chaetothyriales	Herpotrichiellaceae	HC	parasitic or mutualistic symbiont of plants
<i>Fimetariella</i>	0	0.145	0.145	0	0	0	Ascomycota	Sordariomycetes	Sordariales	Lasiosphaeriaceae	HC	saprotroph
<i>Flavodon</i>	0.361	0.29	0.651	2.664	4.065	6.729	Basidiomycota	Agaricomycetes	Polyporales	Meruliaceae	LHC	saprotroph
<i>Fomitopsis</i>	0	0	0	0.062	0	0.062	Basidiomycota	Agaricomycetes	Polyporales	Fomitopsidaceae	HC	saprotroph
<i>Fonsecaea</i>	0.361	0	0.361	0.124	0.517	0.641	Ascomycota	Eurotiomycetes	Chaetothyriales	Herpotrichiellaceae	HC	parasitic or mutualistic symbiont of animals
<i>Fusarium</i>	0.482	0.145	0.627	0.248	0.443	0.691	Ascomycota	Sordariomycetes	Hypocreales	Nectriaceae	HC	parasitic or mutualistic symbiont of plants

...Continued on next page

TABLE 2. (Continued)

Genus	RSDP			SCSP			Phylum	Class	Order	Family	Biochemical Activity ¹	Main Ecological Group
	ITS1	ITS2	Total	ITS1	ITS2	Total						
<i>Galerina</i>	0.482	0.073	0.555	0.434	0.074	0.508	Basidiomycota	Agaricomycetes	Agaricales	Strophariaceae	LHC	saprotroph
<i>Ganoderma</i>	0.602	0	0.602	0.062	0.074	0.136	Basidiomycota	Agaricomycetes	Polyporales	Ganodermataceae	LHC	saprotroph
<i>Gerronema</i>	0	0	0	0.31	0	0.31	Basidiomycota	Agaricomycetes	Agaricales	Marasmiaceae	LHC	saprotroph
<i>Gliocladium</i>	0.241	0	0.241	0	0	0	Ascomycota	Sordariomycetes	Hypocreales	Hypocreaceae	HC	saprotroph
<i>Gloeotinia</i>	0.241	0	0.241	0	0	0	Ascomycota	Leotiomycetes	Helotiales	norank	HC	parasitic or mutualistic
<i>Grammothele</i>	0.241	0	0.241	0	0	0	Basidiomycota	Agaricomycetes	Polyporales	Polyporaceae	LHC	saprotroph
<i>Greeneria</i>	0	0	0	0	0.148	0.148	Ascomycota	Sordariomycetes	Diaporthales	Gnomoniaceae	HC	saprotroph
<i>Gymnopus</i>	0	0	0	1.425	0.296	1.721	Basidiomycota	Agaricomycetes	Agaricales	Omphalotaceae	LHC	saprotroph
<i>Hansfordia</i>	0	0.218	0.218	0	0	0	Ascomycota	Sordariomycetes	Xylariales	norank	HC	saprotroph
<i>Henningsomyces</i>	0	0	0	0.062	0	0.062	Basidiomycota	Agaricomycetes	Agaricales	Schizophyllaceae	LHC	saprotroph
<i>Hyphoderma</i>	0	0	0	0.124	0.148	0.272	Basidiomycota	Agaricomycetes	Polyporales	Meruliaceae	LHC	saprotroph
<i>Hypocrea</i>	0	0.218	0.218	0.372	0.074	0.446	Ascomycota	Sordariomycetes	Hypocreales	Hypocreaceae	HC	saprotroph
<i>Isaria</i>	0.12	0.145	0.265	0	0.296	0.296	Ascomycota	Sordariomycetes	Hypocreales	Cordycipitiaceae	HC	parasitic or mutualistic
<i>Kuraishia</i>	0	0	0	0	0.074	0.074	Ascomycota	Saccharomycetes	Saccharomycetales	Saccharomycetaceae	HC	saprotroph
<i>Lasiodiplodia</i>	0.723	0.218	0.941	0.186	0.517	0.703	Ascomycota	Dothideomycetes	Botryosphaeriales	Botryosphaeriaceae	HC	parasitic or mutualistic
<i>Lasionectria</i>	0	0	0	0	0.074	0.074	Ascomycota	Sordariomycetes	Hypocreales	Bionectriaceae	HC	saprotroph

...Continued on next page

TABLE 2. (Continued)

Genus	RSDP		SCSP		Phylum	Class	Order	Family	Biochemical Activity ¹	Main Ecological Group		
	ITS1	ITS2	Total	ITS1							ITS2	Total
<i>Lecanicillium</i>	0	0.073	0.073	0	0	0	Ascomycota	Sordariomycetes	Hypocreales	Cordycipitaceae	HC	parasitic or mutualistic symbiont of animals
<i>Lepiota</i>	0	0	0	0.062	0	0.062	Basidiomycota	Agaricomycetes	Agaricales	Agaricaceae	LHC	saprotroph
<i>Letendraea</i>	0	0.073	0.073	0	0	0	Ascomycota	Dothideomycetes	Pleosporales	Tubeufiaceae	HC	parasitic or mutualistic symbiont of plants
<i>Leucocoprinus</i>	0	0	0	0.062	0.074	0.136	Basidiomycota	Agaricomycetes	Agaricales	Agaricaceae	LHC	saprotroph
<i>Malassezia</i>	0	0	0	0.062	0.517	0.579	Basidiomycota	Exobasidiomycetes	Malasseziales	Malasseziaceae	HC	parasitic or mutualistic symbiont of animals
<i>Megacollybia</i>	0	0	0	0	0.222	0.222	Basidiomycota	Agaricomycetes	Agaricales	Marasmiaceae	LHC	saprotroph
<i>Meira</i>	0	0	0	0.062	0.074	0.136	Basidiomycota	Exobasidiomycetes	Exobasidiales	Brachybasidiaceae	HC	parasitic or mutualistic symbiont of plants
<i>Metacorytheopsis</i>	0	0.073	0.073	0	0	0	Ascomycota	Sordariomycetes	Hypocreales	Clavicipitaceae	HC	parasitic or mutualistic symbiont of animals
<i>Metarhizium</i>	0	0.29	0.29	0	0.148	0.148	Ascomycota	Sordariomycetes	Hypocreales	Clavicipitaceae	HC	parasitic or mutualistic symbiont of animals

...Continued on next page

TABLE 2. (Continued)

Genus	RSDP		SCSP		Phylum	Class	Order	Family	Biochemical Activity ¹	Main Ecological Group		
	ITS1	ITS2	Total	ITS1							ITS2	Total
<i>Metulocladosporiella</i>	0	0.87	0.87	0	2.513	2.513	Ascomycota	Dothideomycetes	Capnodiales	Mycosphaerellaceae	HC	parasitic or mutualistic symbiont of plants
<i>Meyerozyma</i>	52.771	23.06	75.831	47.77	18.256	66.026	Ascomycota	Saccharomycetes	Saccharomycetales	Debaryomycetaceae	HC	saprotroph
<i>Monascus</i>	0.12	0	0.12	0	0	0	Ascomycota	Eurotiomycetes	norank	Monascaceae	HC	saprotroph
<i>Mortierella</i>	0	0	0	0	0.074	0.074	Zygomycota	norank	Mortierellales	Mortierellaceae	S	saprotroph
<i>Munkovalsaria</i>	0	0	0	0	0.074	0.074	Ascomycota	Dothideomycetes	Pleosporales	Dacampiaceae	HC	saprotroph
<i>Mycosphaerella</i>	0	0.218	0.218	0	0	0	Ascomycota	Dothideomycetes	Capnodiales	Mycosphaerellaceae	HC	parasitic or mutualistic symbiont of plants
<i>Myrothecium</i>	0	0	0	0.062	0	0.062	Ascomycota	Sordariomycetes	Hypocreales	norank	HC	saprotroph
<i>Nigrospora</i>	0	0.073	0.073	0.062	0.37	0.432	Ascomycota	Sordariomycetes	Trichosphaeriales	norank	HC	saprotroph
<i>Ochroconis</i>	0	0	0	0	0.074	0.074	Ascomycota	norank	norank	norank	HC	parasitic or mutualistic symbiont of animals
<i>Paraconiothyrium</i>	0	0.363	0.363	0	0.148	0.148	Ascomycota	Dothideomycetes	Pleosporales	Montagnulaceae	HC	saprotroph
<i>Paradictyoarhrium</i>	0	0.073	0.073	0	0	0	Ascomycota	norank	norank	norank	HC	saprotroph
<i>Penicillium</i>	0	0.145	0.145	0.372	0.887	1.259	Ascomycota	Eurotiomycetes	Eurotiales	Trichocomaceae	HC	saprotroph
<i>Perenniporia</i>	0	0.073	0.073	0	0	0	Basidiomycota	Agaricomycetes	Polyporales	Polyporaceae	LHC	saprotroph
<i>Pestalotiopsis</i>	0	0.363	0.363	0	0.074	0.074	Ascomycota	Sordariomycetes	Xylariales	Amphisphaeriaceae	HC	saprotroph

...Continued on next page

TABLE 2. (Continued)

Genus	RSDP				SCSP			Phylum	Class	Order	Family	Biochemical Activity ¹	Main Ecological Group	
	ITS1		ITS2		Total	ITS1	ITS2							Total
	ITS1	ITS2	ITS1	ITS2										
<i>Phanerochaete</i>	0	0	0	0	0	0.31	0.148	0.458	Basidiomycota	Agaricomycetes	Polyporales	Phanerochaetaceae	LHC	saprotroph
<i>Phialemonium</i>	0	0	0	0	0	0.062	0.222	0.284	Ascomycota	Sordariomycetes	Sordariales	Cephalothecaceae	HC	parasitic or mutualistic symbiont of animals
<i>Phialophora</i>	0.723	0.218	0.941	0.496	0	0.496	0	0.496	Ascomycota	Eurotiomycetes	Chaetothyriales	Herpotrichiellaceae	HC	parasitic or mutualistic symbiont of animals
<i>Phlebia</i>	0.241	0.073	0.314	0	0	0	0	0	Basidiomycota	Agaricomycetes	Polyporales	Meruliaceae	LHC	saprotroph
<i>Phoma</i>	0	0.073	0.073	0	0	0.074	0.074	0.074	Ascomycota	Dothideomycetes	Pleosporales	norank	HC	parasitic or mutualistic symbiont of plants
<i>Pichia</i>	2.41	5.729	8.139	6.258	15.743	22.001	22.001	22.001	Ascomycota	Saccharomycetes	Saccharomycetales	Saccharomycetaceae	HC	saprotroph
<i>Pluteus</i>	0	0	0	0.31	0	0.31	0	0.31	Basidiomycota	Agaricomycetes	Agaricales	Pluteaceae	LHC	saprotroph
<i>Prosthecium</i>	0	0.145	0.145	0	0	0	0	0	Ascomycota	Sordariomycetes	Diaporthales	Melanconidaceae	HC	parasitic or mutualistic symbiont of plants
<i>Pseudocercospora</i>	0	0.073	0.073	0	0	0	0	0	Ascomycota	Dothideomycetes	Capnodiales	Mycosphaerellaceae	HC	parasitic or mutualistic symbiont of plants

...Continued on next page

TABLE 2. (Continued)

Genus	RSDP			SCSP			Phylum	Class	Order	Family	Biochemical Activity ¹	Main Ecological Group
	ITS1	ITS2	Total	ITS1	ITS2	Total						
<i>Ramichloridium</i>	0	0.073	0.073	0	0	0	Ascomycota	Dothideomycetes	Capnodiales	Mycosphaerellaceae	HC	saprotroph
<i>Resinicium</i>	24.578	11.965	36.543	10.037	6.356	16.393	Basidiomycota	Agaricomycetes	Hymenochaetales	Rickenellaceae	LHC	saprotroph
<i>Resupinatus</i>	0	0.145	0.145	1.363	0.148	1.511	Basidiomycota	Agaricomycetes	Agaricales	Tricholomataceae	LHC	saprotroph
<i>Rhinocladiella</i>	0.12	0.29	0.41	0.062	0.37	0.432	Ascomycota	Eurotiomycetes	Chaetothyriales	Herpotrichiellaceae	HC	parasitic or mutualistic symbiont of animals
<i>Rhodotorula</i>	0.602	0	0.602	0.434	0.813	1.247	Basidiomycota	Microbotryomycetes	Sporidiobolales	norank	HC	saprotroph
<i>Scheffersomyces</i>	0.12	0	0.12	0	0	0	Ascomycota	Saccharomycetes	Saccharomycetales	Debaryomycetaceae	HC	saprotroph
<i>Scytalidium</i>	1.687	35.968	37.655	0.991	15.225	16.216	Ascomycota	Leotiomycetes	Helotiales	norank	HC	saprotroph
<i>Seimatosporium</i>	0.12	0	0.12	0	0	0	Ascomycota	Sordariomycetes	Xylariales	Amphisphaeriaceae	HC	saprotroph
<i>Simplicillium</i>	1.325	0	1.325	0.496	0	0.496	Ascomycota	Sordariomycetes	Hypocreales	Cordycipitaceae	HC	saprotroph
<i>Sporothrix</i>	0	0.29	0.29	0.062	0.296	0.358	Ascomycota	Sordariomycetes	Ophiostomatales	Ophiostomataceae	HC	saprotroph
<i>Sugiyamaella</i>	0.12	0.073	0.193	0.186	0	0.186	Ascomycota	Saccharomycetes	Saccharomycetales	Trichomonascaceae	HC	saprotroph
<i>Symbiotaphrina</i>	0	0.073	0.073	0	0	0	Ascomycota	norank	norank	norank	HC	parasitic or mutualistic symbiont of animals
<i>Teratosphaeria</i>	0	0	0	0	0.148	0.148	Ascomycota	Dothideomycetes	Pleosporales	Pleosporaceae	HC	parasitic or mutualistic symbiont of plants

... Continued on next page

TABLE 2. (Continued)

Genus	RSDP		SCSP		Phylum	Class	Order	Family	Biochemical Activity ¹	Main Ecological Group	
	ITS1	ITS2	Total	ITS1							ITS2
<i>Thielavia</i>	0.12	0	0.12	0	0	Ascomycota	Sordariomycetes	Sordariales	Chaetomiaceae	HC	saprotroph
<i>Torulaspota</i>	0	0	0	0	0.074	Ascomycota	Saccharomycetes	Saccharomycetales	Saccharomycetaceae	HC	saprotroph
<i>Trametes</i>	0.241	1.668	1.909	0	0.665	Basidiomycota	Agaricomycetes	Polyporales	Polyporaceae	LHC	saprotroph
<i>Trichoderma</i>	0.12	0.508	0.628	0.434	0.961	Ascomycota	Sordariomycetes	Hypocreales	Hypocreaceae	HC	saprotroph
<i>Tricladium</i>	0.12	0	0.12	0	0	Ascomycota	Leotiomycetes	Helotiales	Helotiaceae	HC	saprotroph
<i>Umberlopsis</i>	0	0	0	0.062	0.284	Zygomycota	norank	Mucorales	Umberlopsidaceae	S	saprotroph
<i>Veronaea</i>	0	0.073	0.073	0.124	0	Ascomycota	norank	norank	norank	HC	parasitic or mutualistic
<i>Verticicola</i>	1.325	6.889	8.214	0.929	6.578	Ascomycota	Sordariomycetes	norank	Annulatascaceae	HC	saprotroph
<i>Verticillium</i>	0	0	0	0.062	0.222	Ascomycota	Sordariomycetes	norank	Plectosphaerellaceae	HC	parasitic or mutualistic
<i>Villosiclava</i>	0.361	0	0.361	0.31	0	Ascomycota	Sordariomycetes	Hypocreales	Clavicipitaceae	HC	symbiont of plants
<i>Wickerhamomyces</i>	0	0.073	0.073	0	0	Ascomycota	Saccharomycetes	Saccharomycetales	Wickerhamomycetaceae	HC	saprotroph
<i>Xylaria</i>	0	0	0	0.124	0.074	Ascomycota	Sordariomycetes	Xylariales	Xylariaceae	HC	saprotroph

¹Note: LHC: able to degrade lignin, cellulose and hemicelluloses (white-rot) in bold; HC: able to degrade hemicelluloses and/or cellulose

Discussion

The internal transcribed spacer (ITS1 and ITS2) and 5.8S regions of the nuclear ribosomal repeat unit have been used as the primary fungal barcode markers for species delimitation (Schoch *et al.* 2012). The full-length ITS region comprises approximately 599 ± 113 bp; the ITS1 region contributes approximately 208 ± 57 and ITS2, 310 ± 67 (Porrás-Alfaro *et al.* 2014). “Ion torrent 316” generates reads of approximately 200 bp in length; thus, it is important to determine the best metabarcodes for fungal metagenomics research. The use of ITS1 and/or ITS2 in the identification of fungal species has been previously evaluated but there is no consensus as to which of them, ITS1 (Wang *et al.* 2015), ITS2 (Bazzicalupo *et al.* 2013) or both (Porrás-Alfaro *et al.* 2014) capture the highest number of fungal taxa. In our study, the full-length ITS region was amplified and submitted to NGS, randomly generating sequences. The sequence data from ITS1 and ITS2 were treated separately, and the obtained read lengths were sufficient to discriminate MOTUs at the genus level at 95% similarity level.

In this study we identified a higher number of genera in SCSP than in RSDP. The first area is located in a region known for a high level of species richness and a high degree of endemism of trees and did not show evidence of recent disturbance, representing the best state of conservation of the original vegetation (Martini *et al.* 2007). By contrast, RSDP showed lower species richness of trees (Lopes *et al.* 2002) and showed several signs of anthropogenic impacts. According to Boddy (2001), a higher species diversity of trees represents increased variability in the amount of woody debris available for decomposition by saproxylic fungi.

The majority of the ascomycotan genera retrieved in our study are directly or indirectly related to the decomposition of plant debris (Zhao *et al.* 2013), and Sordariomycetes exhibited the highest number of genera, as also reported in a global biodiversity study of soil mycobiome (Tedersoo *et al.* 2014). We identified four genera (*Pestalotiopsis*, *Xylaria*, *Hansfordia*, *Seimatosporium*) of Xylariales that comprise the most effective group of wood decomposers in the Sordariomycetes (Pointing *et al.* 2003, Eichlerová *et al.* 2015). Furthermore, many of the ascomycotan saproxylic species belonging to the Hypocreales (*Trichoderma*, *Acremonium*, *Fusarium*, *Simplicillium*, *Hypocrea*, *Villosiclava*, *Gliocladium*, *Myrothecium*, *Chaunopycnis*, *Clonostachys*, *Lasionectria*), Eurotiales (*Aspergillus* and *Penicillium*), Diaporthales (*Greeneria* and *Prosthecia*), Chaetosphaeriales (*Chloridium*) and the Sordariales (*Chaetosphaeria*, *Phiallemonium*, *Fimetariella*, *Thielavia*) were also retrieved. The genus *Sporothrix* (Ophiostomatales), which has been associated with bark and ambrosia beetles, as a causal agent of sap stain in freshly cut wood, and also as a ubiquitous saprobe on rotting leaves, wood, soil (Meyer *et al.* 2008) was also detected. In contrast to the Hypocreales, the majority of which are related to wood decomposition, the genera *Isaria*, *Beauveria*, *Cordyceps*, and *Lecanicillium* are composed of typical insect pathogens (Cannon & Kirk 2007). Moreover, several genera of Eurotiomycetes (*Phialophora*, *Fonsecaea*, *Rhinochrysiella*, *Exophiala*, *Cyphellophora*, and *Cladophialophora*) can occur as saprobes or facultative pathogens of animals that cause opportunistic infections (Cannon & Kirk 2007).

Saccharomycetes are mainly related to sugar decomposition and are present in live and dead plant tissues, including wood (Stokland *et al.* 2012). Furthermore, Saccharomycetes are especially associated with the guts of saproxylic insects involved in lignocellulosic decomposition (Calderon & Berkov 2012). The majority of the genera of this class (*Pichia*, *Candida*, *Sugiyamaella*, *Debaryomyces*, *Kuraishia*, *Scheffersomyces*, *Wickerhamomyces*, *Torulaspora*, and *Meyerozyma*) have also been detected in previous studies of wood-decaying fungi from tropical forests that employed culture-dependent methods (Urbina *et al.* 2013, Davis 2014). Orbiliomycetes (*Arthrobotryx*, *Dactylella*) are typically found as saprotrophs on wood and can also present a nematophagous habit (Cannon & Kirk 2007), which represents a strategy for survival in low-nitrogen environments such as wood (Berg & McClaugherty 2008).

There were proportionally many more genera of white-rot fungi than brown-rot fungi as expected for wood-decaying Basidiomycota in tropical biomes (Ryvarden 1991) (in our study approx. 87:13%). With a few exceptions, the majority of the retrieved genera belonged to Agaricomycetes, a pattern similar to that reported in Tedersoo *et al.* (2014) for soil fungi. This basidiomycotan class includes saprotrophs, pathogens and mutualists, and all of the species are capable of lignin decomposition (Riley *et al.* 2014). Numerous representatives of white-rot genera of Hymenochaetales (*Resinicium*), Polyporales (*Flavodon*, *Ganoderma*, *Phanerochaete*, *Trametes*, *Hyphoderma*, *Phlebia*, *Perenniporia*, *Grammothele*), Russulales (*Asterostoma*) and Agaricales (*Pluteus*, *Resupinatus*, *Gymnopus*, *Leucocoprinus*, *Galerina*, *Lepiota*, *Gerronema*, *Megacollybia*, *Collybia*, *Clitopilus*, *Agaricus*) were detected. Furthermore, brown-rot genera of Polyporales such as *Fomitopsis* (Yoon *et al.* 2005) and *Antrodia* (Clausen & Green 2003) were also present, as well as typical brown-rot taxa of Dacrymycetes (*Calocera*) (Seifert, 1983) and hemicellulolytic taxa associated with saproxylic insects in the order Tremellales (*Cryptococcus*) (Urbina *et al.* 2013). Moreover, two genera (*Meira* and *Malassezia*) belonging to the subphylum Ustilaginomycotina, which are not directly related to wood decomposition,

were also retrieved. *Meira* is typically reported as an endophyte associated with mites (Rush & Aime 2013), and it has been detected in orchid roots (Huang *et al.* 2014). *Malassezia* has been detected in the gut of basidiomata-feeding beetles (Blackwell *et al.* 2007), in addition to occurring in the skin of animals as a commensal and facultative pathogen (Wang *et al.* 2014).

A total of 20% of the genera identified was unequivocally composed of lignin decomposers as the species of these genera possess high-oxidation potential peroxidases, such as LiP, MnP, and VP, based on genetic and/or biochemical evidence according to the CAZY (Lombard *et al.* 2013) and BRENDA (Schomburg *et al.* 2013) databases, as well as specialized literature. Moreover, the majority of the genera presented evidence of bearing hemicellulolytic and/or cellulolytic enzymes or causing brown and soft rot in wood, excluding some genera of typical “sugar fungi”, e.g., *Mortierella* and *Umbelopsis*.

Nine studies have already specifically investigated the mycobiomes of decaying wood using targeted, amplicon-based metagenomics, five in palearctic boreal forests in northern Europe (Ovaskainen *et al.* 2010, Kubartova *et al.* 2012, Ovaskainen *et al.* 2013, Runnel *et al.* 2015) and four in palearctic temperate forests in Europe and Asia (Van der Wal *et al.* 2015, Hoppe *et al.* 2015, Jang *et al.* 2015, Yamashita *et al.* 2015). With the exception of Yamashita *et al.* (2015), which used the V9 region of 18S rDNA, all the other works used the complete ITS (Ovaskainen *et al.* 2010, Kubartova *et al.* 2012, Ovaskainen *et al.* 2013, Hoppe *et al.* 2015, Otosson *et al.* 2015) or the ITS2 (Jang *et al.* 2015, Runnel *et al.* 2015, Van der Wal *et al.* 2015) as the targeted region. Furthermore, all the analyses of these studies were mainly focused on Basidiomycota, limiting a fully global comparison with our study. Nevertheless, some general patterns could be evidenced taking into account the Phylum Basidiomycota. Some genera were retrieved in all forest types (boreal, temperate and tropical) such as *Resinicium*, *Pluteus*, *Ganoderma*, *Calocera*, *Cryptococcus*, *Phanerochaete*, *Galerina*, *Phlebia*, *Megacollybia*, *Fomitopsis* and *Antrodia*, while other genera, such as *Asterostroma*, *Flavodon*, *Resupinatus*, *Gymnopus*, *Leucocoprinus*, *Lepiota*, *Meira*, *Gerronema*, *Collybia*, *Grammothele* and *Agaricus* were only identified in our study in the tropical sites and were not reported in any of the other aforementioned studies based on large-scale DNA sequencing in boreal or temperate forests. The fungal genera identified in all forest types have a wide geographic distribution, occurring in all major climatic zones (Ryvarden 1993), whereas some of the genera detected only in the neotropical forests usually have known geographic distribution restricted to or mainly in tropical and subtropical regions (Suhara *et al.* 2010, Velinga 2004, Johnson & Vilgalys 1998).

There are at least twice as many Ascomycota than Basidiomycota taxa (Hibbett, 2014), and this pattern was also retrieved in our work. In a recent and comprehensive review of the diversity of Brazilian fungi, Maia *et al.* (2015) reported that almost half of the fungal taxa recorded in Brazil was Basidiomycota (47.9%), followed by Ascomycota (32.9%). The same authors suggested that this inverted ratio could be explained by the low number of experts in Ascomycota taxonomy in Brazil.

The number of fungal genera retrieved in our work was little more than half of the fungal genera presented in the continuously updated list of Plants and Fungi of Brazil (Canhos *et al.* 2015) when we filtered this general database for the parameters: Atlantic forest biome and dead wood. Furthermore, approximately 77% of the fungal genera retrieved in our work were not recorded in this trimmed dataset. Nevertheless, some of them are yeasts or microfungi that could not be represented in this database.

Conclusion

This is the first study using NGS to uncover the diversity of wood-decaying fungi in tropical forests. The higher diversity of trees and lower disturbance in the tropical rainforest site could explain the higher number of fungal genera obtained in the tropical rainforest site when compared with the seasonal tropical forest. All genera belonging to the ecological functional groups of saproxylic fungi, except for mycorrhizal fungi, were retrieved. Most prevalent genera were ligninolytic and/or cellulolytic Ascomycota (Sordariomycetes) and Basidiomycota (Agaricomycetes) but plant pathogens, endophytes and facultative or obligatory animal pathogenic fungi were also found. Moreover, the great majority of Basidiomycota genera are white-rotters as expected for tropical regions. As the metabarcoding approach is not dependent on collection, direct observation and/or culturing, this methodology theoretically permits the most thorough access to the diversity of the mycobiome of any kind of substrate. Thus, the metabarcoding as a rapid, large-scale, and theoretically taxonomically unbiased strategy, proved useful to perform comprehensive surveys, however, the clustering and taxonomic identification steps are still challenging. We recommend to test different percentage values for these steps and compare the results with a comprehensive and reliable database.

Acknowledgments

We thank all individuals who contributed directly or indirectly to this work, especially the authorities of RDSP (Rio Doce State Park) and SCSP (Serra do Conduru State Park), CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico), FIOCRUZ-MG (Fundação Oswaldo Cruz, Minas Gerais) and the Graduate Programs of Microbiology (PPG Microbiologia) and Bioinformatics (PPG Bioinformatics) of the Federal University of Minas Gerais (UFMG). A. Góes-Neto is supported by CNPq (contract n. 308148/2013-4). Our research group is part of the Brazilian Microbiome Project (BMP): <http://www.brmicrobiome.org/>.

References

- Bazzicalupo, A.L., Bálint, M. & Schmitt, I. (2013) Comparison of ITS1 and ITS2 rDNA in 454 sequencing of hyperdiverse fungal communities. *Fungal Ecology* 6: 102–109.
- Berg, B. & McLaugherty, C. (2008) Decomposition of fine root and woody litter. In: Berg, B. & McLaugherty, C. (Eds.) *Plant litter: Decomposition, Humus formation, carbon sequestration*. Springer, Berlin, pp. 171–199.
- Blaalid, R., Kumar, S., Nilsson, R.H., Abarenkov, K., Kirk, P.M. & Kauserud, H. (2013) ITS1 versus ITS2 as DNA metabarcodes for fungi. *Molecular Ecology Resources* 13: 218–224.
- Blackwell, M., Suh, S.O. & Nardi, J.B. (2007) Fungi in the hidden environment: the gut of beetles. In: Gadd, G.M., Watkinson, S.C. & Dyer, P.S. (Eds.) *Fungi in the Environment*. British Mycological Society Symposia, Cambridge, pp. 357–370.
- Boddy, L. (2001) Fungal community ecology and wood decomposition processes in angiosperms: from standing trees to complete decays coarse woody debris. *Ecological Bulletins* 49: 43–56.
- Calderon, O. & Berkov, A. (2012) Midgut and fat body bacteriocytes in neotropical cerambycid beetles (Coleoptera: Cerambycidae). *Environmental Entomology* 41: 108–117.
- Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K. & Madden, T.L. (2008) BLAST+: architecture and applications. *BMC Bioinformatics* 10: 421.
- Cannon, P. & Sutton, B. (2004) Microfungi on wood and plant debris. In: Mueller, G., Bills, G. & Foster, M. (Eds.) *Biodiversity of fungi: inventory and monitoring methods*. Elsevier Academic Press, Burlington, pp. 217–239.
- Cannon, P.F. & Kirk, P.M. (2007) *Fungal families of the world*. CABI, United Kingdom, 456 pp.
- Canhos, D.A.L., Sousa-Baena, M.S., Souza, S. de, Maia, L.C., Stehmann, J.R., Canhos, V.P., Giovanni, R. De., Bonacelli, M.B.M., Los, W. & Peterson, A.T. (2015) The importance of biodiversity E-infrastructures for megadiverse countries. *PLoS Biology* 13.7: e1002204.
- Clausen, C.A. & Green, F. (2003) Oxalic acid overproduction by copper-tolerant brown-rot basidiomycetes on southern yellow pine treated with copper-based preservatives. *International Biodeterioration & Biodegradation* 51 (2): 139–144.
- Cuadros-Orellana, S., Leite, L.R., Smith, A., Medeiros, J.D., Badotti, F., Fonseca, P.L.C., Vaz, A.B.M., Oliveira, G. & Góes-Neto, A. (2013) Assessment of Fungal Diversity in the Environment using Metagenomics: a Decade in Review. *Fungal Genomics & Biology* 3: 110.
- Davis, T.S. (2014) The ecology of yeasts in the bark beetle holobiont: A century of research revisited. *Microbial Ecology* 69: 723.
- Edgar, R.C. (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26: 2460–2461.
- Eichlerová, I., Homolka, L., Žifčáková, L., Lisá, L., Dobiášová, P. & Baldrian, P. (2015) Enzymatic systems involved in decomposition reflects the ecology and taxonomy of saprotrophic fungi. *Fungal Ecology* 13: 10–22.
- Evert, R.F. & Eichorn, S.E. (2013) *Raven Biology of Plants*. W.H. Freeman and Company, New York, USA, 880 pp.
- Goodwin, S., McPherson, J.D. & McCombie, W.R. (2016) Coming of age: ten years of next-generation sequencing technologies. *Nature Reviews Genetics* 17: 333–351.
- Góes-Neto, A., Loguerio-Leite, C. & Guerrero, R.T. (2005) DNA extraction from frozen field collected and dehydrated herbarium fungal basidiomata: performance of SDS and CTAB-based methods. *Biotemas* 18: 19–32.
- Hatakka, A. (2001) Biodegradation of lignin. In: Hofrichter, M. & Steinbüchel, A. (Eds.) *Lignin, Humic Substances and Coal (Biopolymers Vol. 1.)*. Wiley-VCH, Weinheim, Germany, pp. 129–180.
- Hibbett, D.S. (2014) Major events in the evolution of the Fungi. In: Losos, J. (Ed.) *Princeton Guide to Evolution*. Princeton University Press.
- Hoppe, B., Purahong, W., Wubet, T., Kahl, T., Bauhus, J., Arnstadt, T., Hofrichter, M., Buscot, F. & Krüger, D. (2015) Linking molecular deadwood-inhabiting fungal diversity and community dynamics to ecosystem functions and processes in Central European forests. *Fungal Diversity* 77: 1–13.

- Huang, C.L., Jian, F.Y., Huang, H.J., Chang, W.C., Wu, W.L., Hwang, C.C., Lee, R.H. & Chiang, T.Y. (2014) Deciphering mycorrhizal fungi in cultivated *Phanaeopsis* microbiome with next-generation sequencing of multiple barcodes. *Fungal Diversity* 66: 77–88.
- Jang, Y., Jang, S., Min, M., Hong, J.H., Lee, H., Lee, H., Lim, Y.W. & Kim, J.J. (2015) Comparison of the Diversity of Basidiomycetes from Dead Wood of the Manchurian fir (*Abies holophylla*) as Evaluated by Fruiting Body Collection, Mycelial Isolation, and 454 Sequencing. *Microbial ecology* 70 (3): 634–645.
- Johnson, J. & Vilgalys, R. (1998) Phylogenetic systematics of *Lepiota* sensu lato based on nuclear large subunit rDNA evidence. *Mycologia*: 971–979.
- Koljalg, U., Nilsson, R.H., Abarenkov, K., Tedersoo, L., Taylor, A.F.S., Bahram, M., Bates, S.T., Bruns, T.D., Bengtsson-Palme, J., Callaghan, T.M., Douglas, B., Drenkhan, T., Eberhardt, U., Dueñas, M., Grebenc, T., Griffith, Hartmann, M., Kirk, P.M., Kohout, P., Larsson, E., Lindahl, B.D., Lücking, R., Martín, M.P., Matheny, P.B., Hguyen, N.H., Niskanen, T., Oja, J., Peay, K.G., Peintner, U., Peterson, M., Pöldmaa, K., Saag, L., Saar, I., Schübler, A., Scott, J.A., Senés, C., Smith, M.E., Suija, A., Taylor, D.L., Telleria, M.T., Weiss, M. & Larsson, K.H. (2013) Towards a unified paradigm for sequence-based identification of fungi. *Molecular Ecology* 22: 5271–5277.
- Keller, M., Palace, M., Asner, G.P., Pereira, R. & Silva, J.N.M. (2004) Coarse woody debris in undisturbed and logged forests in the eastern Brazilian Amazon. *Global Change Biology* 10: 784–795.
- Kubartova, A., Ottosson, E., Dahlberg, A. & Stenlid, J. (2012) Patterns of fungal communities among and within decaying logs, revealed by 454 sequencing. *Molecular Ecology* 21: 4514–4532.
- Lindahl, B.D., Nilsson, R.H., Tedersoo, L., Abarenkov, K., Carlsen, T., Kjølner, R., Kõljalg, U., Pennanen, T., Rosendahl, S., Stenlid, J. & Kausrud, H. (2013). Fungal community analysis by high-throughput sequencing of amplified markers – a user’s guide. *New Phytologist* 199: 288–299.
- Lindner, D.L., Carlsen, T., Henrik, N.R., Davey, M., Schumacher, T. & Kausrud, H. (2013) Employing 454 amplicon pyrosequencing to reveal intragenomic divergence in the internal transcribed spacer rDNA region in fungi. *Ecology and Evolution* 3: 1751–1764.
- Lombard, V., Ramulu, H.G., Drula, E., Coutinho, P.M. & Henrissat, B. (2014) The carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic acids research* 42 (D1): D490–D495.
- Lopes, W. de P., da Silva, A.F., de Souza, A.L. & Neto, J.A.A.M. (2002) Estrutura fitossociológica de um trecho de vegetação arbórea no Parque Estadual do Rio Doce-Minas Gerais, Brasil. *Acta Botanica Brasilica* 16: 443–456.
- Maia, L.C., Carvalho Júnior, A.A.D., Cavalcanti, L.D.H., Gugliotta, A.D.M., Drechsler-Santos, E.R., Santiago, A.L.D.A., Cáceres, M.E.D.A., Gibertoni, T.B., Aptroot, A., Giachini, A.J., Soares, A.M.D.S., Silva, A.C.G., Magnago, A.C., Goto, B.T., Lira, C.R.S. de, Montoya, C.A.S., Pires-Zottarelli, C.L.A., Silva, D.K.A. de, Soares, D.J., Rezende, D.H.C., Luz, E.D.M.N., Gumboski, E.L., Wartchow, F., Karstedt, F., Freire, F.M., Coutinho, F.P., Melo, G.S.N. de, Sotão, H.M.P., Baseia, I.G., Pereira, J., Oliveira, J.J.S. de, Souza, J.F., Bezerra, J.L., Neta, L.S.A., Pfenning, L.H., Gusmão, L.F.P., Neves, M.A., Capelari, M., Jaeger, M.C.W., Pulgarín, M.P., Menolli, N. Jr., Medeiros, P.S. de, Friedrich, R.C.S., Chikowski, R. dos S., Pires, R.M., Melo, R.F., Silveira, R.M.B. da, Urrea-Valencia, S., Cortez, V.G. & Soares, A.M.D.S. (2015) Diversity of Brazilian Fungi. *Rodriguésia* 66 (4): 1033–1045.
- Martin, K.J. & Rygielwicz, P.T. (2005) Fungal-specific PCR primers developed for analysis of the ITS region of environmental DNA extracts. *BMC Microbiology* 5: 28.
- Martini, A.M.Z., Fiaschi, P., Amorim, A.M. & da Paixão, J.L. (2007) A hot-point within a hot-spot: a high diversity site in Brazil’s Atlantic Forest. *Biodiversity and Conservation* 16: 3111–3128.
- Meyer, E.M., De Beer, Z.W., Summerbell, R.C., Moharram, A.M., de Hoog, G.S., Vismar, H.F. & Wingfield, M.J. (2008) Taxonomy and phylogeny of new wood-and soil-inhabiting *Sporothrix* species in the *Ophiostoma stenoceras-Sporothrix schenckii* complex. *Mycologia* 100: 647–661.
- Morgenstern, I., Klopman, S. & Hibbett, D.S. (2008) Molecular evolution and diversity of lignin degrading heme Peroxidase in the Agaricomycetes. *Journal of Molecular Evolution* 66: 243–257.
- Nilsson, R.H., Kristiansson, E., Ryberg, M., Hallenberg, N. & Larsson, K.-H. (2008) Intraspecific ITS variability in the kingdom Fungi as expressed in the international sequence databases and its implications for molecular species identification. *Evolutionary Bioinformatics* 4: 193–201.
- Nilsson, R.H., Veldre, V., Hartmann, M., Unterseher, M., Amend, A., Bergsten, J., Kristiansson, E., Ryberg, M., Jumpponen, A. & Abarenkov, K. (2010) An open source software package for automated extraction of ITS1 and ITS2 from fungal ITS sequences for use in high-throughput community assays and molecular ecology. *Fungal Ecology* 3: 284–287.
- Ottosson, E., Kubartová, A., Edman, M., Jönsson, M., Lindhe, A., Stenlid, J., & Dahlberg, A. (2015) Diverse ecological roles within fungal communities in decomposing logs of *Picea abies*. *FEMS Microbiology Ecology* 91 (3).
- Ovaskainen, O., Nokso-Koivisto, J., Hottola, J., Rajala, T., Pennanen, T., Ali-Kovero, H., Miettinen, O., Oinonen, P., Auvinen, P., Paulin, L., Larsson, K.H. & Mäkipää, R. (2010) Identifying wood-inhabiting fungi with 454 sequencing—what is the probability that BLAST gives the correct species? *Fungal Ecology* 3: 274–283.
- Ovaskainen, O., Schigel, D., Ali-Kovero, H., Auvinen, P., Paulin, L., Nordén, B. & Nordén, J. (2013) Combining high-throughput

- sequencing with fruit body surveys reveals contrasting life-history strategies in fungi. *ISME Journal* 7: 1696–1709.
- Pointing, S.B., Parungao, M.M. & Hyde, K.D. (2003) Production of wood-decay enzymes, mass loss and lignin solubilization in wood by tropical *Xylariaceae*. *Mycological Research* 107: 231–235.
- Porras-Alfaro, A., Liu, K.L., Kuske, C.R. & Xie, G. (2014) From Genus to Phylum: Large-Subunit and Internal Transcribed Spacer rRNA Operon Regions Show Similar Classification Accuracies Influenced by Database Composition. *Applied Environmental Microbiology* 80: 829.
- Pylro, V.S. (2013) Brazilian Microbiome Project: Revealing the Unexplored Microbial Diversity—Challenges and Prospects. *Microbial Ecology*.
- R Development Core Team (2016) *R: a Language and Environment for Statistical Computing*. Available from: <http://www.rproject.org>. (accessed 10 July 2016)
- Ribeiro, M.C., Metzger, J.P., Martensen, A.C., Ponzoni, F.J. & Hirota, M.M. (2009) The Brazilian Atlantic Forest: How much is left, and how is the remaining forest distributed? Implications for conservation. *Biological Conservation* 142: 1141–1153.
- Riley, R., Salamov, A.A., Brown, D.W., Nagy, L.G., Floudas, D., Held, B.W., Levasseur, A., Lombard, V., Morin, E., Otilar, R., Lindquist, E.A., Sun, H., LaButti, K.M., Schmutz, J., Jabbour, D., Luo, H., Baker, S.E., Pisabarro, A.G., Walton, J.D., Blanchette, R.A., Henrissat, B., Martin, F., Cullen, D., Hibbett, D.S. & Grigoriev, I.V. (2014) Extensive sampling of basidiomycete genomes demonstrates inadequacy of the white-rot/brown-rot paradigm for wood decay fungi. *Proceedings of the National Academy of Sciences* 111: 9923–9928.
- Rothberg, J.M., Hinz, W., Rearick, T.M., Schultz, J., Mileski, W., Davey, M., Leamon, J.H., Johnson, K., Milgrew, M.J., Edwards, M., Hoon, J., Simons, J.F., Marran, D., Myers, J.W., Davidson, J.F., Branting, A., Nobile, J.R., Puc, B.P., Light, D., Clark, T.A., Huber, M., Branciforte, J.T., Stoner, I.B., Cawley, S.E., Lyons, M., Fu, Y., Homer, N., Sedova, M., Miao, X., Reed, B., Sabina, J., Feierstein, E., Schorn, M., Alanjary, M., Dimalanta, E., Dressman, D., Kasinskas, R., Sokolsky, T., Fidanza, J.A., Namsaraev, E., McKernan, K.J., Williams, A., Roth, G.T. & Bustillo, J. (2011) An integrated semiconductor device enabling non-optical genome sequencing. *Nature* 475: 348–352.
- Runnel, K., Tamm, H. & Lõhmus, A. (2015) Surveying wood-inhabiting fungi: most molecularly detected polypore species form fruit-bodies within short distances. *Fungal Ecology* 18: 93–99.
- Rush, T.A. & Aime, M.C. (2013) The genus *Meira*: phylogenetic placement and description of a new species. *Antonie van Leeuwenhoek* 103: 1097–1106.
- Ryvarden, L. (1991) *Genera of Polypores: Nomenclature and Taxonomy (Synopsis Fungorum Ser; n5)*. Lubrecht & Cramer Ltd, 373 pp.
- Ryvarden, L. (1993) Tropical polypores. In: British Mycological Society Symposium Series (Vol. 19). Cambridge University Press, pp. 149–149.
- Schwarze, F.W. (2007) Wood decay under the microscope. *Fungal Biology Reviews* 21: 133–170.
- Schwarze, F.W.M.R., Engels, J. & Mattheck, C. (2000) *Fungal strategies of wood decay in trees*. Springer, Berlin, 183 pp.
- Schoch, C.L., Seifert, K.A., Huhndorf, S., Robert, V., Spouge, J.L., Levesque, C.A., Chen, W. & Fungal Barcoding Consortium (2012) Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences USA* 109: 6241–6246.
- Schomburg, I., Chang, A., Placzek, S., Söhlgen, C., Rother, M., Lang, M. & Scheer, M. (2012) BRENDA in 2013: integrated reactions, kinetic data, enzyme function data, improved disease classification: new options and contents in BRENDA. *Nucleic acids research: gks1049*.
- Seifert, K.A. (1983) Decay of wood by the Dacrymycetales. *Mycologia* 75: 1011–1018.
- Stokland, J.N., Siitonen, J. & Jonsson, B.G. (2012) *Biodiversity in dead wood*. Cambridge University Press, London, 521 pp.
- Suhara, H., Maekawa, N., Ushijima, S., Kinjo, K. & Hoshi, Y. (2010) *Asterostroma* species (Basidiomycota) from mangrove forests in Japan. *Mycoscience* 51 (1): 75–80.
- Tedersoo, L., Bahram, M., Põlme, S., Kõljalg, U., Yorou, N.S., Wijesundera, R., Villarreal, R.L., Vasco-Palacios, A.M., Thu, P.Q., Suija, A., Smith, M.E., Sharp, C., Saluveer, E., Saitta, A., Rosas, M., Riit, T. & Abarenkov, K. (2014) Fungal biogeography. Global diversity and geography of soil fungi. *Science* 346: 1256688.
- Urbina, H., Schuster, J. & Blackwell, M. (2013) The gut of Guatemalan passalid beetles: a habitat colonized by cellobiose- and xylose-fermenting yeasts. *Fungal Ecology* 6: 339–355.
- Van der Wal, A., Ottosson, E. & de Boer, W. (2015) Neglected role of fungal community composition in explaining variation in wood decay rates. *Ecology* 96 (1): 124–133.
- Vellinga, E.C. (2004) Ecology and distribution of Lepiotaceous fungi (Agaricaceae)—A Review. *Nova Hedwigia* 78 (3–4): 273–299.
- Wang, Q.M., Theelen, B., Groenewald, M. & Bai, F.Y. (2014) Boekhout T. *Moniliellomycetes* and *Malasseziomycetes*, two new classes in *Ustilaginomycotina*. *Persoonia* 33: 41–47.
- Yamashita, S., Masuya, H., Abe, S., Masaki, T. & Okabe, K. (2015) Relationship between the Decomposition Process of Coarse Woody Debris and Fungal Community Structure as Detected by High-Throughput Sequencing in a Deciduous Broad-Leaved Forest in

Japan. *PLoS ONE* 10 (6): e0131510.

Yoon, J. & Kim, Y. (2005) Degradation of crystalline cellulose by the brown-rot basidiomycete *Fomitopsis palustris*. *The Journal of Microbiology* 43 (6): 487.

Zhao, Z., Liu, H., Wang, C. & Xu, J.R. (2013) Correction: Comparative analysis of fungal genomes reveals different plant cell wall degrading capacity in fungi. *BMC Genomics* 15: 6.

Zhou, S. & Ingram, L.O. (2000) Synergistic hydrolysis of carboxymethyl cellulose and acid-swollen cellulose by two endoglucanases (CelZ and CelY) from *Erwinia chrysanthemi*. *Journal of Bacteriology* 82: 5676–5682.