



Lectotypification of the name *Helianthus dentatus* Cav., basionym of *Viguiera dentata* (Cav.) Spreng. (Asteraceae: Heliantheae)

MARA ANGELINA GALVÃO MAGENTA¹, BENOÎT LOEUILLE², DAVID J. NICHOLAS HIND³
& JOSÉ RUBENS PIRANI²

¹ Departamento de Botânica, Universidade Santa Cecília, Rua Oswaldo Cruz 277, Boqueirão, 11045-90, Santos, SP, Brazil; email: maramagenta@unisanta.br

² Departamento de Botânica, Instituto de Biociências, Universidade de São Paulo, Rua do Matão 277, 05508-090, São Paulo, SP, Brazil; email: benoit_loeuille@yahoo.fr; pirani@ib.usp.br

³ Herbarium, Library, Art & Archives, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, U.K; email: n.Hind@kew.org

A recent molecular phylogenetic analysis (Schilling & Panero 2011) proposed to narrow the concept of the paraphyletic genus *Viguiera* Kunth (1818: 176; Asteraceae: Heliantheae) to embrace only the type, *V. dentata* (Cavanilles 1794: 10) Sprengel (1826: 615). Robinson (1977) emphasized the peculiarities of *Viguiera dentata* as the “distinctive and unique” presence of hairs on the filaments of the anther, the disk corollas throat less than twice as long as the lobes and densely scabrous below, a combination of features that differs from most member of the genus. During taxonomic studies of *Viguiera* and related genera, it was discovered that *V. dentata* has never been typified. The basionym, *Helianthus dentatus*, was described by Antonio José Cavanilles in *Icones et Descriptiones Plantarum* based on plants grown in the Real Jardín Botánico from seed sent by Martín Sessé from Mexico (Blanco 2000). Previous workers including Blake (1918) and Robinson (1977) have accepted the plate in Cavanilles as the type for *Helianthus dentatus* Cav.

Helianthus dentatus Cavanilles (1794: 10)

Type:—'Habitat in Imperio Mexicano cum praecedente. Florebat in Regio horto Matriense mensibus Octobri et Novembri' [Helianthus linearis Cavanilles—'Habitat in imperio Mexicano, prope oppidum Hapam'], *A. J. Cavanilles* s.n. (lectotype: MA 475778 [scan seen], designated here).≡ *Viguiera dentata* (Cav.) Sprengel. Fig. 1.

Of the four sheets of *H. dentatus* in the Cavanilles herbarium, only the sheet MA 475778 has label information annotated by Cavanilles (Garilleti 1993) and with an indication that the plant was cultivated in the Real Jardín Botánico (Fig. 1). The plate number from the *Icones* (T. 220) also appears on the label. The plate of Cavanilles does not exactly match with none of the four sheets of *H. dentatus* in the Cavanilles herbarium. But careful observations of the biggest fragment on the sheet MA 475778 allow to observe a great similarity between the plant's drawing (T. 220) and the exsicata fragment (which is in an inverted position, in relation to the plant's drawing). The label on the sheet MA 475779 was not written by Cavanilles but by J. D. Rodríguez (Blanco *et al.* 2010) and is dated 1801 (after the publication of the *Icones*). The two remaining sheets have the same number, MA 475777, one of them with two handwritten labels, but neither in Cavanilles' hand.

Viguiera dentata shows remarkable phenotypic plasticity and considerable resistance to drought and heat (Sarquis *et al.* 2010). It occurs in dry areas (slopes, canyons, fields, and roadside ditches) in the southwestern United States, Mexico, West Indies (Cuba) and Central America (Schilling 2006). More studies are necessary to determine if the four varieties currently accepted for *V. dentata* should be elevated to the species level (Blake 1918, Schilling & Panero 2011). This emphasizes the importance of selecting a type for *V. dentata*.




FIGURE 1. Lectotype of *Viguiera dentata* Cavanilles (MA 475778).

Acknowledgments

The authors would like to thank the Keeper and staff of the Real Jardín Botánico de Madrid for supplying the digital images of the syntypes of *Helianthus dentatus*.

References

- Blanco, P. (2000) Colecciones históricas: el herbario de la Real Expedición Botánica de Nueva España (1787–1803). *Boletín de la Asociación de Herbarios Ibero-Macaronésicos* 5: 5–10.
- Blanco, P., Serna, A.E., Espejo, M.A & López-Ferrari, A.R. (2010) *Catálogo del herbario de la real expedición botánica de Nueva España, 1787–1803, conservado en el Real Jardín Botánico de Madrid*. Real Jardín Botánico, CSIC, Madrid.
- Blake, S.F. (1918) *A revision of the genus Viguiera*. Contributions of the Gray Herbarium of Harvard 54: 1–205.
- Cavanilles, A.J. (1794) [1795] *Icones et descriptiones plantarum, quae aut sponte in Hispania crescunt, aut in hortis hospitantur*. Typographia regia, Madrid. Vol. 3(1).
- Garinetti, R. (1993) Herbarium cavanillesianum seu enumeratio plantarum aliquo ad novitates cavanillesianas pertinetium, quae in Horta Regii Matritensis atque Londinenses Societatis Linnaeanae herbariis assercantur. *Fontqueria* 38: 6–248.
- Kunth, C.S. (1818) In: Humboldt, A., Bonpland, A. & Kunth, C.S. (eds), *Voyage de Humboldt et Bonpland. Sixième Partie. Botanique. Nova Genera et Species Plantarum*. Ed. Folio. Maze, Paris. Vol. 4, pp. 247 & tabs 301–412.
- Robinson, H. (1977) Studies in Heliantheae (Asteraceae) VIII. Notes on genus and species limits in the genus *Viguiera*. *Phytologia* 36: 201–215.
- Sarquis, J.I., Coria, N., González-Rodríguez, H. (2010) Physiology of photosynthesis in chilmalacate (*Viguiera dentata*) in the zapotitlan de las salinas Valley of the Tehuacan biosphere reserva in puebla, Mexico. *Tropical and Subtropical Agroecosystems* 12: 361–371.
- Schilling, E.E. (2006) *Viguiera*. In: Flora of North America Editorial Committee, ed., *Flora of North America, Magnoliophyta: Asteridae (in part): Asteraceae, Vol. 21, Part 3*. Oxford University Press Inc., New York, pp 172–174.
- Schilling, E.E. & Panero, J.L. (2011) A revised classification of subtribe Helianthinae (Asteraceae: Heliantheae) II. Derived lineages. *Botanical Journal of the Linnean Society* 167: 311–331.
- Sprengel, C. (1826) *Systema vegetabilium*. Ed. 16. [sumptibus Librariae Dieterichiana] Göttingae [Göttingen]. Vol. 3.