

<https://doi.org/10.11646/phytotaxa.725.1.12>

XIN LI, XIULAN HE, WEIXIONG SHI, SHUYU ZUO, XUE LI, DI PENG, ZHAOHUI GUO & LEI SU (2025) *Penicillium sinense*, a new species of endophytic fungi from *Citrus trifoliata* with inhibition activities to pathogenic fungi. *Phytotaxa* 720 (3): 224–236.

Please note the **Results** and **Taxonomy** on page 228 and 230:

Presently TABLE 2 and FIGURE 3:

TABLE 2. Inhibition rate of strain CGMCC3.27641 on pathogenic fungi

Pathogenic fungi	Colony diameter r(cm)		Fungistatic rate (%)
	control	CGMCC3.27641	
<i>Alternaria alternata</i>	7.76±0.04b	2.54±0.18b	67.28±2.16b
<i>Fusarium solani</i>	7.98±0.04b	2.83±0.07b	64.60±1.02c
<i>Fusarium oxysporum</i> f.sp.cucumerinum	6.76±0.22c	2.20±0.11c	67.45±1.78b
<i>Colletotrichum camelliae</i>	7.71±0.12b	2.22±0.11c	71.27±1.54a
<i>Fusarium oxysporum</i>	8.43±0.15a	3.68±0.21a	56.40±2.49c

Note: Data in the table were presented as mean±SD, data with the different lowercases letters indicated significantly different ($P < 0.05$).




FIGURE 3.

Should Changed:

TABLE 2. Inhibition rate of strain CGMCC3.27641 on pathogenic fungi

Pathogenic fungi	Colony diameter r(cm)		Fungistatic rate (%)
	control	CGMCC3.27641	
<i>Alternaria alternata</i>	7.76±0.04c	2.54±0.18c	67.28±2.16b
<i>Fusarium solani</i>	7.98±0.04b	2.83±0.07b	64.60±1.02c
<i>Fusarium oxysporum</i> f.sp.cucumerinum	6.76±0.22d	2.20±0.11d	67.45±1.78b
<i>Colletotrichum camelliae</i>	7.71±0.12c	2.22±0.11d	71.27±1.54a
<i>Fusarium oxysporum</i>	8.43±0.15a	3.68±0.21a	56.40±2.49d

Note: Data in the table were presented as mean±SD, data with the different lowercases letters indicated significantly different ($P < 0.05$).




FIGURE 3.