

<https://doi.org/10.11646/phytotaxa.774.3.10>

BU-YUN ZHANG, LI LI, LIAN-SHENG XU, YU-LIN CHEN, ZHU-QIU SONG, QI-LONG YAN, CHUN-YU ZOU, GUO-DI CHEN, XIAO-YUN CHEN & YOU-SHENG CHEN (2026) Two new endangered species of *Ormosia* (Fabaceae) from Guangdong, China. *Phytotaxa* 772 (2): 121–139.

Please note the **Diagnosis** of *Ormosia yangjiangensis* Y. S. Chen, *sp. nov.* on page 127:

Presently reads:

Diagnosis. *Ormosia yangjiangensis* resembles *O. xylocarpa* and *O. pubescens* in having obtuse or acute leaflet apices and densely appressed sericeous legume valves at maturity. However, it can be clearly distinguished from *O. xylocarpa* by its conspicuously or sparsely appressed short-pubescent branchlets (vs. densely appressed yellow-brown pubescent), thinly coriaceous leaflets (vs. thickly coriaceous), axillary or pseudoterminal racemes (vs. terminal panicles or axillary racemes), a standard claw ca. 2 mm long (vs. ca. 4 mm), and legumes lacking a rib along the adaxial sutures with valves 1–3 mm thick (vs. with a prominent rib along the adaxial sutures, valves 3–7 mm thick).

Should read:

Diagnosis. *Ormosia yangjiangensis* resembles *O. xylocarpa* and *O. pubescens* in having obtuse or acute leaflet apices and densely appressed sericeous legume valves at maturity. However, it can be clearly distinguished from *O. xylocarpa* by its conspicuously or sparsely appressed short-pubescent branchlets (vs. densely appressed yellow-brown pubescent), thinly coriaceous leaflets (vs. thickly coriaceous), axillary or pseudoterminal racemes (vs. terminal panicles or axillary racemes), a standard claw ca. 2 mm long (vs. ca. 4 mm), and legumes lacking a rib along the adaxial sutures with valves 1–3 mm thick (vs. with a prominent rib along the adaxial sutures, valves 2–6 mm thick).

Please note the **Diagnosis** of *Ormosia taishanensis* Y. S. Chen, *sp. nov.* on page 131:

Presently reads:

Diagnosis. *Ormosia taishanensis* resembles *O. eugeniifolia* and *O. indurata* in having obovate, oblanceolate, or elliptic leaflets, and legumes with 3–5 ovules that become glabrescent at maturity. However, it differs from *O. eugeniifolia* by its axillary or pseudoterminal racemes (vs. terminal panicles or axillary racemes), legumes with a prominent rib along the adaxial suture (vs. without a rib), valves 6–11 mm thick (vs. 2–4 mm), and seed hilum 3–4 mm long (vs. ca. 2 mm). It differs from *O. indurata* by its thickly coriaceous leaflets (vs. thinly coriaceous), axillary or pseudoterminal racemes (vs. terminal panicles or axillary racemes), ovate or top-shaped flower buds (vs. oblong or obovate), exauriculate wings at the base (vs. 1-auriculate), distinctly turgid legumes with a prominent rib along the adaxial suture (vs. slightly turgid, without a rib), valves 6–11 mm thick (vs. 3–4 mm), and seed hilum 3–4 mm long (vs. ca. 2 mm).

Should read:

Diagnosis. *Ormosia taishanensis* resembles *O. eugeniifolia* and *O. indurata* in having obovate, oblanceolate, or elliptic leaflets, and legumes with 3–5 ovules, the legumes becoming glabrescent at maturity. However, it differs from *O. eugeniifolia* by its axillary or pseudoterminal racemes (vs. terminal panicles or axillary racemes), legumes with a prominent rib along the adaxial suture (vs. without a rib), valves 6–11 mm thick (vs. 2–4 mm), and seed hilum 3–4 mm long (vs. ca. 2 mm). It differs from *O. indurata* by its thickly coriaceous leaflets (vs. thinly coriaceous), axillary

or pseudoterminal racemes (vs. terminal panicles or axillary racemes), ovate or top-shaped flower buds (vs. oblong or obovate), exauriculate wings at the base (vs. 1-auriculate), distinctly turgid legumes with a prominent rib along the adaxial suture (vs. slightly turgid, without a rib), valves 6–11 mm thick (vs. 3–5 mm), and seed hilum 3–4 mm long (vs. ca. 2 mm).

Please note the **FIGURE 4** on page 132:

Presently reads:

FIGURE 4. *Ormosia taishanensis*. A. Habit; B. Leafy branch; C. Flowering branch; D. Fruiting branch; e. Leaves; F. Immature legumes; G. Dissected flower (G1. Calyx; G2. Standard; G3. Wings; G4. Keels; G5. Stamens and Ovary); H. Opened legumes at maturity; I. Seeds. A–C, E, G: photos by Y. S. Chen based on Voucher specimen, Y. S. Chen, C. Y. Zou, B. Y. Zhang ZL2025105 (IBSC); D, F: photos by Y. S. Chen based on Voucher specimen, Y. S. Chen & Q. L. Yan ZL2024236 (IBSC); H, I: photos by L. Li based on type specimen, Y. S. Chen, L. Li, Q. L. Yan ZL2024313 (IBSC).

Should read:

FIGURE 4. *Ormosia taishanensis*. A. Habit; B. Leafy branch; C. Flowering branch; D. Fruiting branch; E. Leaves; F. Immature legumes; G. Dissected flower (G1. Calyx; G2. Standard; G3. Wings; G4. Keels; G5. Stamens and Ovary); H. Opened legumes at maturity; I. Seeds. A–C, E, G: photos by Y. S. Chen based on Voucher specimen, Y. S. Chen, C. Y. Zou, B. Y. Zhang ZL2025105 (IBSC); D, F: photos by Y. S. Chen based on Voucher specimen, Y. S. Chen & Q. L. Yan ZL2025236 (IBSC); H, I: photos by L. Li based on type specimen, Y. S. Chen, L. Li, Q. L. Yan ZL2024313 (IBSC).

Please note the **FIGURE 7** on page 135:

Presently reads:

FIGURE 7. Morphological comparison of *Ormosia taishanensis*, *O. indurata* and *O. eugeniifolia*. A. Leaves (A1: *O. taishanensis*, A2: *O. indurata*, A3: *O. eugeniifolia*); B. Inflorescences (B1: *O. yangjiangensis*, B2: *O. indurata*); C. Immature fruit (C1: *O. taishanensis*, C2: *O. indurata*, C3: *O. eugeniifolia*). A1: photos by Y. S. Chen based on Voucher specimen, Y. S. Chen, C. Y. Zou, B. Y. Zhang, Q. C. Li ZL2025105 (IBSC); A2, C2: photos by B. Y. Zhang, based on Voucher specimen, B. Y. Zhang, L. Li, Y. L. Chen ZL2026214 (IBSC); A3: photos by B. Y. Zhang, based on Voucher specimen, Y. S. Chen, B. Y. Zhang, Q. L. Yan ZL2026100 (IBSC); B1: photos by B. Y. Zhang, based on Voucher specimen, Y. S. Chen, C. Y. Zou, B. Y. Zhang, Q. C. Li ZL2025105 (IBSC); B2: photos by S. P. Dong, based on Voucher specimen, Z. Q. Song & S. P. Dong DK2026193 (IBSC); C3: photos by J. H. Liang.

Should read:

FIGURE 7. Morphological comparison of *Ormosia taishanensis*, *O. indurata* and *O. eugeniifolia*. A. Leaves (A1: *O. taishanensis*, A2: *O. indurata*, A3: *O. eugeniifolia*); B. Inflorescences (B1: *O. taishanensis*, B2: *O. indurata*); C. Immature fruit (C1: *O. taishanensis*, C2: *O. indurata*, C3: *O. eugeniifolia*). A1: photos by Y. S. Chen based on Voucher specimen, Y. S. Chen, C. Y. Zou, B. Y. Zhang, Q. C. Li ZL2025105 (IBSC); A2, C2: photos by B. Y. Zhang, based on Voucher specimen, B. Y. Zhang, L. Li, Y. L. Chen ZL2026214 (IBSC); A3: photos by B. Y. Zhang, based on Voucher specimen, Y. S. Chen, B. Y. Zhang, Q. L. Yan ZL2026100 (IBSC); B1: photos by B. Y. Zhang, based on Voucher specimen, Y. S. Chen, C. Y. Zou, B. Y. Zhang, Q. C. Li ZL2025105 (IBSC); B2: photos by S. P. Dong, based on Voucher specimen, Z. Q. Song & S. P. Dong DK2026193 (IBSC); C1: photos by Y. S. Chen, based on Voucher specimen, Y. S. Chen, L. Li, Y. L. Chen ZL2025197 (IBSC); C3: photos by J. H. Liang.

Please note the **Additional specimens examined** (Paratypes): on page 135:

Presently reads:

Additional specimens examined (Paratypes):—China, Guangdong, Jiangmen city, Taishan county, Beifengshan forest park, alt. 279 m, riversides in evergreen forests, 23 April 2025, *You-Sheng CHEN, Chun-Yu ZOU, Bu-Yun ZHANG ZL2025105* (IBSC); Jiangmen city, Taishan county, Chixi Town, Luobokeng, alt. 31 m, riversides, 21 April 2025, *You-Sheng CHEN, Chun-Yu ZOU, Bu-Yun ZHANG ZL2025082* (IBSC); China, Guangdong, Jiangmen city, Taishan county, Beifengshan forest park, alt. 279 m, riversides in evergreen forests, 7 July 2025, *You-Sheng CHEN, Li LI, Yu-Lin CHEN ZL2024197* (IBSC); Jiangmen city, Taishan county, Beifengshan forest park, alt. 279 m, riversides in evergreen forests, 27 September 2025, *You-Sheng CHEN, Qi-Long YAN ZL2024236* (IBSC); China, Guangdong, Jiangmen city, Taishan county, Gudoushan mountain, alt. 527 m, 23 March 2024, *Shuang-Wen DENG, Mei LU, Peng-Huan WANG DSW3430* (IBSC).

Should read:

Additional specimens examined (Paratypes):—China, Guangdong, Jiangmen city, Taishan county, Beifengshan forest park, alt. 279 m, riversides in evergreen forests, 23 April 2025, *You-Sheng CHEN, Chun-Yu ZOU, Bu-Yun ZHANG ZL2025105* (IBSC); Jiangmen city, Taishan county, Chixi Town, Luobokeng, alt. 31 m, riversides, 21 April 2025, *You-Sheng CHEN, Chun-Yu ZOU, Bu-Yun ZHANG ZL2025082* (IBSC); China, Guangdong, Jiangmen city, Taishan county, Beifengshan forest park, alt. 279 m, riversides in evergreen forests, 7 July 2025, *You-Sheng CHEN, Li LI, Yu-Lin CHEN ZL2025197* (IBSC); Jiangmen city, Taishan county, Beifengshan forest park, alt. 279 m, riversides in evergreen forests, 27 September 2025, *You-Sheng CHEN, Qi-Long YAN ZL2025236* (IBSC); China, Guangdong, Jiangmen city, Taishan county, Gudoushan mountain, alt. 527 m, 23 March 2024, *Shuang-Wen DENG, Mei LU, Peng-Huan WANG DSW3430* (IBSC).

Please note the **TABLE 2.** on page 125:

Presently reads:

Character	<i>O. yangjiangensis</i>	<i>O. taishanensis</i>	<i>Ormosia pubescens</i>	<i>O. xylocarpa</i>	<i>O. indurata</i>	<i>O. eugeniifolia</i>
Branchlets	conspicuously or sparsely appressed short pubescent	densely appressed pubescent	densely appressed shortly pubescent	densely appressed-pubescent	densely appressed-pubescent	densely appressed pubescent
Petiolule	(2–)3–5 mm	(3–)4–9 mm	(3–)4–8(–10) mm	(3–)6–13(–14) mm	3–6 mm	3–6 mm

Should read:

Character	<i>O. yangjiangensis</i>	<i>O. taishanensis</i>	<i>Ormosia pubescens</i>	<i>O. xylocarpa</i>	<i>O. indurata</i>	<i>O. eugeniifolia</i>
Branchlets	conspicuously or sparsely appressed short pubescent	densely appressed pubescent	densely appressed shortly pubescent	densely appressed-pubescent	densely appressed-pubescent	densely appressed pubescent
Petiolule	(2–)3–5 mm	2–7 mm	(3–)4–8(–10) mm	(3–)6–13(–14) mm	3–6 mm	3–6 mm