



Notes on Early Land Plants Today. 30. Transfer of some taxa from *Anastrophyllum* (Anastrophyllaceae, Marchantiophyta)

JIŘÍ VÁŇA¹, LARS SÖDERSTRÖM², ANDERS HAGBORG³ & MATT VON KONRAT³

¹Department of Botany, Charles University, Benátská 2, CZ-12801 Praha 2, Czech Republic; vana@natur.cuni.cz

²Department of Biology, Norwegian University of Science and Technology, N-7491 Trondheim, Norway; lars.soderstrom@bio.ntnu.no

³Department of Botany, The Field Museum, 1400 South Lake Shore Drive, Chicago, IL 60605–2496, USA; hagborg@pobox.com, mvonkonrat@fieldmuseum.org

The genus *Anastrophyllum* Stephani (1893: 139) was treated in a broad sense by e.g. Schuster (1969) and has usually been so until recently. De Roo *et al.* (2007) showed that the genus is polyphyletic and removed *Sphenolobus* (Lindberg 1874: 369) Berggren (1898: 22) and *Crossocalyx* Meylan (1939: 266). Subsequently Váňa & Engel (2013) removed subgenus *Zantenia* Hattori (1966: 342). However, two other elements remain, that also deserve to be removed.

Anastrophyllum involutifolium and at least also *Anastrophyllum subcomplicatum* (supposedly also *Anastrophyllum revolutum*) belong to a group of taxa which should be separated from the genus *Anastrophyllum* on generic level. Both taxa are characterized by the very concave (hemispherically concave, not canaliculately concave) leaves with slightly interlocking dorsal merophytes, dorsally falcate-secund leaves, broadly overlapping dorsally and extending across the width of stem and decurrent, etc.

Schuster (2002) designated the species *Anastrophyllum schismoides* (treated as synonym of *Anastrophyllopsis subcomplicata* here) as the type of his new section *Anastrophyllum* sect. *Anastrophyllopsis*. However, he also included in this section the taxa *Anastrophyllum piligerum* (Reinwardt *et al.* 1824: 414) Stephani (1893: 140) and *Anastrophyllum appendiculatum* Kitagagawa (1970: 210), “whose sectional disposition remains uncertain” (Schuster 2002: 313). At least *Anastrophyllum appendiculatum* (accepted by JV as conspecific with *Anastrophyllum auritum* (Lehmann 1829: 368) Stephani (1901: 1137)) does not belong to this group.

The necessary new combinations are made here.

Formal treatment

The format of this note follows what is outlined in Söderström *et al.* (2012).

***Anastrophyllopsis* (R.M.Schust.) Váňa et L.Söderstr., comb. et stat. nov.**

Basionym:—*Anastrophyllum* sect. *Anastrophyllopsis* R.M.Schust., *Beih. Nova Hedwigia* 119: 310, 2002 (Schuster 2002).

Type:—*Anastrophyllum schismoides* (Mont.) Steph. [= *Anastrophyllum subcomplicatum* (Lehm. et Lindenb.) Steph.].

***Anastrophyllopsis involutifolia* (Mont. ex Gottsche, Lindenb. et Nees) Váňa et L.Söderstr. comb. nov.**

Basionym:—*Jungermannia involutifolia* Mont. ex Gottsche, Lindenb. et Nees, *Syn. Hepat.* 1: 81, 1844 (Gottsche *et al.* 1844).

Lectotype (Váňa & Engel 2013):—CHILE. Prov. Magallanes: Bahía San Nicolas, Hombron, (PC!, isolectotypes BM!, S!).

≡ *Anastrophyllum involutifolium* (Mont. ex Gottsche, Lindenb. et Nees) Steph., *Hedwigia* 32: 140, 1893 (Stephani 1893).

***Anastrophyllopsis revoluta* (Steph.) Váňa et L.Söderstr. comb. nov.**

Basionym:—*Anastrophyllopsis revolutum* Steph., *Hedwigia* 32: 139, 1893.

Type:—PAPUA NEW GUINEA. Mt. Suckling, 1891, *Wm. McGregor* (holotype G-67269! [=G-11073!]¹, isotype BM!).

***Anastrophyllopsis subcomplicata* (Lehm. et Lindenb.) Váňa et L.Söderstr. comb. nov.**

Basionym:—*Jungermannia subcomplicata* Lehm. et Lindenb., *Nov. Stirp. Pug.* 7: 4, 1838 (Lehmann 1838).

Lectotype (Váňa & Engel 2013):—MARIANA IS. Habitat in insulis Marianis. Specimina benevole communicavit Dr. Bongard, Professsor Petropolitanus celeberrimus (W-Lindenb. Hep. 1597!, isolectotype S!).

≡ *Anastrophyllopsis subcomplicatum* (Lehm. et Lindenb.) Steph., *Hedwigia* 32: 140, 1893 (Stephani 1893).

= *Anastrophyllopsis schismoides* (Mont.) Steph., *Hedwigia* 32: 140, 1893 (Stephani 1893), syn. in Váňa & Engel (2013).

Basionym: *Jungermannia schismoides* Mont., *Ann. Sci. Nat. Bot. (ser. 2)* 19: 250, 1843 (Montagne 1843).

Type:—NEW ZEALAND. Auckland I., 1840, *Hooker*, Voy. of H.M.S. Erebus & Terror (isotypes BM!, FH!, G! S!, W-Lindenb. Hep. 1624!, 1625!).

= *Anastrophyllopsis obtusatum* (Hook.f. et Taylor) Steph., *Bull. Herb. Boissier ser. 2*, 1: 1136 (*Spec. Hep.* 2: 119), 1901 (Stephani 1901), syn. in Gottsche *et al.* (1845). Basionym:—*Jungermannia obtusata* Hook.f. et Taylor, *London J. Bot.* 4: 80, 1845 (Hooker & Taylor 1845). Type:—ST. HELENA. 1844, *Hooker* (isotype FH!).

= *Anastrophyllopsis tristanianum* J.J.Engel, *Fieldiana, Bot.* 38: 71, 1977 (Engel 1977), syn. in Váňa (1982). Type:—TRISTAN DA CUNHA. Above Burntwood, 650 m, *Christophersen & Mejland* 889 (holotype O!).

= *Anastrophyllopsis schismoides* var. *crassulum* J.J.Engel, *J. Bryol.* 20: 379, 1998 (Engel & Braggins 1998), *syn. nov.* Holotype:—AUSTRALIA. Tasmania: Gordon River, vicinity of Sir John Falls, just up river from Butler Island, ca. 50 m, Engel 14742 (holotype F, isotype HO).

***Schizophyllum* (R.M.Schust.) Váňa et L.Söderstr. comb. et stat. nov.**

Basionym:—*Anastrophyllopsis* subgen. *Schizophyllum* R.M.Schust., *Hepat. Anthocerotae N. Amer.* 2: 739, 1969 (Schuster 1969).

Type:—*Anastrophyllopsis sphenoloboides* R.M.Schust.

***Schizophyllum aristatum* (Herzog ex N.Kitag.) Váňa et L.Söderstr. comb. nov.**

Basionym:—*Anastrophyllopsis bidens* var. *aristatum* Herzog ex N.Kitag., *J. Hattori Bot. Lab.* 33: 216, 1970 (Kitagawa 1970).

Type:—BORNEO. Sarawak: Dulit Ridge, 1300 m, on tree-trunk, *P. W. Richards* 1908 (holotype JE!).

≡ *Anastrophyllopsis aristatum* (Herzog ex N.Kitag.) A.E.D.Daniels, D.G.Long, Karyappa et P.Daniel., *J. Bryol.* 34: 148, 2012 (Daniels *et al.* 2012).

***Schizophyllum bidens* (Reinw., Blume et Nees) Váňa et L.Söderstr. comb. nov.**

Basionym:—*Jungermannia bidens* Reinw., Blume et Nees, *Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur.* 12: 208, 1825 [1824] (Reinwardt *et al.* 1824).

Type:—INDONESIA. Java (holotype STR, isotype W-Lindenb. Hep. 1618!).

***Schizophyllum lancilobum* (Steph.) Váňa et L.Söderstr. comb. nov.**

Basionym:—*Anastrophyllopsis lancilobum* Steph., *Sp. Hepat. (Stephani)* 6: 107, 1917 (Stephani 1917).

Type:—TAHITI. *Temarii*, hb Levier 3161b. (holotype G-10890!, isotypes FH!, FI!).

Note:—This taxon is close to *A. bidens*, but the type specimen is poor and difficult to place.

***Schizophyllum papillosum* (J.J.Engel et Braggins) Váňa et L.Söderstr. comb. nov.**

Basionym:—*Anastrophyllopsis papillosum* J.J.Engel et Braggins, *J. Bryol.* 20: 381, 1998 (Engel & Braggins 1998).

1. Citation of specimens in G should preferably use the barcode (M. Price, pers. comm.) but for comparability the numbers printed on the specimen, which have often been cited by previous authors, are also given here in square brackets.

Type:—NEW ZEALAND. North Is.: Wellington Prov., Tongariro Natl. Park, Soda Sings, Mangatepopo Stream, N. E. of Whakapapa Village, 1350 m, Braggins 97/106B (holotype F, isotypes AK, CHR-558817).

***Schizophyllum sphenoloboides* (R.M.Schust.) Váňa et L.Söderstr. comb. nov.**

Basionym:—*Anastrophylloides sphenoloboides* R.M.Schust., *Hepat. Anthocerotae N. Amer.* 2: 741, 1969 (Schuster 1969).
Type:—NW. GREENLAND. Kangerlugssuak, Inglefield Bay, RMS 45984 (F-1079638, isotypes C!, TNS!).

Acknowledgement

We thank Lucia Kawasaki (F) for helping us searching for R.M.Schuster types. The Early Land Plants Today project (ELPT) has been generously supported in part by the Global Biological Information Facility (GBIF) Seed Money Award No.2007-41, activities facilitated in part by funding from the Biodiversity Synthesis Center of the Encyclopedia of Life (BioSynC), partial funding from the National Science Foundation (Award No's 0749762, 1115002), the Warwick Foundation, and the Negaunee Foundation.

References

- Berggren, S. (1898) *On New Zealand Hepaticae. I.* E. Malmström, Lund, 48 pp.
- Daniels, A.E.D., Long, D.G., Kariyappa, K.C. & Daniel, P. (2012) *Anastrophylloides aristatum* (Herzog ex N.Kitag.) A.E.D. Daniels et al., comb. et stat. nov. (Marchantiophyta: Anastrophylaceae) from India. *Journal of Bryology* 34: 146–149. <http://dx.doi.org/10.1179/1743282012Y.0000000002>
- Engel, J.J. (1977) Austral Hepaticae IX. *Anastrophylloides tristanianum*, a new species from Tristan da Cunha. *Fieldiana, Botany* 38: 71–74.
- Engel, J.J. & Braggins, J.E. (1998) Austral Hepaticae. 27. The genus *Anastrophylloides* (Spruce) Steph. (Jungermanniales) in Australasia, with a synopsis of austral taxa. *Journal of Bryology* 20: 371–388.
- Gottschke, C.M., Lindenberg, J.B.G. & Nees von Esenbeck, S.G. (1844) *Synopsis Hepaticarum, fasc. 1.* Meissner, Hamburg, pp. 1–144.
- Gottschke, C.M., Lindenberg, J.B.G. & Nees von Esenbeck, S.G. (1845) *Synopsis Hepaticarum, fasc. 2.* Meissner, Hamburg, pp. 145–304.
- Grolle, R. (1968) Einige ostmalesische Lebermoose. *Nova Hedwigia* 16: 147–159.
- Hattori, S. (1966) A remarkable *Anastrophylloides* (Hepaticae) from New Guinea. *Botanical Magazine [Shokubutsu-gaku zasshi]*, Tokyo 79: 342–344.
- Herzog, T. (1950) Hepaticae borneenses (Oxford University expeditions to Sarawak, 1932). *Transactions of the British Bryological Society* 1: 275–326.
- Hooker, J.D. & Taylor, T. (1845) Hepaticae Antarcticae, supplementum. *London Journal of Botany* 4: 79–97.
- Kitagawa, N. (1970) Lophoziaeae of North Borneo. *Journal of the Hattori Botanical Laboratory* 33: 203–221.
- Lehmann, J.G.C. (1829) Hepaticarum Capensium a C.F. Ecklon collectarum brevem recensionem cum Schlechtendalio suo. *Linnaea* 4: 357–371.
- Lehmann, J.G.C. (1838) *Novarum et minus cognitarum stirpium pugillus VII addita enumeratione plantarum omnium in his pugillis descriptarum.* Meissner, Hamburg, pp. 1–41.
- Lindberg, S.O. (1874) Manipulus muscorum secundus. *Notiser ur Sällskapets pro Fauna et Flora Fennica Förhandlingar* 13: 353–417.
- Meylan, C. (1939) Localités nouvelles pour la flore des muscinées de la Suisse. *Bulletin de la société vaudoise de sciences naturelles* 60: 261–276.
- Montagne, J.F.C. (1843) Quatrième centurie de plantes cellulaires exotiques nouvelles, décades I–VI. *Annales des Sciences Naturelles, Botanique, ser. 2* 19: 238–266.
- De Roo, R.T., Hedderson, T.A. & Söderström, L. (2007) Molecular insights into the phylogeny of the leafy liverwort family Lophoziaeae Cavers. *Taxon* 56: 301–314.
- Reinwardt, C.G.C., Blume, C.L. & Nees von Esenbeck, C.G. (1824) Hepaticae Iavanicae, editae coniunctis studiis et opera. *Nova Acta Physico-medica Academiae Caesareae Leopoldino-Carolinae Naturae Curiosorum* 12 “1825”: 181–238.
- Schiffner, V. (1893) Ueber exotische Hepaticae, hauptsächlich aus Java, Amboina und Brasilien, nebst einigen morphologischen und kritischen Bemerkungen über *Marchantia*. *Nova Acta Academiae Caesareae Leopoldino-*

- Carolinae Germanicae Naturae Curiosorum* 60: 219–316.
- Schuster, R.M. (1969) *The Hepaticae and Anthocerotae of North America. vol. II*. Columbia University Press, New York, 1062 pp.
- Schuster, R.M. (1978) Studies on Venezuelan Hepaticae. I. *Phytologia* 39: 239–251.
- Schuster, R.M. (2002) Austral Hepaticae, part II. *Beihefte zur Nova Hedwigia* 119: 1–606.
- Söderström, L. Hagborg, A. & von Konrat, M. (2012) Notes on Early Land Plants Today. *Phytotaxa* 65: 41–42.
- Spruce, R. (1876) On *Anomoclada*, a new genus of Hepaticae, and on its allied genera, *Odontoschisma* and *Adelanthus*. *Journal of Botany* 14: 230–235.
- Stephani, F. (1893) Hepaticarum species novae. Pars II. *Hedwigia* 32: 137–147.
- Stephani, F. (1901) Species Hepaticarum 2. *Bulletin de l'Herbier Boissier, série 2* 1: 1022–1140.
- Stephani, F. (1917) *Species Hepaticarum* 6. George & Cie, Genève & Bale, pp. 1–128.
- Váňa, J. (1982) Notes on some African hepatic genera 1–5. *Folia Geobotanica et Phytotaxonomica* 17: 63–87.
- Váňa, J. & Engel, J.J. (2013). The liverworts and hornworts of the Tristan da Cunha group. *Memoirs of the New York Botanical Garden* 105: 1–138.